

International Journal of Nutrition and Metabolism

Volume 4 Number 4 April 2012

ISSN 2141-2340



*Academic
Journals*

ABOUT IJNAM

The **International Journal of Nutrition and Metabolism (IJNAM)** is published monthly (one volume per year) by Academic Journals.

International Journal of Nutrition and Metabolism (IJNAM) is an open access journal that provides rapid publication (monthly) of articles in all areas of the subject such as Thermic effect of food, Anthropogenic metabolism, calorimetry, flavonoids etc.

Submission of Manuscript

Submit manuscripts as e-mail attachment to the Editorial Office at: ijnam@acadjournals.org. A manuscript number will be mailed to the corresponding author shortly after submission.

The International Journal of Nutrition and Metabolism will only accept manuscripts submitted as e-mail attachments.

Please read the **Instructions for Authors** before submitting your manuscript. The manuscript files should be given the last name of the first author.

Editors

Dr. Mohamed Fawzy Ramadan Hassanien,
*Biochemistry Department,
Faculty of Agriculture,
Zagazig University,
Egypt.*

Dr. Ahmed Mohamed El-Waziry,
*Alexandria University,
Faculty of Agriculture,
Dept. of Animal Production,
Egypt.*

Prof. Bechan Sharma,
*Visiting Professor of Biochemistry,
Christopher S. Bond Life Science Center,
Department of Microbiology and Immunology,
University of Missouri-Columbia,
1210 Rollins Street,
Columbia 65201,
USA.*

Prof. Malay Chatterjee,
*Jadavpur University, Kolkata,
India.*

Dr. Wei Wang,
*School of Public Health and Family Medicine,
Capital Medical University,
China.*

Dr. Kedar Nath Mohanta,
*ICAR Research Complex for Goa,
Goa.*

Dr. Birinchi Kumar Sarma,
*Banaras Hindu University,
Varanasi,
India.*

Editorial Board

Prof. Alonzo A. Gabriel

*University of the Philippines,
Diliman, Quezon City
Philippines.*

Dr. Michael Elliott

*Washington University in St. Louis,
USA.*

Prof. Satyesh Chandra Roy,

*University of Calcutta,
India.*

Dr. Hena Yasmin

*University of Swaziland,
Swaziland.*

Dr. Neveen B. Talaat

*Department of Plant Physiology,
Faculty of Agriculture,
Cairo University,
Egypt.*

Dr. V.Sivajothi

*karpagam college of pharmacy
othakkalmandapam, coimbatore,
Tamilnadu,
India.*

Dr. M. Manjoro Nee Mwale,

*University of Fort Hare,
South Africa.*

Dr. Adewumi, Gbenga Adedeji,

*University Of Lagos, Akoka,
Lagos,
Nigeria.*

Dr. Iheanyi O. Okonko,

*University of Ibadan,
Ibadan,
Nigeria.*

Dr. Ashok Kumar Tiwari,

*Indian Institute of Chemical Technology,
India.*

Dr. Mukund Adsul,

*National Chemical Laboratory, Pune,
India.*

Dr. Fengdi Ji,

*Beijing Institute of Food & Brewing,
China.*

Dr. Charles Tortoe,

*CSIR-Food Research Institute,
Ghana.*

Dr. Mridula Devi,

*Food Grains and Oilseeds Processing Division,
Central Institute of Post Harvest Engineering and
Technology (CIPHET),
Ludhiana-141 004, (Punjab),
India.*

Dr. Faiyaz Ahmed,

*DOS in Food Science and Nutrition,
University of Mysore,
India.*

Dr. Samie A,

*University of Venda,
South Africa.*

Dr. Giampaolo Papi,

*Department of Internal Medicine,
Azienda USL Modena,
Italy.*

Ahmad Taher Azar,

*Institution Modern Science and Arts University (MSA),
6th of October City,
Egypt.*

Dr. T. Poongodi Vijayakumar,

*Department of Food Science,
Periyar University,
Salem, Tamil Nadu,
India.*

Dr. Radhakrishnan Ramaraj,

*University of Arizona,
Cedars Sinai Hospital 1501 N Campbell Avenue Tucson, AZ
85724,
United States.*

Dr. Chaman Farzana,

*Mount Carmel college, Bangalore,
India.*

Dr. Hesham Mahyoub Al-Mekhlafi,

*University of Malaya,
Malaysia.*

Dr. Amal Ahmed Ali Abdul-Aziz,

*National Research Center,
Textile Devisision,
Egypt.*

Instructions for Author

Electronic submission of manuscripts is strongly encouraged, provided that the text, tables, and figures are included in a single Microsoft Word file (preferably in Arial font).

The **cover letter** should include the corresponding author's full address and telephone/fax numbers and should be in an e-mail message sent to the Editor, with the file, whose name should begin with the first author's surname, as an attachment.

Article Types

Three types of manuscripts may be submitted:

Regular articles: These should describe new and carefully confirmed findings, and experimental procedures should be given in sufficient detail for others to verify the work. The length of a full paper should be the minimum required to describe and interpret the work clearly.

Short Communications: A Short Communication is suitable for recording the results of complete small investigations or giving details of new models or hypotheses, innovative methods, techniques or apparatus. The style of main sections need not conform to that of full-length papers. Short communications are 2 to 4 printed pages (about 6 to 12 manuscript pages) in length.

Reviews: Submissions of reviews and perspectives covering topics of current interest are welcome and encouraged. Reviews should be concise and no longer than 4-6 printed pages (about 12 to 18 manuscript pages). Reviews are also peer-reviewed.

Review Process

All manuscripts are reviewed by an editor and members of the Editorial Board or qualified outside reviewers. Authors cannot nominate reviewers. Only reviewers randomly selected from our database with specialization in the subject area will be contacted to evaluate the manuscripts. The process will be blind review.

Decisions will be made as rapidly as possible, and the journal strives to return reviewers' comments to authors as fast as possible. The editorial board will re-review manuscripts that are accepted pending revision. It is the goal of the IJNAM to publish manuscripts within weeks after submission.

Regular articles

All portions of the manuscript must be typed double-spaced and all pages numbered starting from the title page.

The Title should be a brief phrase describing the contents of the paper. The Title Page should include the authors' full names and affiliations, the name of the corresponding author along with phone, fax and E-mail information. Present addresses of authors should appear as a footnote.

The Abstract should be informative and completely self-explanatory, briefly present the topic, state the scope of the experiments, indicate significant data, and point out major findings and conclusions. The Abstract should be 100 to 200 words in length.. Complete sentences, active verbs, and the third person should be used, and the abstract should be written in the past tense. Standard nomenclature should be used and abbreviations should be avoided. No literature should be cited. Following the abstract, about 3 to 10 key words that will provide indexing references should be listed.

A list of non-standard **Abbreviations** should be added. In general, non-standard abbreviations should be used only when the full term is very long and used often. Each abbreviation should be spelled out and introduced in parentheses the first time it is used in the text. Only recommended SI units should be used. Authors should use the solidus presentation (mg/ml). Standard abbreviations (such as ATP and DNA) need not be defined.

The Introduction should provide a clear statement of the problem, the relevant literature on the subject, and the proposed approach or solution. It should be understandable to colleagues from a broad range of scientific disciplines.

Materials and methods should be complete enough to allow experiments to be reproduced. However, only truly new procedures should be described in detail; previously published procedures should be cited, and important modifications of published procedures should be mentioned briefly. Capitalize trade names and include the manufacturer's name and address. Subheadings should be used. Methods in general use need not be described in detail.

Results should be presented with clarity and precision. The results should be written in the past tense when describing findings in the authors' experiments. Previously published findings should be written in the present tense. Results should be explained, but largely without referring to the literature. Discussion, speculation and detailed interpretation of data should not be included in the Results but should be put into the Discussion section.

The Discussion should interpret the findings in view of the results obtained in this and in past studies on this topic. State the conclusions in a few sentences at the end of the paper. The Results and Discussion sections can include subheadings, and when appropriate, both sections can be combined.

The Acknowledgments of people, grants, funds, etc should be brief.

Tables should be kept to a minimum and be designed to be as simple as possible. Tables are to be typed double-spaced throughout, including headings and footnotes. Each table should be on a separate page, numbered consecutively in Arabic numerals and supplied with a heading and a legend. Tables should be self-explanatory without reference to the text. The details of the methods used in the experiments should preferably be described in the legend instead of in the text. The same data should not be presented in both table and graph form or repeated in the text.

Figure legends should be typed in numerical order on a separate sheet. Graphics should be prepared using applications capable of generating high resolution GIF, TIFF, JPEG or Powerpoint before pasting in the Microsoft Word manuscript file. Tables should be prepared in Microsoft Word. Use Arabic numerals to designate figures and upper case letters for their parts (Figure 1). Begin each legend with a title and include sufficient description so that the figure is understandable without reading the text of the manuscript. Information given in legends should not be repeated in the text.

References: In the text, a reference identified by means of an author's name should be followed by the date of the reference in parentheses. When there are more than two authors, only the first author's name should be mentioned, followed by 'et al'. In the event that an author cited has had two or more works published during the same year, the reference, both in the text and in the reference list, should be identified by a lower case letter like 'a' and 'b' after the date to distinguish the works.

Examples:

Abayomi (2000), Agindotan et al. (2003), (Kelebeni, 1983), (Usman and Smith, 1992), (Chege, 1998;

1987a,b; Tijani, 1993,1995), (Kumasi et al., 2001) References should be listed at the end of the paper in alphabetical order. Articles in preparation or articles submitted for publication, unpublished observations, personal communications, etc. should not be included in the reference list but should only be mentioned in the article text (e.g., A. Kingori, University of Nairobi, Kenya, personal communication). Journal names are abbreviated according to Chemical Abstracts. Authors are fully responsible for the accuracy of the references.

Examples:

Chikere CB, Omoni VT and Chikere BO (2008). Distribution of potential nosocomial pathogens in a hospital environment. *Afr. J. Biotechnol.* 7: 3535-3539.

Moran GJ, Amii RN, Abrahamian FM, Talan DA (2005). Methicillinresistant *Staphylococcus aureus* in community-acquired skin infections. *Emerg. Infect. Dis.* 11: 928-930.

Pitout JDD, Church DL, Gregson DB, Chow BL, McCracken M, Mulvey M, Laupland KB (2007). Molecular epidemiology of CTXM-producing *Escherichia coli* in the Calgary Health Region: emergence of CTX-M-15-producing isolates. *Antimicrob. Agents Chemother.* 51: 1281-1286.

Pelczar JR, Harley JP, Klein DA (1993). *Microbiology: Concepts and Applications*. McGraw-Hill Inc., New York, pp. 591-603.

Short Communications

Short Communications are limited to a maximum of two figures and one table. They should present a complete study that is more limited in scope than is found in full-length papers. The items of manuscript preparation listed above apply to Short Communications with the following differences: (1) Abstracts are limited to 100 words; (2) instead of a separate Materials and Methods section, experimental procedures may be incorporated into Figure Legends and Table footnotes; (3) Results and Discussion should be combined into a single section.

Fees and Charges: Authors are required to pay a \$550 handling fee. Publication of an article in the International Journal of Nutrition and Metabolism is not contingent upon the author's ability to pay the charges. Neither is acceptance to pay the handling fee a guarantee that the paper will be accepted for publication. Authors may still request (in advance) that the editorial office waive some of the handling fee under special circumstances.

Copyright: © 2012, Academic Journals.

All rights Reserved. In accessing this journal, you agree that you will access the contents for your own personal use but not for any commercial use. Any use and or copies of this Journal in whole or in part must include the customary bibliographic citation, including author attribution, date and article title.

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract or as part of a published lecture, or thesis) that it is not under consideration for publication elsewhere; that if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher.

Disclaimer of Warranties

In no event shall Academic Journals be liable for any special, incidental, indirect, or consequential damages of any kind arising out of or in connection with the use of the articles or other material derived from the IJNAM, whether or not advised of the possibility of damage, and on any theory of liability.

This publication is provided "as is" without warranty of any kind, either expressed or implied, including, but not limited to, the implied warranties of merchantability, fitness for a particular purpose, or non-infringement. Descriptions of, or references to, products or publications does not imply endorsement of that product or publication. While every effort is made by Academic Journals to see that no inaccurate or misleading data, opinion or statements appear in this publication, they wish to make it clear that the data and opinions appearing in the articles and advertisements herein are the responsibility of the contributor or advertiser concerned. Academic Journals makes no warranty of any kind, either express or implied, regarding the quality, accuracy, availability, or validity of the data or information in this publication or of any other publication to which it may be linked.

ARTICLES

Research Article

Proximate composition and breeding description of an unidentified Cichlid of Epe Lagoon, Lagos, Southwest, Nigeria commonly called 'Wesafu'

57

H. Fashina-Bombata and I. Megbowon

Full Length Research Paper

Proximate composition and breeding description of an unidentified Cichlid of Epe Lagoon, Lagos, Southwest, Nigeria commonly called 'Wesafu'

H. Fashina-Bombata^{1*} and I. Megbowon²

¹Department of Fisheries, Faculty of Science, Lagos State University, Lagos, Nigeria.

²Biotechnology Section, Nigerian Institute for Oceanography and Marine Research, Victoria Island, P. M. B. 12729 Lagos, Nigeria.

Accepted 1 March, 2012

A common cichlid of the Epe Lagoon, Lagos, Nigeria is an important food fish and viable candidate for culture in meeting the millennium goal and vision 2020 in the subregion. The intense fisheries of this ecotype also pose a conservation issue. The comparative protein contents of the cichlid fishes were 21.80, 18.34, 19.13, 18.71 and 18.08% for Wesafu, *Sarotherodon galileus*, *Sarotherodon melanotheron*, *Tilapia guineensis* and *Tilapia mariae*, respectively. The highest percentage of protein was obtained for Wesafu. The results of the lipid determination showed that Wesafu had 0.9% lipid which is higher than others with *T. guineensis* having the least (0.4%) value for lipid. The result showed that all the fish studied belong to low oil category and therefore can be classified as lean fish. The major component of fish muscle is moisture. There was a progressive increase in the number of ripe eggs (absolute fecundity) from larger fish to smaller one. However, the relative fecundity followed a reverse order. Female Wesafu produce only a few hundreds to less than 2,000 eggs per spawn but inter-spawning period varied from 25 to 30 days in the present study. It was observed that this parameter (spawning frequency) was not affected by the weight of the broodstock. Although Wesafu could be seen as a commercially important fish species based on its credentials of deeper body, high economic value, in terms of price, white flesh and good taste, its aquaculture potential may be affected by its low fecundity. On the courtship behavior, Wesafu exhibits a characteristic golden yellow radiation during breeding period. During courtship, the fish display a characteristics kissing lifestyle. This appears different from that of *Oreochromis niloticus* which displays a poking courtship behavior with the female swimming in front while the male pokes the side. The result of this finding showed that of all the brooders in all hapas, only females that had eggs/hatchlings in their mouthparts. This comes to show that Wesafu probably belong to the genus, *Oreochromis*. Although this study has identified Wesafu as a probable member of the genus, *Oreochromis*, there is the need to identify the fish at a molecular level to see if it belongs to any of the known species of *Oreochromis*, or perhaps a simple strain.

Key words: Ecotype cichlid, Wesafu, *Tilapiine* spp., breeding behavior, proximate composition.

INTRODUCTION

Throughout the world, aquaculture is seen as a panacea for meeting the increasing demand for fish, as catches

from open waters are declining due to over exploitation and degradation of fish habitats. Fish production from inland open waters declined due to over exploitation, intensive agriculture, industrial development, erosion and siltation, reclamation of land for human settlement, pollution, destruction of mangrove forests, etc. Fish farming therefore appears to hold the key for improved

*Corresponding author. E-mail: bombatta2002@gmail.com.
Tel: +2348021186098.

global fish production.

Among the numerous species of fish for culture, tilapia is widely recognized as one of the popular species for a wide range of aquaculture systems worldwide. It is an ideal candidate for warm water aquaculture (Tahoun et al., 2008). Tilapia generally differs greatly in size and taxonomic group (Olojo et al., 2003). Worldwide, Nile tilapia (*Oreochromis niloticus*) culture increased during the year, 2001 to 2006 from 1,113,737 to 1,988,726 MT representing a growth of 79% (FAO, 2008), thus making it one of the fastest growing freshwater aquaculture species.

Tilapia spawn easily in captivity, use a wide range of natural foods as well as formulated diets, tolerate poor water quality and grows well at warm temperatures. These attributes, along with relatively low input costs, have made tilapia the most widely cultured freshwater fish in tropical and subtropical countries (Borgeson et al., 2006; El-Saidy and Gaber, 2005; Fasakin et al., 2005). Although the potential for tilapia culture is high, the production in Africa and more importantly, Nigeria is very low; the draw-back being the early maturity, uncontrolled reproduction in ponds leading to increased competition for food, reduction in growth rate which results in a phenomenon referred to as stunting (Fashina-Bombata et al., 2006, Baroiller and Toguyemi, 1996). There are generally low productions of fry of *O. niloticus* which is attributable to low fecundity, inadequate sex ratio, spawning techniques, broodstock nutrition and high fry mortality (Tahoun et al., 2008). Reproductive success in many species has been shown to be influenced by the broodstock, sex ratio, stocking density, age, size, nutrition and feeding regime. Broodstock productivity clearly represents the most significant constraint to commercial tilapia production in Nigeria.

Although *O. niloticus* is the leading species of tilapia cultured globally, there exists an unidentified cichlid 'Wesafu', in Epe lagoon, Lagos, Nigeria where it is highly priced. This unidentified cichlid grows to 1,500 g, 414 mm in the wild. At present, there is paucity of information on the origin of the species of fish.

A number of studies have been conducted on the morphometric and meristic characteristics, age and growth, food and feeding habit, nutritional requirement, amino acid profile and biochemical characterization of the fish (Fashina-Bombata and Hammed, 2010; Hammed et al., 2010, 2011, Fashina-Bombata et al., 2005a, 2005b, 2006, 2008). These studies showed some unique characteristics in this cichlid. However, the proximate composition, breeding characteristic and DNA studies have not been addressed and could provide some basic information and clue as to the identity of this fish. This study therefore focused on the proximate composition, breeding characterization and placing Wesafu in the appropriate genera; RAPD technique was employed to show the genetic diversity and relatedness with other cichlid in Epe lagoon, Nigeria.

MATERIALS AND METHODS

Sampling

Fresh specimens of Tilapiine fishes were purchased at Ebute Chief landing site, Epe lagoon, Lagos State. The samples were collected aseptically and transported in an insulated container under chilled condition to the Fish Technology Laboratory, Nigerian Institute for Oceanography and Marine Research Victoria Island, for analysis.

Proximate composition

Proximate analyses were carried out in triplicate on the samples after dressing. These analyses included moisture content, total lipid, crude protein and ash.

Moisture content

The moisture content was determined by drying the sample to constant mass using the oven-dry method (AOAC, 1994). At first, the initial weight of the samples was taken. Then samples were dried in the oven (Memmet 854 Schwabach) at about $105 \pm 2^\circ\text{C}$ for about 8 to 10 h until a constant weight was reached and cooled in a desiccator and weighed again. Then the samples were minced in an electric grinder. The percentage of moisture content was calculated by the following equation:

Percentage (%) of moisture = (weight loss/original weight of sample) \times 100.

Determination of fat content

Dried samples were finely ground and the fat was extracted with a non-polar solvent, ethyl ether using the modified Bligh and Dyer procedure. After extraction, the solvent was evaporated and the extracted materials weighed. The percentage of fat content was calculated as:

Percentage (%) of fat = (weight of extract/weight of sample) \times 100.

Determination of protein

Protein content of the fish was determined by micro-Kjeldahl method. It involves conversion of organic nitrogen to ammonium sulphate by digestion with concentrated sulphuric acid in a micro-Kjeldahl flask. The digest was diluted, made alkaline with sodium hydroxide and distilled. The liberated ammonia was collected in a boric acid solution and was determined titrimetrically. The percentage of protein in the sample was calculated by the following equation:

Percentage (%) of protein = (c-b) \times 14 \times d \times 6.25/a \times 1000 \times 100

where, a = Sample weight (g); b = volume of NaOH required for back titration and neutralize 25 ml of 0.1 N H_2SO_4 (for sample); c = volume of NaOH required for back titration and neutralize 25 ml of 0.1 N H_2SO_4 (for blank); d = normality of NaOH used for titration; 6.25 = conversion factor of N to protein while 14 = atomic weight of N.

Determination of ash

The ash content was determined by igniting the samples at about




Figure 1. Moisture content of 5 species of tilapia fishes from Epe Lagoon, Lagos.

550 to 600°C for 5 to 6 h until the sample were completely free of carbon particles (residue become white) in a Carbolite Sheffield LMF3 muffle furnace. The percent of ash was calculated as follows:

Percentage (%) of ash = (weight of ash / weight of sample) × 100

Statistical analysis

Data were analyzed by using SPSS 12 statistical programme with five percent level of significance.

Breeding characterization

The breeding study involved single pair mating of gravid broodstock (143.1 to 310.6 g) in triplicate batches in 1 m × 1 m × 1 m hapas installed in 0.1 ha fish pond in a private fish farm in Badagry. Female broodstocks were divided into 4 groups based on weight: Less than 150 g (A), 150 to 200 g (B), 200 to 250 g (C) and 250 g and above (D). The hapas were covered with net (2 cm mesh size) to prevent fish from jumping out. Fish were daily fed 3 to 5% body weight with Coppens^(c) feed, imported from Netherland and the mouth of both sexes observed regularly for the presence of eggs/hatchlings and the condition of the genital papilla which may still be swollen even few days after the female has released eggs. The courtship behavior was observed from time to time and documented. Individual weight of fish was recorded every 2 weeks. The sex of fish with eggs/hatchlings in the mouth was noted and recorded for each spawn. The number of eggs fertilized were also counted and recorded. The inter-spawning period (number of days) were also observed and recorded for each hapa. The following reproductive parameters were determined according to (Mair et al., 2004):

- i. Absolute fecundity: The number of seed per spawning per female
- ii. Relative fecundity: The number of seed per unit weight of female
- iii. Inter-spawning period: Number of days between successive spawning.

RESULTS

Proximate analysis

Figures 1 to 4 show the results of proximate composition of sample of cichlid fishes collected from Epe lagoon. The major component of fish muscle was moisture (Figure 2). The moisture content ranged between 78.50 and 81.70% with *Tilapia guineensis* having the highest value and Wesafu the least. The values were 79.33, 80.35, 81.70, 80.83 and 78.50% for *Sarotherodon galileaus*, *Sarotherodon galileaus*, *T. guineensis*, *Tilapia mariae* and Wesafu, respectively.

The result of lipid content showed a range of 0.4 to 0.9% with Wesafu having the highest lipid content (0.9%) and *T. guineensis* the least (0.4%). Others are *S. galileaus* (0.65%), *S. melanotheron* (0.82%) and *T. mariae* (0.6%) (Figure 3).

The ash content of the sample were 1.75, 1.75, 1.00, 1.30 and 1.20% for *S. galileaus*, *S. melanotheron*, *T. guineensis*, *T. mariae* and Wesafu, respectively (Figure 4). The values of crude protein were 18.34, 19.13, 18.71, 18.08 and 21.80% for *S. galileaus*, *S. melanotheron*, *T. guineensis*, *T. mariae* and Wesafu, respectively. The result showed Wesafu to have the highest crude protein value (Figure 5).

Breeding characterization

The result of this study shows a progressive increase in absolute fecundity with corresponding increase in fish weight. However, the relative fecundity decreased with increase in the fish weight (Table 1). It was observed that




Figure 2. Lipid content (%) of 5 species of tilapia fishes obtained from Epe Lagoon, Lagos.




Figure 3. Ash content of 5 species of tilapia fishes obtained from Epe lagoon, Lagos.




Figure 4. Crude protein content of 5 tilapia fishes obtained from Epe lagoon, Lagos.




Figure 5. Crude protein content of 5 tilapiine fishes obtained from Epe lagoon, Lagos.

Table 1. Effect of fish weight on absolute and relative fecundity of an ecotype cichlid ‘Wesafu’ bred in hapa.

Treatment	Wt. of female broodstock (g)	Wt. of male broodstock (g)	Absolute fecundity	Relative fecundity	No. of fertilized eggs	Fertilization (%)	Av. no. of swim - up fry/ g of female
A	150.0±8.9	150.0±6.1	951±18.7	6.63±1.9	876±9.9	91.2	6.04±1.7
B	190.0±12.7	180.0±15.7	1042±14.7	5.8±1.8	894±9.7	88.3	4.67±1.2
C	240.0±11.6	225.0±14.9	1078±18.5	5.05±1.1	921±9.5	83.2	4.14±0.7
D	290.0±16.3	290.0±14.2	1165±19.7	3.98±1.2	1003±11.7	85.5	3.42±1.1

Table 2. Inter-spawning period and brooding sex of the cichlid.

Treatment	Wt. of female broodstock (g)	Wt. of male broodstock (g)	Range of Inter-spawning period (days)	Brooding sex
A	150.0±8.9	150.0±6.1	25-28	F
B	190.0±12.7	180.0±15.7	25-29	F
C	240.0±11.6	225.0±14.9	26-29	F
D	290.0±16.3	290.0±14.2	25-29	F

the average number of fry per gram weight of female decreased with increase in female body weight. The results further show that percentage fertilization was highest in Group A than B, C and D with slight numeric differences. Furthermore, there appeared to be no significant variation in the inter-spawning period among the four groups. The range was between 25 and 30 days, (Table 2). It is worthy of note that in all the single pair mating carried out; the brooding sex was female, suggesting that the unidentified cichlid is most probably, a maternal mouth brooder. It comes to confirm the fish as a member of the genera, *Oreochromis*.

DISCUSSION

The major component of fish muscle is moisture. The moisture content of the fishes studied varied between

78.80 and 80.83%. The moisture content for fish sample was within reasonable range as reported by Gallagher et al. (1991). FAO (1999) reported that moisture and lipid content in fish muscle were inversely related and their sum is approximately 80% with other components accounting for the remaining 20% which partially agreed with the finding of the present study. In summary, the finding of this present study reveals the nutritional significance of *Wesafu* among other cichlids of Epe Lagoon.

The protein contents of the cichlid fishes were 21.80, 18.34, 19.13, 18.71 and 18.08% for *Wesafu*, *S. galileaus*, *S. melanotheron*, *T. guineensis* and *T. mariae*, respectively. The values varied considerably from species to species. This is in agreement with Alkobaby et al. (2008) who reported that the chemical composition of fish muscle varies greatly from species to species even among individuals within the same species. According to

the author, such variation depends on age, size, sex, environment and season. Sallam et al. (2007) further stressed that the variation is closely related to feed intake, migratory, swimming and sexual changes associated with spawning. The highest percentage of protein was obtained for Wesafu in this study, which is probably due to the presence of certain amino acids in Wesafu which may be absent in other cichlids studied. Hammed et al. (2010) reported that Wesafu has the highest value of phenylalanine, isoleucine, leucine and valine than other cichlid studies from the lagoon.

The results of the lipid determination showed that Wesafu had 0.9% lipid which is higher than others with *T. guineensis* having the least (0.4%) value for lipid. The result showed that all the fish studied belong to low oil category and therefore can be classified as lean fish. Stansby (1982) and Ackman (1989) reported that fishes with lipid content below 5% are lean fishes.

There was a progressive increase in the number of ripe eggs (absolute fecundity) from larger fish to smaller one. However, the relative fecundity followed a reverse order. This may be attributed to the age of the brooders. Smith et al. (1991) reported that older two year old red tilapia female produced more seed per clutch but fewer seed per unit weight than younger, one year old females. The author further established that despite a large average clutch size, seed production was lower in older 2 year old female due to lowering spawning frequency, suggesting fewer reproductively active individuals of longer inter-spawning periods. Furthermore, Little and Hulata (2000) reported that, larger, older fish can perform well in intensive hatchery systems, although frequency of spawning and clutch size decline with size and age. El-Sayed and Gaber (2005) reported that small Nile Tilapia broodfish produced many more eggs, with shorter inter-spawning period than large fish. It was also found that large eggs contained more yolk and led to larger fry with better growth and higher resistance to starvation and severe environmental conditions. One may therefore ask whether it is better to use small fish, which could collectively yield more eggs per culture unit, or larger fish with higher individual egg production but longer intervals. Poor handling efficiency of large fish is a problem in hapa and tank-based hatchery systems where broodstocks are individually handled to harvest mouthbrooded seed. In practice, fish larger than 300 g are difficult to handle quickly and efficiently, more importantly for a highly jumping fish like Wesafu capable of jumping over 80 cm above the flood plane in hapa. Extra care is required to prevent fish escape.

Female Wesafu produce only a few hundreds to less than 2,000 eggs per spawn but inter-spawning period varied from 25 to 30 days in the present study. It was observed that this parameter (spawning frequency) was not affected by the weight of the broodstock. This is in agreement with the report of Popma and Lovshin (1996) which stated that, under appropriate environmental

conditions, tilapias spawn frequently (every 4 to 6 weeks) and at young age (usually less than 6 months) but overall fecundity is low. Although Wesafu could be seen as a commercially important fish species based on its credentials of deeper body, high economic value, in terms of price, white flesh and good taste (Fashina-Bombata et al., 2005b), its aquaculture potential may be affected by its low fecundity (Fashina-Bombata and Somotun, 2008). This is not a desirable aquacultural trait because greater number of female brood stock is required to sustain a commercial aquaculture production. On the positive side, however, the need for many broodstock decreases the risk of inbreeding depression, an important management problem in the farming of more fecund fish species.

On the courtship behavior, Wesafu exhibits a characteristic golden yellow radiation during breeding period. Smaller fish have a more or less silvery colouration. During courtship, the fish display a characteristics kissing lifestyle which the opposite sex quickly understands. During the period, the male and female bring their opened mouth close to touch each other and then separate. This appears different from that of *O. niloticus* which displays a poking courtship behavior with the female swimming in front while the male pokes the side. The different courtship behavior of the two species has thwarted hybridization effort for now but still remains an option for tilapia programme development in Nigeria.

This study showed that only females had eggs/hatchlings in their mouthparts. This comes to show that Wesafu probably belong to the genus, *Oreochromis*. Popma and Lovshin (1996) reported that all maternal mouth brooders are classified as *Oreochromis* since 1983. Although this study has identified Wesafu as a probable member of the genus, *Oreochromis*, there is still the need to identify at a molecular level to see if it belongs to any of the known species of *Oreochromis* or simply a variant of any of them. This opens a window of opportunity for researchers in tilapia genetics, for which collaboration is sought.

REFERENCES

- Ackman R (1989). Nutritional composition of fats in Seafoods. *Progressive Nutr. Sci.*, 13: 161-241.
- Alkobaby AI, Sami, AS, Ghada, IA (2008). Effect of protein sources on characteristics and quality traits of Nile Tilapia (*O. niloticus*). 8th International Symposium on Tilapia Aquaculture, Egypt, 12-14, 1: 551-567.
- Association of Official Methods of Analytical Chemists (AOAC). (1994). Official methods of Analysis of the Association of Official Chemists. Association of Analytical chemist, Airlington, 1(11): 129
- Baroiller JF, Toguyemi A (1996). Comparative effects of natural androgen 11 α hydro androstenedione and a synthetic androgen, 17 α -methyltestosterone on the sex ratio of *O. niloticus* 238-245. In Pullin, R.S.V., Lazard L., Legendre, J., Amon Konthias, J.B. and Pauly, D. (eds.). The third International Symposium on Tilapia in Aquaculture. ICLARM Conference proceedings, 41: 575.
- Borgeson TL, Racz V, Wilkie DC, White LJ, Drew MD (2006). Effect of

- replacing fish meal and oil with simple or complex mixtures of vegetable ingredients in diets fed to Nile tilapia (*Oreochromis niloticus*). *Aquacult. Nutr.*, 12: 141-149
- EI-Sayed DMS, Gaber MMA (2005). Effect of dietary protein levels and feeding rates on growth performance, production traits and body composition of Nile tilapia, *O. niloticus* in concrete tanks. *Aquacult. Res.*, 36 (2): 163-171.
- FAO (1999). World Production of Crustaceans and Molluscs by major fishing areas. Fisheries Information Data and Statistics Unit (FIDI). Fisheries Department, FAO, Rome, pp. 33.
- FAO (2008). FAO Fisheries Department, Fisheries Information, Data and Statistics unit, pp. 214-218.
- Fasakin EA, Serwata RD, Davies SJ (2005). Comparative utilization of rendered animal derived products with or without composite mixture of soya bean meal in hybrid tilapia (*O. niloticus* X *O. mossambicus*). *Aquaculture*, 249 (1-4): 329-338.
- Fashina-Bombata HA, Ajepe GR, Hammed AM (2008). Age and Growth Pattern of the Endemic Cichlid of Epe Lagoon Commonly Called 'Wesafu'. *Glob. J. Agric. Sci.*, 7(1): 105-109
- Fashina-Bombata HA, Ajepe RG, Hammed AM (2006) Food and feeding habits of "Wesafu" An Ecotype Cichlid of Epe-Lagoon, Nigeria. *World Aquacult.*, 37:1 63-66.
- Fashina-Bombata HA, Hammed AM (2010). Determination of Nutrient Requirements of an Ecotype Cichlid of Epe Lagoon, Southwest Nigeria. *Glob. J. Agric. Sci.*, 9(2): 57-61
- Fashina-Bombata HA, Somotun AO (2008). The Effect of Lyophilized Goat Testes Meal as First Feed on the Growth of 'Wesafu': An Ecotype Cichlid of Epe-Lagoon, in Lagos State, Nigeria. *Pak. J. Nutr.*, 7 (5): 686-688
- Fashina-Bombata HA, Ajepe RG, Hammed AM, Jimoh AA (2005a). Characterization of an ecotype cichlid commonly referred to as Wesafu, endemic to Epe lagoon, Nigeria. *World Aquaculture*, 36: 20-22.
- Fashina-Bombata HA, Hammed AM, Ajepe RG, Jimoh AA (2005b). Comparative Organoleptic Assessment of "Wesafu" (an ecotype Cichlid of Epe Lagoon), *Oreochromis niloticus* and *Tilapia mariae*. *Proceedings of the 4th Faculty of Science Annual Conference*, 28-29
- Gallagher ML, Harrell ML, Rurilson RA (1991). Variation in Lipid and fatty acid contents of Atlantic croakers, Striped mullet and Summer Flounder. *Trans. Amer. Fish.*, 614-619.
- Hammed AM, Fashina-Bombata HA, Fajana OO (2010) Tissue and Blood Amino Acids Composition of an Ecotype Cichlid 'Wesafu', *Tilapia zilli* and *Oreochromis niloticus* Using Paper Chromatography. *Pak. J. Nutr.*, 9 (7):724-727
- Hammed AM, Fashina-Bombata HA, Omitogun GO (2011). Biochemical characterization of an Ecotype Cichlid from Epe Lagoon 'Wesafu' and *Oreochromis niloticus* using Sodium dodecyl Sulphate Polyacrylamide gel electrophoresis (SDS- PAGE). *Res. Biotechnol.*, 2(5): 56-62
- Little DC, Hulata G (2000). Strategies for tilapia seed production . P267-326, In: Beveridge M.C.M and B.J McAndrew (Editors). *Tilapia: Biology and Exploitation*. Kluwer Acad. Publ., Dordrecht, The Netherlands.
- Mair CG, Lakapunrat S, Jere WL, Bart A (2004). Comparison of Reproductive Parameters Among Improved Strains of Nile Tilapia *Oreochromis niloticus* L. In: Bolivar, R. B.; Mair, G. C. and Fitzsimmons, K. (Editors). *New dimensions in farmed tilapia*. Proc. 6th Inter. Sympos. Tilapia in aquaculture, Roxas Boulevard, Manila, Philippines, pp.12-16.
- Olojo EA, Olurin KB, Osikoya OJ (2003). Food and feeding habit of *Cynodontis nigrita* from Osun river, S W. Nig. *World fish centre Quarterly (NAGA)*, 26(4): 21-24.
- Popma TJ, Lovshin LL (1996). World wide prospects for commercial production of tilapia. Research and Development, series no. 41. International centre for Aquaculture and Aquatic environments, Department of Fisheries and Allied Aquaculture, Auburn University, Alabama, pp. 23.
- Sallam KI, Ahmed AM, Elgazzar MM, Eldaly EA (2007). Chemical quality and sensory attributes of marinated saury (*Colobalis saira*).during vacuum packaged storage at 4°C. *Food chem.*, 102: 1061-1070.
- Smith SJ, Watanabe WO , Chan JR, Ernst DH, Wicklund RI, Olla BL (1991). Hatchery production of Florida red tilapia seed in brackishwater tanks: the influence of broodstock age. *Aquacult. Fish. Manage.*, 22 (2): 141-147.
- Stansby ME (1982). Properties of fish oil and their application to handling of fish. *Ayi Publishing Co. Westport C.T.*
- Tahoun AM, Ibrahim MAR, Hammounda YF, Eid MS, Zaki-Eldin MMA, Magouz FI (2008). Effect of age and stocking density on spawning performance of Nile tilapia (*Oreochromis niloticus*) broodstock reared in hapas. (ISTA VIII), Egypt, *Proceed. Intl. symposium Tilapia Aquaculture*, pp. 329-344.

UPCOMING CONFERENCES

**16th International Congress on Renal Nutrition and Metabolism
(ICRNM) Honolulu, USA, 26 Jun 2012**



***Academy of Nutrition and Dietetics Food & Nutrition Conference & Expo,
Philadelphia, USA, 6 Oct 2012***



Conferences and Advert

September 2012

30th Annual Scientific Meeting of The Obesity Society, San Antonio, USA, 20 Sep 2012

October 2012

Academy of Nutrition and Dietetics Food & Nutrition Conference & Expo, Philadelphia, USA, 6 Oct 2012

International Journal of Nutrition and Metabolism

Related Journals Published by Academic Journals

- *Clinical Reviews and Opinions*
- *Journal of Medicinal Plant Research*
- *African Journal of Pharmacy and Pharmacology*
- *Journal of Dentistry and Oral Hygiene*
- *Journal of Parasitology and Vector Biology*
- *Journal of Pharmacognosy and Phytotherapy*

academicJournals