

# Journal of Bacteriology Research

Volume 6 Number 2 April 2014

ISSN 2006-9871



*Academic  
Journals*

## ABOUT JBR

The **Journal of Bacteriology Research (JBR)** (ISSN 2006-9871) is published Monthly (one volume per year) by Academic Journals.

**Journal of Bacteriology Research (JBR)** is a peer reviewed journal. The journal is published per article and covers all areas of the subject such as: Bacterial physiology, Bacterial flora for human, Prokaryotes of the environment, Bacterial endotoxin, Cell signalling.

## Submission of Manuscript

Please read the **Instructions for Authors** before submitting your manuscript. The manuscript files should be given the last name of the first author

[Click here to Submit manuscripts online](#)

If you have any difficulty using the online submission system, kindly submit via this email [jbr@academicjournals.org](mailto:jbr@academicjournals.org).

With questions or concerns, please contact the Editorial Office at [jbr@academicjournals.org](mailto:jbr@academicjournals.org).

## Editor

**Dr. Colleen Olive**

*Queensland Institute of Medical Research  
PO Royal Brisbane Hospital  
Brisbane,  
Australia.*

**Dr. Lyuba Doumanova**

*The Stephan Angeloff Institute of Microbiology  
Bulgarian Academy of Sciences  
Sofia,  
Bulgaria.*

**Dr. Imtiaz Wani**

*S.M.H.S Hospital  
Amira Kadal,  
India.*

**Dr. Aamir Shahzad**

*Max F. Perutz Laboratories  
University of Vienna  
Vienna,  
Austria.*

**Dr. Ömür Baysal**

*West Mediterranean Agricultural Research  
Institute (BATEM)  
Antalya,  
Turkey.*

## Associate Editors

**Dr. Chang-Gu Hyun**

*Jeju Biodiversity Research Institute (JBRI) and Jeju Hi-Tech Industry Development Institute (HiDI)  
Jeju,  
Korea.*

**Dr. Ramasamy Harikrishnan**

*Jeju National University  
Department of Aquatic Life Medicine  
College of Ocean Science  
Korea.*

**Prof. Salah M. Azwai**

*Al Fateh University  
Tripoli,  
Libya.*

**Dr. Osman Radwan**

*University of Illinois  
Urbana, IL  
USA.*

**Prof. Abdulghani Alsamarai**

*Tikrit University College of Medicine  
Tikrit,  
Iraq.*

**Dr. Nuno Cerca**

*University of Minho  
Braga,  
Portugal.*

**Dr. Mohammad Reza Shakibaie**

*Department of Microbiology and Immunology  
Kerman University of Medical Sciences  
Kerman,  
Iran.*

## Editorial Board

**Dr. Bojarajan Senthilkumar**

*Institute for Ocean Management  
Anna University  
Chennai,  
India.*

**Dr. Gurdeep Rastogi**

*Department of Plant Pathology  
University of California  
Davis, CA  
USA.*

**Dr. Asis Khan**

*Washington University  
St. Louis, MO  
USA.*

**Saikat Kumar Basu**

*University of Lethbridge  
Lethbridge, AB  
Canada.*

**Dr. Sivaramaiah Nallapeta**

*ONAN Centre for Molecular Investigations  
Secunderabad,  
India.*

**Dr. Yajnavalka Banerjee**

*Max-Planck Institute for Biophysical Chemistry  
Goettingen,  
Germany.*

**Dr. Petr Slama**

*Mendel University of Agriculture and Forestry  
Department of Animal Morphology, Physiology  
and Genetics  
Brno,  
Czech Republic.*

**Dr. Petros V. Vlastarakos**

*Lister Hospital  
Stevenage,  
UK.*

**Dr. Lee Seong Wei**

*Department Fishery Science and Aquaculture  
Faculty Agrotechnology and Food Science  
Universiti Malaysia Terengganu  
Terengganu,  
Malaysia.*

# Instructions for Author

**Electronic submission** of manuscripts is strongly encouraged, provided that the text, tables, and figures are included in a single Microsoft Word file (preferably in Arial font).

The **cover letter** should include the corresponding author's full address and telephone/fax numbers and should be in an e-mail message sent to the Editor, with the file, whose name should begin with the first author's surname, as an attachment.

## Article Types

Three types of manuscripts may be submitted:

**Regular articles:** These should describe new and carefully confirmed findings, and experimental procedures should be given in sufficient detail for others to verify the work. The length of a full paper should be the minimum required to describe and interpret the work clearly.

**Short Communications:** A Short Communication is suitable for recording the results of complete small investigations or giving details of new models or hypotheses, innovative methods, techniques or apparatus. The style of main sections need not conform to that of full-length papers. Short communications are 2 to 4 printed pages (about 6 to 12 manuscript pages) in length.

**Reviews:** Submissions of reviews and perspectives covering topics of current interest are welcome and encouraged. Reviews should be concise and no longer than 4-6 printed pages (about 12 to 18 manuscript pages). Reviews are also peer-reviewed.

## Review Process

All manuscripts are reviewed by an editor and members of the Editorial Board or qualified outside reviewers. Authors cannot nominate reviewers. Only reviewers randomly selected from our database with specialization in the subject area will be contacted to evaluate the manuscripts. The process will be blind review.

Decisions will be made as rapidly as possible, and the journal strives to return reviewers' comments to authors as fast as possible. The editorial board will re-review manuscripts that are accepted pending revision. It is the goal of the AJFS to publish manuscripts within weeks after submission.

## Regular articles

All portions of the manuscript must be typed double-spaced and all pages numbered starting from the title page.

**The Title** should be a brief phrase describing the contents of the paper. The Title Page should include the authors' full names and affiliations, the name of the corresponding author along with phone, fax and E-mail information. Present addresses of authors should appear as a footnote.

**The Abstract** should be informative and completely self-explanatory, briefly present the topic, state the scope of the experiments, indicate significant data, and point out major findings and conclusions. The Abstract should be 100 to 200 words in length. Complete sentences, active verbs, and the third person should be used, and the abstract should be written in the past tense. Standard nomenclature should be used and abbreviations should be avoided. No literature should be cited.

Following the abstract, about 3 to 10 key words that will provide indexing references should be listed.

A list of non-standard **Abbreviations** should be added. In general, non-standard abbreviations should be used only when the full term is very long and used often. Each abbreviation should be spelled out and introduced in parentheses the first time it is used in the text. Only recommended SI units should be used. Authors should use the solidus presentation (mg/ml). Standard abbreviations (such as ATP and DNA) need not be defined.

**The Introduction** should provide a clear statement of the problem, the relevant literature on the subject, and the proposed approach or solution. It should be understandable to colleagues from a broad range of scientific disciplines.

**Materials and methods** should be complete enough to allow experiments to be reproduced. However, only truly new procedures should be described in detail; previously published procedures should be cited, and important modifications of published procedures should be mentioned briefly. Capitalize trade names and include the manufacturer's name and address. Subheadings should be used. Methods in general use need not be described in detail.

**Results** should be presented with clarity and precision. The results should be written in the past tense when describing findings in the authors' experiments. Previously published findings should be written in the present tense. Results should be explained, but largely without referring to the literature. Discussion, speculation and detailed interpretation of data should not be included in the Results but should be put into the Discussion section.

**The Discussion** should interpret the findings in view of the results obtained in this and in past studies on this topic. State the conclusions in a few sentences at the end of the paper. The Results and Discussion sections can include subheadings, and when appropriate, both sections can be combined.

**The Acknowledgments** of people, grants, funds, etc should be brief.

**Tables** should be kept to a minimum and be designed to be as simple as possible. Tables are to be typed double-spaced throughout, including headings and footnotes. Each table should be on a separate page, numbered consecutively in Arabic numerals and supplied with a heading and a legend. Tables should be self-explanatory without reference to the text. The details of the methods used in the experiments should preferably be described in the legend instead of in the text. The same data should not be presented in both table and graph form or repeated in the text.

**Figure legends** should be typed in numerical order on a separate sheet. Graphics should be prepared using applications capable of generating high resolution GIF, TIFF, JPEG or Powerpoint before pasting in the Microsoft Word manuscript file. Tables should be prepared in Microsoft Word. Use Arabic numerals to designate figures and upper case letters for their parts (Figure 1). Begin each legend with a title and include sufficient description so that the figure is understandable without reading the text of the manuscript. Information given in legends should not be repeated in the text.

**References:** In the text, a reference identified by means of an author's name should be followed by the date of the reference in parentheses. When there are more than two authors, only the first author's name should be mentioned, followed by 'et al'. In the event that an author cited has had two or more works published during the same year, the reference, both in the text and in the reference list, should be identified by a lower case letter like 'a' and 'b' after the date to distinguish the works.

Examples:

Abayomi (2000), Agindotan et al. (2003), (Kelebeni, 1983), (Usman and Smith, 1992), (Chege, 1998;

1987a,b; Tijani, 1993,1995), (Kumasi et al., 2001) References should be listed at the end of the paper in alphabetical order. Articles in preparation or articles submitted for publication, unpublished observations, personal communications, etc. should not be included in the reference list but should only be mentioned in the article text (e.g., A. Kingori, University of Nairobi, Kenya, personal communication). Journal names are abbreviated according to Chemical Abstracts. Authors are fully responsible for the accuracy of the references.

Examples:

Chikere CB, Omoni VT and Chikere BO (2008). Distribution of potential nosocomial pathogens in a hospital environment. *Afr. J. Biotechnol.* 7: 3535-3539.

Moran GJ, Amii RN, Abrahamian FM, Talan DA (2005). Methicillinresistant *Staphylococcus aureus* in community-acquired skin infections. *Emerg. Infect. Dis.* 11: 928-930.

Pitout JDD, Church DL, Gregson DB, Chow BL, McCracken M, Mulvey M, Laupland KB (2007). Molecular epidemiology of CTXM-producing *Escherichia coli* in the Calgary Health Region: emergence of CTX-M-15-producing isolates. *Antimicrob. Agents Chemother.* 51: 1281-1286.

Pelczar JR, Harley JP, Klein DA (1993). *Microbiology: Concepts and Applications.* McGraw-Hill Inc., New York, pp. 591-603.

### Short Communications

Short Communications are limited to a maximum of two figures and one table. They should present a complete study that is more limited in scope than is found in full-length papers. The items of manuscript preparation listed above apply to Short Communications with the following differences: (1) Abstracts are limited to 100 words; (2) instead of a separate Materials and Methods section, experimental procedures may be incorporated into Figure Legends and Table footnotes; (3) Results and Discussion should be combined into a single section.

Proofs and Reprints: Electronic proofs will be sent (e-mail attachment) to the corresponding author as a PDF file. Page proofs are considered to be the final version of the manuscript. With the exception of typographical or minor clerical errors, no changes will be made in the manuscript at the proof stage.

**Fees and Charges:** Authors are required to pay a \$550 handling fee. Publication of an article in the Journal of Bacteriology Research is not contingent upon the author's ability to pay the charges. Neither is acceptance to pay the handling fee a guarantee that the paper will be accepted for publication. Authors may still request (in advance) that the editorial office waive some of the handling fee under special circumstances

**Copyright: © 2014, Academic Journals.**

All rights Reserved. In accessing this journal, you agree that you will access the contents for your own personal use but not for any commercial use. Any use and or copies of this Journal in whole or in part must include the customary bibliographic citation, including author attribution, date and article title.

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract or as part of a published lecture, or thesis) that it is not under consideration for publication elsewhere; that if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher.

**Disclaimer of Warranties**

In no event shall Academic Journals be liable for any special, incidental, indirect, or consequential damages of any kind arising out of or in connection with the use of the articles or other material derived from the JBR, whether or not advised of the possibility of damage, and on any theory of liability.

This publication is provided "as is" without warranty of any kind, either expressed or implied, including, but not limited to, the implied warranties of merchantability, fitness for a particular purpose, or non-infringement. Descriptions of, or references to, products or publications does not imply endorsement of that product or publication. While every effort is made by Academic Journals to see that no inaccurate or misleading data, opinion or statements appear in this publication, they wish to make it clear that the data and opinions appearing in the articles and advertisements herein are the responsibility of the contributor or advertiser concerned. Academic Journals makes no warranty of any kind, either express or implied, regarding the quality, accuracy, availability, or validity of the data or information in this publication or of any other publication to which it may be linked.

**ARTICLES**

**Incidence and antibiotic susceptibility pattern of *Vibrio* species isolated from sea foods sold in Port-Harcourt, Nigeria**

Chijioke A. Nsofor, Samuel T. Kemajou and Chibuzor M. Nsofor



## Full Length Research Paper

# Incidence and antibiotic susceptibility pattern of *Vibrio* species isolated from sea foods sold in Port-Harcourt, Nigeria

Chijioke A. Nsofor<sup>1\*</sup>, Samuel T. Kemajou<sup>2</sup> and Chibuzor M. Nsofor<sup>2</sup>

<sup>1</sup>Department of Biotechnology, Federal University of Technology, Owerri, Imo State, Nigeria.

<sup>2</sup>Department of Medical Laboratory Science, Madonna University, Elele, Rivers State, Nigeria.

Accepted 6 February, 2014

Most *Vibrio* infections are associated with the consumption of raw or undercooked sea foods or exposure of wounds to warm seawater. In this study, 63 samples of a variety of sea foods viz: shrimps, periwinkle, prawn, crayfish collected from two major household markets in Port-Harcourt were cultivated using standard bacteriological method on Thiosulphate Citrate Bile Sucrose (TCBS) agar. A total of 63 *Vibrios* belonging to seven different species were isolated with *Vibrio fluvialis* recorded as the highest percentage frequency occurrence and the most predominant species, 30 (47.6%), followed by *Vibrio paraheamolyticus* with 19 (30.2%), *Vibrio vulnificus* with 4 (6.34%), *Vibrio metschnikovi* with 4 (6.34%), *Photobacterium* spp. with 3 (4.76%), *Vibrio cholerae* with 2 (3.17%) and the least was *Vibrio mimicus* with 1 (1.60%). Statistically, there was a significant different ( $P < 0.05$ ) in prawn sold in Mile 1 market as compared to prawn sold in Mile 3 markets. These isolates were subjected to their susceptibility patterns using agar diffusion method, which recorded susceptible to most antibiotics used. The presence of these pathogenic strains of *Vibrios* in commonly consumed sea foods is of public concern.

**Key word:** Vibrios, sea foods, Nigeria.

## INTRODUCTION

*Vibrio* species are Gram negative, facultative anaerobic motile asporogenous rod or curved rod-shaped bacteria with a single polar flagellum. The genus contains at least twelve species pathogenic to human, eight of which can cause or are associated with food-borne illness (Dickinson et al., 2013). The majority of the food-borne illness is caused by *Vibrio paraheamolyticus*, *Vibrio vulnificus*, *Vibrio fluvialis* and *Vibrio cholerae*. *V. vulnificus* is responsible for 95% of sea food related deaths while immune suppressed individuals are most

susceptible to other *Vibrio* infection (Miyoshi, 2013). Marine *Vibrios* are ubiquitous in the marine environment; therefore, it is not surprising that many are pathogenic for various seafood hosts which are harvested for human consumption (Kaysner and DePaola, 2004).

Most countries have guidelines for detecting *V. paraheamolyticus*, *V. cholera* O<sub>1</sub> and O<sub>139</sub> in seafood, whereas few have guidelines for *V. vulnificus*. Thus, there are routine microbiological analysis of seafood includes testing for *V. paraheamolyticus*, *V. cholera* O<sub>1</sub> and O<sub>139</sub>,

\*Corresponding author. E-mail: nsoforac@gmail.com. Tel: +2348038783955.

**Table 1.** Mean *Vibrio* species from sea foods bought in Mile 1 and Mile 3 markets in Port Harcourt represented as log<sub>10</sub> cfu/ml.

Sample	Mile 1	Mile 3
Periwinkle	67.0±44.2	20.4±14.5
Shrimp	52.0±21.6	21.0±11.1
Prawn	24.4±13.9	49.4±41.6
Crayfish	15.0±9.8	17.7±11.5

but seldom for *V. vulnificus* (Park et al., 2013).

Some species are primarily associated with gastrointestinal illness (*V. cholerae* and *V. parahaemolyticus*) while others can cause non-intestinal illness, such as septicemia (*V. vulnificus*). In tropical and temperate regions, disease-causing species of *Vibrio* occur naturally in marine, coastal and estuarine environments and are most abundant in estuarine. Pathogenic *Vibrios*, in particular *V. cholerae*, can also be recovered from freshwater reaches of estuaries (Lutz et al., 2013). The occurrence of these bacteria does not generally correlate with numbers of fecal coli forms and depurations of shellfish may not reduce their numbers. However, a positive correlation between fecal contamination and levels of *V. cholerae* may be found in areas experiencing cholera outbreaks (Amita et al., 2003).

Cholera epidemics outbreaks have killed millions of people and continue to be the major public health concern worldwide (Islam et al., 2013). Fluid and electrolytes replacements are the major treatment of cholera patients, however, patients with severe disease require antibiotic treatment to reduce the duration of illness and reduce replacement fluid intake. Like most bacteria of clinical and public health significance, *V. cholerae* is continuously becoming resistance to a variety of antimicrobial agents, necessitating use of newer drugs which are more expensive and have more adverse effect. Spread of cholera epidemics worldwide has been associated with the emergence of multiple drug resistance among a large number of *V. cholera* strains (Dunstan et al., 2013).

Sea foods especially shellfish is a substrate for some zoonotic *Vibrios* of which these microorganisms, cause food poisoning and diarrhea in human. Sea foods are prone to bacterial contamination and could cause health risk to human consumers (Lutz et al., 2013). *Vibrios* are associated with live sea foods as they form part of the indigenous micro flora of the marine environment. *V. parahaemolyticus* is the top causative agent among all the reported seafood poisoning outbreaks in Nigeria (Eyisi et al., 2013). There is large number of public frozen seafood processing services distributed in the country, where a considerable number of people buy their fresh sea food products daily. Serious consequences relating

to national productivity and development can arise from lack of hygiene (Eyisi et al., 2013). In this study, we evaluated the incidence and antibiotic susceptibility pattern of *Vibrio* spp. isolated from commonly consumed sea foods in Port-Harcourt, Nigeria.

## MATERIALS AND METHODS

### Sample collection, cultivation and identification of *Vibrio* spp.

Fresh samples of variety of sea foods including prawn, periwinkles, crayfish and shrimp were collected randomly from different sellers at the two major household markets (Mile 1 and Mile 3) in Port-Harcourt city, Rivers State, Nigeria. The samples were transported to the laboratory immediately in a screw cap bottles containing enrichment medium (alkaline peptone water) with alkalinity ranging from 8.6- 9.0 and incubated at 37°C for 8-10 h. The samples were divided into 2 sets, one set of the samples was directly cultured onto the Thiosulphate Citrate Bile Salt Agar and the other set was carried out with a pour plate technique using a 10 fold serial dilution method with thiosulphate citrate bile salt agar. The inoculated plates were incubated overnight at 37°C, no antibiotics were added to the TCBS plates. *Vibrio* spp. were identified using conventional microbiological tests: Oxidase positive, rapid and darting motility, sugar fermentation (which differentiates the various *Vibrio* spp.) and salt tolerance (Cheesbrough, 2000).

### Antibiotics susceptibility testing

The antibiotics susceptibility pattern of the isolates was determined using the disk diffusion method (Cheesbrough, 2000), on Mueller-Hinton agar (Oxoid, England). Inhibition zone diameter values were interpreted using standard recommendations of the Clinical Laboratory Standard Institute (CLSI, 2006). Susceptibility was tested against Ofloxacin (10 µg), Gentamycin (10 µg), Ciprofloxacin (10 µg), Ampicillin (30 µg), Streptomycin (30 µg), Pefloxacin (10 µg), Augmentin (30 µg), Septrin (30 µg), Ceporex (10 µg) (Oxoid, England). *Escherichia coli* ATCC 25922 was included as a reference strain.

### Statistical analysis

Analysis of variance (ANOVA) and students T-test were used to compare means, and values were considered significant at  $p < 0.05$ . Post hoc multiple comparisons for the ANOVA were done using least significant difference (LSD).

## RESULTS AND DISCUSSION

In this study, 63 samples of sea foods: 15 crayfishes, 18 prawns, 15 shrimps and 15 periwinkles were analyzed following standard microbiological method on TCBS Agar (Oxoid, England). The mean bacterial counts of the sea foods obtained from the two markets are shown in Table 1. Table 2 shows that there was no significant difference ( $P > 0.05$ ) in the frequency of occurrence of *Vibrio* species in the sea food collected from Mile 1 and Mile 3 markets. Table 3 shows the antibiotic susceptibility pattern of the *Vibrio* isolates from sea foods bought in Mile 1 and Mile 3 market were significantly ( $P < 0.05$ ) more susceptible to Ciprofloxacin, Ofloxacin, Pefloxacin, Gentamycin and

**Table 2.** Frequency of occurrence of *Vibrio* species isolated from prawns, periwinkle, crayfish and shrimp purchased in Mile 1 and Mile 3 markets.

Isolates	Mile 1					Mile 3			
	TNI	PR	PE	CR	SH	PR	PE	CR	SH
<i>V. cholerae</i>	2	0(0.00)	0(0.00)	0(0.00)	1(50)	1(50)	0(0.00)	0(0.00)	0(0.00)
<i>V. fluvialis</i>	30	5(16.7)	4(13.3)	3(10)	2(6.67)	4(13.3)	5(16.7)	4(13.3)	3(10)
<i>V. parahaemolyticus</i>	19	3(15.9)	1(5.3)	3(10)	2(10.5)	3(15.9)	3(15.9)	1(5.3)	3(15.9)
<i>V. vulnificus</i>	4	1(25)	0(0.00)	0(0.00)	0(0.00)	0(0.00)	1(50)	1(50)	0(0.00)
<i>V. mimicus</i>	1	0(0.00)	1(50)	0(0.00)	0(0.00)	0(0.00)	0(0.00)	0(0.00)	0(0.00)
<i>V. metschnikovi</i>	4	0(0.00)	0(0.00)	0(0.00)	0(0.00)	0(0.00)	0(0.00)	0(0.00)	0(0.00)
<i>Photobacterium</i> spp.	3	1(33.3)	0(0.00)	0(0.00)	0(0.00)	0(0.00)	0(0.00)	1(33.3)	0(0.00)

TNI = Total number of isolates; PR = Prawn; PE = periwinkle; CR = crayfish; SH = shrimp.

**Table 3.** The antibiotic susceptibility pattern of *Vibrio* species isolated from sea foods bought from Mile 1 and Mile 3 markets.

Isolates	TNI	OFX	PEF	CN	AU	CPX	SXT	S	PN	CEP
<i>V. cholerae</i>	2	2(100)	2(100)	2(100)	2(100)	2(100)	2(100)	2(100)	2(100)	1(50)
<i>V. fluvialis</i>	30	27(91.3)	28(93.3)	27(90)	29(96.4)	30(100)	23(76.7)	30(100)	26(86.7)	26(86.7)
<i>parahaemolyticus</i>	19	19(100)	19(100)	17(89.5)	16(84.4)	19(100)	15(78.9)	19(100)	12(63.2)	17(89.5)
<i>V. vulnificus</i>	4	4(100)	4(100)	4(100)	2(50)	4(100)	4(100)	3(75)	2(50)	4(100)
<i>V. mimicus</i>	1	1(100)	1(100)	1(100)	1(100)	1(100)	0(0.00)	1(100)	1(100)	0(0.00)
<i>V. metschnikovi</i>	4	2(50)	2(50)	4(100)	2(50)	4(100)	0(0.00)	4(100)	2(50)	2(50)

TNI: Total number of isolates; OFX: ofloxacin (10 µg); CN: Gentamycin (10 µg); CPX: Ciprofloxacin (10 µg); PN: Ampicillin (30 µg) S: Streptomycin (30 µg); PEF: Pefloxacin (10 µg); AU: Augmentin (30 µg); SXT: Septrin (30 µg); CEP: Ceporex (10 µg).

Streptomycin as compared to Augmentin, Septrin, Ampicillin and Ceporex.

This study indicates high occurrence of *V. fluvialis* and *V. parahaemolyticus* in the sea foods studied. Similar observations were made by Nwachukwu (2006) in studies on the pathogenic characteristics of non O<sub>1</sub> *Vibrio cholerae* isolated from sea foods in Nigeria and in Malaysia by Elhadi et al. (2004). High occurrence of *V. cholerae* has also been reported in Bangladesh by Faraque et al. (2003). This high prevalence may be attributed to their high salt tolerance ability. In contrast, Caldini et al. (1997) reported that 150 *Vibrio* species isolated from the Aron River basin in Italy, recorded *V. cholerae* non O<sub>1</sub> as the most prevalent specie with 82% occurrence. This was attributed to the fact that River Basin is fresh water.

In Tanzania, data collected during cholera epidemics in 1990 and 1991 showed that all *V. cholerae* O<sub>1</sub> strains were sensitive to Streptomycin, Gentamycin, Ciprofloxacin (Webber et al., 1998). Rajkarnitar (2000) reported that 25 strains of *V. fluvialis* were susceptible to Tetracycline, Streptomycin while 84% were resistance to Ampicillin, the results are in line with this research.

However, it is important to note that *V. fluvialis*, *V. parahaemolyticus*, *V. vulnificus*, *V. metschnikovi*, *V. cholerae*, *V. mimicus* and *Photobacterium* spp. in periwinkles, prawns, crayfish and shrimp and other sea foods that are rampant in Port-Harcourt might be

significantly responsible of gastroenteritis and severe diarrhea, wound infection cases among the population of the city considering the sources and high occurrence among the sea foods that were studied in this research.

Based on this research, the *Vibrio* species isolated showed different susceptibility to the antibiotics used. It was recorded that the *Vibrio* species isolated from the sea food sample showed 100% susceptibility to Streptomycin and Ciprofloxacin, followed by Ofloxacin, Pefloxacin, Gentamycin, Ceporex, Septrin and Augmentin and the least susceptible was Ampicillin. The antimicrobial susceptibility pattern of *Vibrio* species can be predicted easily, due to control in prescription of antimicrobial agents. Therefore it is clearly necessary to put more emphasis on food hygiene; thereby surveillance of potential contaminant bacteria in harvested seafood is crucial for sustenance of public health.

To prevent *Vibrio* infections, proper hygiene cooking, treatment of water supply and avoidance of eating raw food should be encouraged seriously; there should be establishment of a well separated sewage treatment infrastructure. Warning on possible cholera or other *Vibrio* spp. contamination should be reported around contaminated water sources with direction on how to decontaminate the water and sea foods for possible use. Effective sanitations practices should be practiced, if instituted and adhered to in time they are usually sufficient to stop an epidemic. Patients who present

diarrhea symptoms should be referred to the health centre or hospital for immediate diagnosis and treatment.

In conclusion, the *Vibrio* species isolated from sea foods (periwinkles, prawns, crayfish and shrimp) showed high incidence in *V. fluvialis*, followed by *V. parahaemolyticus* while the least incidence was observed in *V. mimicus* among sea foods sold in some market in Port-Harcourt, Nigeria. It is important to note that this research is laying serious emphasis on the need for high level of hygiene and proper cooking of sea foods before eating. It is also important to note the need to establish regular nationwide antibiotic susceptibility surveillance of *Vibrio* species in different parts of the country so as to provide guidance on the best option in different situation.

## REFERENCES

- Amita F, Choudhury SR, Thungapathr M, Ramamurthy T, Nair GB, Gosh A (2003). Etiological studies in hospital on patients with secretory diarrhea in Calcutta J. Indian Med Assoc. 96:14-15.
- Caldini G, Neri A, Cresti S, Boddi V, Rossolini GM, Lanciotti E (1997). High prevalence of *Vibrio cholerae* non-O1 carrying heat-stable enterotoxin-encoding genes among *Vibrio* isolates from a temperate-climate river basin of Central Italy. J. Appl. Environ. Microbiol. 63:2934-2939.
- Cheesbrough M (2000). District Laboratory Practice in Tropical Countries, Part 2. Cambridge University Press, Cambridge, UK; pp. 434.
- Clinical Laboratory Standards Institute (2006). Performance standards for Antimicrobial susceptibility testing. National committee for clinical laboratory standards, Wayne pa.
- Dickinson G, Lim KY, Jiang SC (2013). Quantitative Microbial Risk Assessment of Pathogenic Vibrios in Marine Recreational Waters of Southern California. Appl. Environ. Microbiol. 79(1):294-302.
- Dunstan RA, Heinz E, Wijeyewickrema LC, Pike RN, Purcell AW, Evans TJ, Praszker J, Robins-Browne RM, Strugnell RA, Korotkov KV, Lithgow T (2013). Assembly of the Type II Secretion System such as Found in *Vibrio cholerae* Depends on the Novel Pilotin AspS. PLoS Pathog. 9(1):303-309.
- Elhadi N, Radu S, Chen CH, Nishibuchi M (2004). Prevalence of potentially pathogenic *Vibrio* species in the seafood marketed in Malaysia. J. Food Protocol 67:1469-1475.
- Eyisi OA, Nwodo UU, Iroegbu CU (2013). Distribution of *Vibrio* species in shellfish and water samples collected from the Atlantic coastline of south-east Nigeria. J. Health Popul. Nutr. 31(3):314-320.
- Faraque SM, Chowdlury N, Kamaruzzamam M, Ahmed QS, Faraque AS, Salam MA (2003). Re-emergence of epidemic *Vibrio cholerae* 0139, Bangladesh. Emerg. Infect. Dis. 9:1116-1122.
- Islam A, Labbate M, Djordjevic SP, Alam M, Darling A, Melvold J, Holmes AJ, Johura FT, Cravioto A, Charles IG, Stokes HW (2013). Indigenous *Vibrio cholerae* strains from a non-endemic region are pathogenic Open Biol. 3(2):120-181.
- Kaysner CA, DePaola JA (2004). Vibrios In Food and Drug Administration Bacteriological Analytical Manual. Retrieved from on 3rd October, 2006.
- Lutz C, Erken M, Noorian P, Sun S, McDougald D (2013). Environmental reservoirs and mechanisms of persistence of *Vibrio cholerae*. Front Microbiol. 4:375-380.
- Miyoshi SI (2013). Extracellular proteolytic enzymes produced by human pathogenic *Vibrio* species, Front Microbiol. 4:339-345.
- Nwachukwu E (2006). Studies on the pathogenic characteristics of non O1 *Vibrio cholerae* isolated from seafoods in Nigeria. J. Eng. Appl. Sci. 1:284-287.
- Park JY, Jeon S, Kim JY, Park M, Osong SK (2013). Multiplex Real-time Polymerase Chain Reaction Assays for Simultaneous Detection of *Vibrio cholerae*, *Vibrio parahaemolyticus*, and *Vibrio vulnificus*. Public Health Res. Perspect. 4(3):133-139.
- Rajkarnitar A (2000). Antibiotics resistant of *Vibrio cholerae* isolated from Kathmandu valley and characterization of the isolates by biotyping and serotyping. A dissertation presented to control Dept of Microbiology.
- Webber JT, Minta ED, Canizaries R, Semigua A, Gomez I, Sempertugi R, Davila A, Greene KD, Cameron DN (1998). Epidemic cholera in Ecuador: Multi drug resistance and transmission by water and sea foods. Epidemiol. Infect. 112:1-11.



# Journal of Bacteriology Research

Related Journals Published by Academic Journals

- *Journal of Cell and Animal Biology*
- *International Journal of Genetics and Molecular Biology*
- *African Journal of Microbiology Research*
- *Journal of Evolutionary Biology Research*
- *International Journal of Biodiversity and Conservation*
- *Journal of Enzymology and Metabolism*
- *Journal of Environmental Microbiology and Microbial Ecology*

**academic**Journals