

African Journal of Mathematics and Computer Science Research

Volume 8 Number 3 June 2015

ISSN 1684-5315



ABOUT AJMCSR

The African Journal of Mathematics and Computer Science Research (ISSN 2006-9731) is published bi-monthly (one volume per year) by Academic Journals.

The African Journal of Mathematics and Computer Science Research (AJMCSR) (ISSN:2006-9731) is an open access journal that publishes high-quality solicited and unsolicited articles, in all areas of the subject such as Mathematical model of ecological disturbances, properties of upper fuzzy order, Monte Carlo forecast of production using nonlinear econometric models, Mathematical model of homogenous tumour growth, Asymptotic behavior of solutions of nonlinear delay differential equations with impulse etc. All articles published in AJMCSR are peer-reviewed.

Contact Us

Editorial Office: ajmcsr@academicjournals.org

Help Desk: helpdesk@academicjournals.org

Website: <http://www.academicjournals.org/journal/AJMCSR>

Submit manuscript online <http://ms.academicjournals.me/>

Editors

Prof. Mohamed Ali Toumi

*Département de Mahtématiques
Faculté des Sciences de Bizerte
7021, Zarzouna, Bizerte
Tunisia.*

Associate Professor Kai-Long Hsiao,

*Department of Digital Entertainment,
and Game Design,
Taiwan Shoufu University,
Taiwan,
R. O. C.*

Dr. Marek Galewski

*Faculty of Mathematics and Computer Science,
Lodz University
Poland.*

Prof. Xianyi Li

*College of Mathematics and Computational Science
Shenzhen University
Shenzhen City
Guangdong Province
P. R. China.*

Editorial Board

Dr. Rauf, Kamilu

*Department of mathematics,
University of Ilorin,
Ilorin, Nigeria.*

Dr. Adewara, Adedayo Amos

*Department of Statistics,
University of Ilorin.
Ilorin.
Kwara State.
Nigeria.*

Dr. Johnson Oladele Fatokun,

*Department of Mathematical Sciences
Nasarawa State University, Keffi.
P. M. B. 1022, Keffi. Nigeria.*

Dr. János Toth

*Department of Mathematical Analysis,
Budapest University of Technology and
Economics.*

Professor Aparajita Ojha,

*Computer Science and Engineering,
PDPM Indian Institute of Information
Technology,
Design and Manufacturing, IT Building,
JEC Campus, Ranjhi, Jabalpur 482 011 (India).*

Dr. Elsayed Elrifai,

*Mathematics Department,
Faculty of Science,
Mansoura University,
Mansoura, 35516, Egypt.*

Prof. Reuben O. Ayeni,

*Department of Mathematics,
Ladoke Akintola University,
Ogbomoso,
Nigeria.*

Dr. B. O. Osu,

*Department of Mathematics,
Abia State University,
P. M. B. 2000,
Uturu,
Nigeria.*

Dr. Nwabueze Joy Chioma,

*Abia State University,
Department of Statistics,
Uturu,
Abia State,
Nigeria.*

Dr. Marchin Papzhytski,

*Systems Research Institute,
Polish Academy of Science,
ul. Newelska 6 0-60-66-121-66,
03-815 Warszawa,
POLAND.*

Amjad D. Al-Nasser,

*Department of Statistics,
Faculty of Science, Yarmouk University,
21163 Irbid,
Jordan.*

Prof. Mohammed A. Qazi,

*Department of Mathematics,
Tuskegee University,
Tuskegee,
Alabama 36088,
USA.*

Professor Gui Lu Long

*Dept. of Physics,
Tsinghua University,
Beijing 100084,
P. R. China.*

Prof. A. A. Mamun, Ph. D.

*Ruhr-Universitaet Bochum,
Institut fuer Theoretische Physik IV,
Fakultaet fuer Phyik und Astronomie,
Bochum-44780,
Germany.*

Prof. A. A. Mamun, Ph. D.

*Ruhr-Universitaet Bochum,
Institut fuer Theoretische Physik IV,
Fakultaet fuer Phyik und Astronomie,
Bochum-44780,
Germany.*

Full Length Research Paper

On two-stage fuzzy random programming for water resources management

H. A. Khalifa

Department of Operations Research, Institute of Statistical Studies and Research, Cairo University, Cairo, Egypt.

Received 11 March, 2015; Accepted 3 June, 2015

In this paper, a two-stage fuzzy random programming for a management problem in terms of water resources allocation having fuzzy random variable coefficients and decision vector of random variables is studied. The first results show the fact that a fuzzy pseudorandom optimal solution of a two-stage fuzzy random programming may be resolved into a two of pseudorandom optimal solutions of relative two-stage random programmings. The subsequent results present a method that a fuzzy pseudorandom optimal solution of two-stage fuzzy random programming is structured by a two of pseudorandom optimal solutions of a relative two two-stage random programmings. A numerical example was given to clarify the obtained results.

Key words: Two-stage programming, fuzzy parameters, water resources management, interval analysis, fuzzy random variable, fuzzy approach.

INTRODUCTION

Stochastic programming deals with situations where some or all the parameters of a mathematical programming problem are described by stochastic variables rather than by deterministic quantities. Several models have been presented in the field of stochastic programming (Stancu-Minasian et al., 1976).

Contini (1978) developed an algorithm for stochastic goal programmings when the random variables are normally distributed with known means and variances. He transformed the stochastic problem into an equivalent deterministic quadratic programming problem, where the objective functions consisted of maximizing the probability of a vector of goals lying in the confidence region of a predefined size.

Teghem et al. (1986) and Leclercq (1982) have presented interactive methods in stochastic

programming. Two major approach to stochastic programming as recognized by Goicoechea et al. (1982) and Kambo (1984) are:

1. Chance constrained programming,
2. Two-stage programming.

In fuzzy decision making problems, the concept of a maximizing decision was proposed by Bellman and Zadeh (1990). Fuzzy linear programming problem with coefficients was formulated by Negoita (1970) and called robust programming.

Dubois and Prade (1982) investigated linear fuzzy constraints. Tanaka and Asai (1984) also proposed a formulation of fuzzy linear programming with fuzzy coefficients. Wang and Wang (1985) built another theory

*Corresponding author. E-mail: hamiden_2008@yahoo.com

Author(s) agree that this article remain permanently open access under the terms of the [Creative Commons Attribution License 4.0 International License](https://creativecommons.org/licenses/by/4.0/)

of fuzzy programming on the basis of "Satisfaction degree" which has been used in engineering design. Wang and Qiao (1993) put forward some model linear programming with fuzzy random variable coefficients, and studied the solution and distribution problems of a fuzzy random programming. Over the past decades, the conflict-laden issues of water resources allocation among competing municipal, industrial and agricultural interests have been of increasing concerns (Huang and Chang, 2003; Wang et al., 2003). The competition among water users has been intensified due to growing population shifts, shrinking water availabilities, varying natural conditions, and deteriorating quality of water resources (Li and Huang, 2008).

The increased water demands and the inadequate water supplies have exacerbated the shortage of water resources; this has been considered as a major obstacle to sustainable water resources management. When the essential demand cannot be satisfied due to insufficient resources, losses can hardly be avoided resulting in a variety of a diverse impact on Socio-economic development (Lu et al., 2010).

Wang and Adams (1986) proposed a two-stage optimization framework of planning reservoir operations, inflows were modeled as a periodic Markov processes. Eiger and Shamir (1991) developed a model for an optimal multi-period operation within a multi-reservoir system. Ferreto et al. (1998) examined a long-term by drothermal scheduling of multi-reservoir systems using a two-stage dynamic programming approach.

In many real-world problems, however, the quality and quantity of uncertain information is often not satisfactory enough to be presented as probabilities distribution. Even if such distributions for uncertain parameters are available, reflecting them in large-scale optimization models can be extremely challenging (Huang and Loucks, 2000).

Wang and Huang (2011) have also developed an interactive two-stage stochastic fuzzy programming approach through incorporating an interactive fuzzy resolution method within an inexact two-stage stochastic programming framework.

This paper will further be devoted to a model proposed in Wang and Huang (2011). Under a certain condition we discuss the solution method of a fuzzy two-stage stochastic programming, which has fuzzy random variable coefficients and the decision vector of random variables. Ammar and Khalifa (2014) introduced a fuzzy programming approach for treating an interactive two-stage stochastic rough interval water resource management.

PRELIMINARIES

In order to discuss our problem conveniently, we shall state some necessary results on interval arithmetic, fuzzy numbers and fuzzy random variables. Moore (1979)

introduced the concept of closed interval numbers.

$$\text{Let } I(R) = \{[a^L, a^U]\}, a^L, a^U \in R = (-\infty, \infty), a^L \leq a^U\}$$

denote the set of all closed interval numbers on R .

Definition 1:

Suppose that $[a^L, a^U], [b^L, b^U] \in I(R)$.

We define :

1. $[a^L, a^U](+)[b^L, b^U] = [a^L + b^L, a^U + b^U]$.
2. $[a^L, a^U](-)[b^L, b^U] = [a^L - b^L, a^U - b^U]$.
3. $[a^L, a^U](\cdot)[b^L, b^U] = [\min(a^L \cdot b^L, a^L \cdot b^U, a^U \cdot b^L, a^U \cdot b^U), \max(a^L \cdot b^L, a^L \cdot b^U, a^U \cdot b^L, a^U \cdot b^U)]$
 $= [a^L \cdot b^L \wedge a^L \cdot b^U \wedge a^U \cdot b^L \wedge a^U \cdot b^U,$
 $= [a^L \cdot b^L \vee a^L \cdot b^U \vee a^U \cdot b^L \vee a^U \cdot b^U]$.
4. The order relation " \leq " in $I(R)$ is defined by $[a^L, a^U](\leq)[b^L, b^U]$ if and only if $a \leq b^L, a^U \leq b^U$.
5. If $a_t \in I(R), t \in T$, where T is an index set, we define $\bigvee_{t \in T} a_t = \sup\{a_t : t \in T\}; \bigwedge_{t \in T} a_t = \inf\{a_t : t \in T\}$.

In particular, for $[a_t^L, a_t^U] \in I(R), t \in T$, we define:

$$\bigvee_{t \in T} [a_t^L, a_t^U] = [\bigvee_{t \in T} a_t^L, \bigvee_{t \in T} a_t^U]; \bigwedge_{t \in T} [a_t^L, a_t^U] = [\bigwedge_{t \in T} a_t^L, \bigwedge_{t \in T} a_t^U]$$

Throughout this paper, $F_0(R)$ denote the set of all bounded closed fuzzy numbers (that is, compact on R , for any $f \in F_0(R), f$ satisfies:

1. There exists $x \in R$ such that $f(x) = 1$;
2. For any $\alpha \in (0, 1], f_\alpha = [f_\alpha^L, f_\alpha^U]$ is a closed interval number on R ,

And observed that $R \subset I(R) \subset F_0(R)$.

Definition 2:

Assume that $*$ is an algebraic operation on $R; f \cdot g \in F_0(R)$.

The algebraic operation $*$ on $F_0(R)$ is defined $(f * g)(z) = \bigvee_{z=x*y} (f(x) \wedge g(y))$.

where * may be "+", "-", "·", etc.

The order relation " \leq " on $F_0(R)$ is defined by $f \leq g$ if and

only if and only if $f_\alpha \leq g_\alpha$ for any $\alpha \in (0, 1]$.

Definition 3:

Let $\{f_t, t \in T\} \subset F_0(R)$, $\alpha \in (0, 1]$.

- (a) $\bigwedge_{t \in T} (f_t)_\alpha$,
- (b) $\bigvee_{t \in T} f_t$ is defined by a fuzzy number $g \in F_0(R)$ such that $g_\alpha = \bigvee_{t \in T} (f_t)_\alpha$, where T is an index set.

Definition 4:

Let $f, g \in F_0(R)$ then for any $\alpha \in (0, 1]$, we have $(f * g)_\alpha = f_\alpha * g_\alpha$,

where * may be continuous algebraic operation (Qiao et al., 1994).

Definition 5:

Let (M, N, P) be a probability measure space.

A mapping $\tilde{a} : M \rightarrow F_0(R)$ is called a fuzzy random variable on (M, N) , if for any $\alpha \in (0, 1]$ $\tilde{a}_\alpha(w) = \{\alpha : x \in R, \tilde{a}(w)(*)\alpha\} = [a_\alpha^L(w), a_\alpha^U(w)]$ is a random variables on (M, N) (Wang and Zhang, 1992). Denote the set of all fuzzy random variables on (M, N) by $FR(M)$ (where $w \in M$).

Moreover, let * be an algebraic operation on $F_0(R)$. The algebraic operation * on $FR(M)$ may be defined by $(\tilde{a} * \tilde{b})(w) = \tilde{a}(w) * \tilde{b}(w)$ for any $w \in M$, and $\tilde{a} \leq \tilde{b}$ if and only if $\tilde{a}(w) \leq \tilde{b}(w)$ for any $w \in M$, where $\tilde{a}, \tilde{b} \in FR(M)$

If $\{\tilde{a}_t : t \in T\} \subset FR(M)$, then $(\bigwedge_{t \in T} \tilde{a}_t)(w) = \bigwedge_{t \in T} \tilde{a}_t(w)$, for any $w \in M$.

STATEMENT OF PROBLEM

A typical two-stochastic programming model for water resources management is (Wang and Huang (2011) as follows:

$$\begin{aligned} \max \quad & f = \sum_{i=1}^m NB_i T_i - E \left[\sum_{i=1}^m C_i S_{iQ} \right] \\ \text{s.t.} \quad & \sum_{i=1}^m (T_i - S_{iQ})(1 + \delta) \leq Q \end{aligned} \tag{1}$$

(Water availability constraints)

$$S_{iQ} \leq T_i \leq T_{i\max}, \forall i$$

(Water- allocation target constraints)

$$S_{iQ} \geq 0, \forall i$$

(Non- negativity and technical constraints)

Where f = system benefit (\$); NB_i = net benefit to user i per m^3 of water allocated ($\$/m^3$) (first- stage revenue parameters); T_i = allocated target for water that is promised to user $i(m^3)$ (first- stage decision variables); $E[.]$ = expected value of a random variable ; C_i = loss to user i per m^3 of water not delivered , $C_i > NB_i$ ($\$/m^3$) (second- stage cost parameters); S_{iQ} = shortage of water to user i when the seasonal flow is $Q(m^3)$ (second – stage decision variables); Q = total amount of seasonal flow (m^3) (random variables); δ_i = rate of water loss during transportation; $T_{i\max}$ = maximum allowable allocation amount for user $i(m^3)$; m = total number of water users; i = water user , $i = 1, 2, 3$, where $i = 1$ for the municipality, $i = 2$ for the industrial user, and $i = 3$ for the agricultural sector.

Based on the findings of Huang and Loucks (2000), the problem (1) can be reformulated as follows:

$$\max \quad f = \sum_{i=1}^m NB_i T_i - \sum_{i=1}^m \sum_{j=1}^n p_j C_i S_{ij}$$

$$\text{s.t. } \sum_{i=1}^m (T_i - S_{ij})(1 + \delta) \leq q_j, \forall j \quad (2)$$

$$S_{ij} \leq T_i \leq T_{i\max}, \forall i, j,$$

$$S_{ij} \geq 0, \forall i, j,$$

Where S_{ij} denotes the amount by which the water-allocation target (T_i) is not met when the seasonal flow is q_j with probability P_j .

In this paper, we study the problem (2) with fuzzy random variable coefficients and decision vector of random variables as:

Model 1: (two-stage fuzzy random programming)

$$\text{Find } (S, T) \text{ to max } f = \sum_{i=1}^m \tilde{N}B_i T_i - \sum_{i=1}^m \sum_{j=1}^n p_j \tilde{C}_i S_{ij} \text{ subject to:}$$

$$\sum_{i=1}^m (T_i - S_{ij})(1 + \delta) \leq q_j, \forall j$$

$$S_{ij} \leq T_i \leq T_{i\max}, \forall i, j, \quad S_{ij} \geq 0, \forall i, j,$$

Where, $\tilde{N}B_i, \tilde{C}_i, \tilde{\delta}, q_j \in FR(M)$.

The solution method of Model 1 will be discussed under the condition $\tilde{N}B_i \geq 0, \tilde{C}_i \geq 0$, and $\tilde{\delta} \geq 0$.

Model 1 may be rewritten as follows:

Model 2: Find (S^*, T^*) from \tilde{G} such that

$$\sum_{i=1}^m \tilde{N}B_i T_i^* - \sum_{i=1}^m \sum_{j=1}^n p_j \tilde{C}_i S_{ij}^* = \max \left(\sum_{i=1}^m \tilde{N}B_i T_i^* - \sum_{i=1}^m \sum_{j=1}^n p_j \tilde{C}_i S_{ij}^* \right),$$

$$(S, T) \in \tilde{G}$$

where

$$\tilde{G} = \{(S, T) : \sum_{i=1}^m (T_i - S_{ij})(1 + \tilde{\delta}) \leq \tilde{q}, \quad j = 1, \dots, m;$$

$$S_{ij} \leq T_i \leq T_{i\max}, i = 1, 2, \dots, m; S_{ij} \geq 0, i = 1, 2, \dots, m; j = 1, 2, \dots, n\}$$

Definition 6:

The (S^*, T^*) which satisfy the condition in Model 2, is called a fuzzy random optimal solution of Model 2.

Observe that Where, $\tilde{N}B_i, \tilde{C}_i, \tilde{\delta}, q_j \in FR(M)$, for any $\alpha \in (0, 1]$ their α -level cuts are random values denoted by

$$(\tilde{N}B_i)_\alpha = [(NB_i)_\alpha^L, (NB_i)_\alpha^U], (\tilde{C}_i)_\alpha = [(C_i)_\alpha^L, (C_i)_\alpha^U], (\tilde{\delta})_\alpha = [(\delta)_\alpha^L, (\delta)_\alpha^U], \mathbf{a}$$

$$\text{nd } (\tilde{q}_j)_\alpha = [(q_j)_\alpha^L, (q_j)_\alpha^U], \quad I = 1, 2, \dots, m; j = 1, 2, \dots, n.$$

Corresponding to Model 2, we structure the following programming problems:

Model 2.1: Find (S^*, T^*) from \tilde{G}_α such that

$$\sum_{i=1}^m (NB_i)_\alpha^L T_i^* - \sum_{i=1}^m \sum_{j=1}^n p_j (c_i)_\alpha^U S_{ij}^*$$

$$= \max_{(S, T) \in \tilde{G}_\alpha^U} \left(\sum_{i=1}^m (NB_i)_\alpha^L T_i - \sum_{i=1}^m \sum_{j=1}^n p_j (c_i)_\alpha^U S_{ij} \right)$$

where

$$\tilde{G}_\alpha^U = \{(S, T) : \sum_{i=1}^m (T_i - S_{ij})(1 + (\delta)_\alpha^U) \leq (q_j)_\alpha^U$$

$$S_{ij} \leq T_i \leq T_{i\max}, \quad i = 1, \dots, m; j = 1, \dots, n;$$

$$S_{ij} \geq 0, \quad i = 1, \dots, m; j = 1, \dots, n\}$$

for any $\alpha \in (0, 1]$.

Model 2.2: Find (S^*, T^*) from \tilde{G}_α such that

$$\sum_{i=1}^m (NB_i)_\alpha^U T_i^* - \sum_{i=1}^m \sum_{j=1}^n p_j (c_i)_\alpha^L S_{ij}^*$$

$$= \max_{(S, T) \in \tilde{G}_\alpha^L} \left(\sum_{i=1}^m (NB_i)_\alpha^U T_i - \sum_{i=1}^m \sum_{j=1}^n p_j (c_i)_\alpha^L S_{ij} \right)$$

for any $\alpha \in (0, 1]$.

For any given $\alpha \in (0, 1]$ Models 2.1-2.2 are linear programming problems with random variable coefficients; we may use the simple method discussed in Wang and Qiao (1993) to find random optimal solutions of the two models.

Lemma 1.

Let $\tilde{\delta} \geq 0$. If $(S, T) \in \tilde{G}$, then $(S^L, T^L) \in \tilde{G}_\alpha^L, (S^U, T^U) \in \tilde{G}_\alpha^U$ for any $\alpha \in (0, 1]$.

Proof:

$$(S, T) \in \tilde{G} \text{ implies } \sum_{i=1}^m (T_i - S_{ij})(1 + \tilde{\delta}) \leq \tilde{q}_j, j = 1, \dots, n; S_{ij} \leq T_i \leq T_i^{\max},$$

$$i = 1, \dots, m; j = 1, \dots, n; S_{ij} \geq 0, i = 1, \dots, m; j = 1, \dots, n$$

It follows that from Definitions 2 and 5:

$$\sum_{i=1}^m (T_i - S_{ij})(1 + (\tilde{\delta})_\alpha) = \sum_{i=1}^m (T_i - S_{ij})(1 + [\delta_\alpha^L, \delta_\alpha^U])$$

$$= \sum_{i=1}^m [(T_i - S_{ij})(1 + \delta_\alpha^L), (T_i - S_{ij})(1 + \delta_\alpha^U)]$$

$$= \left[\sum_{i=1}^m (T_i - S_{ij})(1 + \delta_\alpha^L), \sum_{i=1}^m (T_i - S_{ij})(1 + \delta_\alpha^U) \right].$$

Thus we obtain

$$\sum_{i=1}^m (T_i - S_{ij})(1 + \delta_\alpha^L) \leq (q_j^L)_\alpha, \quad j = 1, \dots, n,$$

$$\sum_{i=1}^m (T_i - S_{ij})(1 + \delta_\alpha^U) \leq (q_j^U)_\alpha, \quad j = 1, \dots, n,$$

for every $i = 1, \dots, m; j = 1, \dots, n$.

Therefore, $\sum_{i=1}^m (T_i - S_{ij})(1 + \delta_\alpha^U) \leq (q_j^U)_\alpha$, and

$$\sum_{i=1}^m (T_i - S_{ij})(1 + \delta_\alpha^L) \leq (q_j^L)_\alpha$$

That is to say, $(S^L, T^L) \in \tilde{G}_\alpha^L, (S^U, T^U) \in \tilde{G}_\alpha^U$.

Theorem 2.

Assume that

$$\tilde{\delta} \geq 0, \tilde{N}B_i \geq 0, \tilde{C}_i \geq 0, \tilde{G}_\alpha^L \subset \{(S^L, T^L) : (S, T) \in \tilde{G}\}$$

and $\tilde{G}_\alpha^U \subset \{(S^U, T^U) : (S, T) \in \tilde{G}\}$ where $\alpha \in (0, 1]$.

If (S^*, T^*) is a random optimal solution of Model 2, then for any $\alpha \in (0, 1]$, we have:

- (1) (S^{*L}, T^{*L}) is a random optimal solution of Model 2.2,
- (2) (S^{*U}, T^{*U}) is a random optimal solution of Model 2.1,

$$(3) \quad W_\alpha^L = \max \left(\sum_{i=1}^m (\tilde{N}B_i)_\alpha^L T_i - \sum_{i=1}^m \sum_{j=1}^n p_j (\tilde{C}_i)_\alpha^U S_{ij} \right), (S, T) \in \tilde{G}_\alpha^U,$$

$$W_\alpha^U = \max \left(\sum_{i=1}^m (\tilde{N}B_i)_\alpha^U T_i - \sum_{i=1}^m \sum_{j=1}^n p_j (\tilde{C}_i)_\alpha^L S_{ij} \right), (S, T) \in \tilde{G}_\alpha^L,$$

Where

$$\tilde{W} = \max \left(\sum_{i=1}^m (\tilde{N}B_i)_\alpha T_i - \sum_{i=1}^m \sum_{j=1}^n p_j \tilde{C}_i S_{ij} \right), (S, T) \in \tilde{G}, \tilde{W}_\alpha = [W_\alpha^L, W_\alpha^U].$$

Proof:

Suppose that (S^*, T^*) is a fuzzy random optimal solution of Model 2, then $(S^*, T^*) \in \tilde{G}$, and

$$\sum_{i=1}^m \tilde{N}B_i T_i^* - \sum_{i=1}^m \sum_{j=1}^n p_j \tilde{C}_i S_{ij}^* = \max_{(S, T) \in \tilde{G}} \left(\sum_{i=1}^m \tilde{N}B_i T_i - \sum_{i=1}^m \sum_{j=1}^n p_j \tilde{C}_i S_{ij} \right)$$

Since $\tilde{\delta} \geq 0$, by using Lemma 1, we can see that

$$(S^{*L}, T^{*L}) \in \tilde{G}_\alpha^L, (S^{*U}, T^{*U}) \in \tilde{G}_\alpha^U,$$

$$\tilde{G}_\alpha^L = \{(S^L, T^L) : (S, T) \in \tilde{G}\}$$

and

$$\tilde{G}_\alpha^U = \{(S^U, T^U) : (S, T) \in \tilde{G}\}.$$

It follows by Theorem 1, and Definition 1, and observing the conditions $\tilde{N}B_i \geq 0, \tilde{C}_i \geq 0, \tilde{\delta} \geq 0$ that is

$$\max \sum_{i=1}^m (\tilde{N}B_i)_\alpha T_i - \sum_{i=1}^m \sum_{j=1}^n p_j (\tilde{C}_i)_\alpha S_{ij}, (S, T) \in \tilde{G}$$

$$= \max \left(\sum_{i=1}^m (\tilde{N}B_i)_\alpha^L T_i - \sum_{i=1}^m \sum_{j=1}^n p_j (\tilde{C}_i)_\alpha^U S_{ij}, \sum_{i=1}^m (\tilde{N}B_i)_\alpha^U T_i - \sum_{i=1}^m \sum_{j=1}^n p_j (\tilde{C}_i)_\alpha^L S_{ij} \right),$$

$(S, T) \in \tilde{G}$, this leads to (3) of the Theorem 2 is true.

Numerical example

Consider the Model 1 with the following economic data illustrated in Table 1. At $\alpha = 0.5$.

The solution of Model 2.2 is

$$T_1 = 3, T_2 = 4, T_3 = 4, S_{11} = S_{12} = S_{13} = S_{21} = S_{22} = S_{23} = S_{33} = 0,$$

$$S_{31} = 9, S_{32} = 3.8, p^L = 373.6 \$.$$

The solution of Model 2.1 is

$$T_1 = 1, T_2 = 2, T_3 = 7, S_{11} = S_{12} = S_{13} = S_{22} = S_{23} = S_{33} = S_{31} = 0, S_{21} = 7.9365, S_{32} = 7.4348$$

$$p^U = 5606519\$.$$

The solution of Model 2 is

$$T_1 = [1, 3], T_2 = [2, 4], T_3 = [4, 7], S_{11} = S_{12} = S_{13} = S_{22} = S_{23} = S_{33} = 0$$

$$S_{21} = [0, 7.9365], S_{31} = [0, 9], S_{32} = [7.4348, 3.8],$$

Table 1. Model 1 economic data.

Activity	User		
	Municipal ($i=1$)	Industrial ($i=2$)	Agricultural ($i=3$)
Maximum allowable allocation $T_{i\max}$	8	8	8
Water-allocation target T_i	[1, 3]	[2, 4]	[4, 7]
Net benefit when water demand is satisfied $\tilde{N}B_i$	(80, 90, 100)	(50, 60, 70)	(30, 31, 32)
Reduction of net benefit when demand is not delivered \tilde{C}_i	(220, 240, 260)	(70, 80, 90)	(40, 50, 60)
Flow level	Probability p_j (%)	Seasonal flow \tilde{q}_j	
Low ($j=1$)	0.2	(2, 3, 4)	
Medium ($j=2$)	0.6	(8, 10, 12)	
High ($j=3$)	0.2	(16, 18, 20)	
Water loss $\tilde{\delta}$	-	(0.1, 0.2, 0.3)	

and $f = [f^L, f^U] = [3736, 5606519]\$$.

Conclusions

In this paper, a two-stage fuzzy random programming for water resources management having fuzzy random variable coefficients and decision vector of random variables has been presented. The results have shown that the fuzzy random optimal solution of the problem under consideration has relative two-stage random programmings.


Conflict of Interest

The author has not declared any conflict of interest.

REFERENCES

- Ammar EE, Khalifa HA (2014). Interactive two-stage stochastic fuzzy rough programming for water resources management. *J. Adv. Phy.* 4(3):622-642.
- Bellman RE, Zadeh LA (1990). Decision making in a fuzzy environment. *Manage. Sci.* 17:141-164.
- Contini B (1978). A stochastic approach to goal programming. *Oper. Res.* 16:576-586.
- Dubois D, Prade H (1982). System linear fuzzy constraints. *Fuzzy Sets Syst.* 13:1-10.
- Eiger G, Shamir U (1991). Optimal operation of reservoirs by stochastic programming. *Eng. Optim.* 17:293-312.
- Ferreto RW, Rivera JF, Shahidehpour SM (1998). A dynamic programming two-stage algorithm for long-term hydrothermal scheduling of multi reservoir systems. *IEEE Trans. Power Syst.* 13:1534-1540.
- Goicoechea A, Hansen DR, Duckstein L (1982). *Multiobjective Decision Analysis with Engineering and Business applications* (Wiley, New York).
- Huang GH, Chang NB (2003). The perspectives of environmental informatics and systems analysis. *Environ. Inform.* 1:1-6.

- Huang GH, Loucks DP (2000). An inexact two-stage stochastic programming model for water resources management under uncertainty. *Civ. Eng. Environ. Syst.* 17:95-118.
- Kambo NS (1984). *Mathematical Programming Techniques* (Affiliated East-West Press Pvt. Ltd)
- Leclercq JP (1982). Stochastic programming: an interactive multi criteria approach. *Eur. J. Oper. Res.* 10:33-41.
- Li YP, Huang GH (2008). Interval parameter resources management under uncertainty. *Water Res. Manage.* 22:681-698.
- Lu HW, Huang GH, He L (2010). Development of an interval-valued fuzzy linear programming method based on infinite α -cuts for water resources management. *Environ. Model. Softw.* 25:354-361.
- Moore RE (1979). *Method and applications of Interval Analysis* (SIAM, Philadelphia, PA).
- Negoita CV (1970). *Fuzziness in Management*, OPSA/TIMS, Miami.
- Qiao Z, Zaban Y, Wang G (1994). On fuzzy random linear programming. *Fuzzy Sets Syst.* 65:31-49.
- Stancu-Minasian I, Wels MJ (1976). A research bibliography in stochastic programming. *Oper. Res.* 24:1078-1119.
- Tanaka H, Asai K (1984). Fuzzy linear programming problem with fuzzy numbers. *Fuzzy Sets Syst.* 13:1-10.
- Teghem J Jr., Dufrance D, Thauvoye M, Kunch P (1986). An interactive method for multi-objective linear programming under uncertainty. *Eur. J. Oper. Res.* 26:65-82.
- Wang G, Wang W (1985). Fuzzy optimum design of structure. *Eng. Optim.* 8:291-300.
- Wang D, Adams BJ (1986). Optimization of real-time reservoir operations with Markov decision processes. *Water Res.* 22:345-352.
- Wang G, Zhang Y (1992). The Theory of fuzzy stochastic processes, *Fuzzy Sets Syst.* 51:161-178.
- Wang G, Qiao Z (1993). Convergence of sequences of fuzzy random variables and its application. *Fuzzy Sets Syst.* 59:295-311.
- Wang LZ, Fang L, Hipel KW (2003). Water resources allocation: A cooperative game theoretic approach. *J. Environ. Inform.* 2:11-22.
- Wang S, Huang GH (2011). Interactive two-stage stochastic fuzzy programming for water resources management. *J. Environ. Manag.* 92:1986-1995.



African Journal of Mathematics and Computer Science Research

Related Journals Published by Academic Journals

- *African Journal of Pure and Applied Chemistry*
- *International Journal of Physical Sciences*
- *Journal of Geology and Mining Research*
- *Journal of Environmental Chemistry and Ecotoxicology*
- *Journal of Internet and Information Systems*
- *Journal of Oceanography and Marine Science*
- *Journal of Petroleum Technology and Alternative Fuels*

academicJournals