

The background of the cover is a photograph showing the backs of several zebras standing in a shallow waterhole. The zebras are facing away from the camera, and their heads are lowered towards the water. The water reflects the zebras' black and white stripes. The foreground is filled with green grass.

Journal of Ecology and The Natural Environment

Volume 7 Number 9, September 2015

ISSN 2006-9847



*Academic
Journals*

ABOUT JENE

The **Journal of Ecology and the Natural Environment (JENE)** (ISSN 2006-9847) is published Monthly (one volume per year) by AcademicJournals.

Journal of Ecology and the Natural Environment (JENE) provides rapid publication (monthly) of articles in all areas of the subject such as biogeochemical cycles, conservation, paleoecology, plant ecology etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in JENE are peer-reviewed.

Submission of Manuscript

Please read the **Instructions for Authors** before submitting your manuscript. The manuscript files should be given the last name of the first author

[Click here to Submit manuscripts online](#)

If you have any difficulty using the online submission system, kindly submit via this email jene@academicjournals.org.

With questions or concerns, please contact the Editorial Office at jene@academicjournals.org.

Editors

Dr. Abd El-Latif Hesham

*Genetics Department, Faculty of Agriculture, Assiut University,
Assiut 71516,
Egypt*

Dr. Ahmed Bybordi

East Azarbaijan Research Centre for Agriculture and Natural Resources, Tabriz, Iran

Dr. Sunil Kumar

*Natural Resource Ecology Laboratory,
Colorado State University
1499 Campus Delivery, A204 NESB, Fort Collins,
Colorado-80526,
USA*

Prof. Gianfranco Rizzo

*University of Palermo
Dipartimento DREAM – Viale delle Scienze - Building
9. 90128
Palermo,
Italy*

Dr. Bahman Jabbarian Amiri

*Kiel University, Germany,
Ökologie-Zentrum der CAU
Abt. Hydrologie und Wasserwirtschaft
Olhausen Straße, 75
Kiel,
Germany*

Dr. Bikramjit Sinha

*National Institute of Science Technology and Development Studies,
Pusa Gate, Dr. KS Krishnan Marg, New Delhi 110012,
India*

Prof. Gianfranco Rizzo

*University of Palermo
Dipartimento DREAM – Viale delle Scienze - Building
9. 90128
Palermo,
Italy*

Associate Editors

Dr. Marko Sabovljevic

*Dept. Plant Ecology, Faculty of Biology,
University of Belgrade
Takovska 43, 11000 Belgrade,
Serbia*

Dr. Sime-Ngando Téléspore

*CNRS
LMGE, UMR 6023, Université Blaise Pascal, 63177,
Aubière Cedex
France*

Dr. Bernd Schierwater

*ITZ, Ecology and Evolution, TiHo Hannover
Büenteweg 17d, 30559 Hannover,
Germany*

Dr. Bhattacharyya Pranab

*North-East Institute of Science & Technology
Medicinal, Aromatic & Economic Plant Division,
North-East Institute of Science & Technology,
Jorhat-785006, Assam,
India*

Prof. Marian Petre

*University of Pitesti, Faculty of Sciences
1 Targul din Vale Street, Pitesti, 110040, Arges
County,
Romania.*

Prof. R.C. Sihag

*CCS Haryana Agricultural University
Department of Zoology & Aquaculture, Hisar-125004,
India*

Prof. Kasim Tatic

*School of Economics and Business, University of
Sarajevo
Trg oslobođenja 1, 71000 SARAJEVO,
Bosnia and Herzegovina*

Dr. Zuo-Fu Xiang

*Central South University of Forestry & Technology,
498 Shaoshan Nanlu,
Changsha, Hunan, China.*

Editorial Board

Dr. Zuo-Fu Xiang

*Central South University of Forestry & Technology,
498 Shaoshan Nanlu,
Changsha, Hunan, China.*

Dr. Pankaj Sah

*Higher College of Technology, Muscat,
Department of Applied Sciences,
(Applied Biology) Higher College of Technology,
Al-Khuwair, PO Box 74,
PC 133, Muscat
(Sultanate of Oman)*

Dr. Arti Prasad

*Mohan Lal Sukhadia University,
Udaipur, Rajasthan, India.
123, Vidya Nagar, Hiran Magri,
Sector-4, Udaipur, Rajasthan,
India*

Parviz Tarikhi

*Mahdasht Satellite Receiving Station
(Postal): No. 80, 14th Street, Saadat Abad Avenue,
Tehran 1997994313,
Iran*

Bharath Prithiviraj

*Post Doctoral Research Associate
Knight Lab, Dept. of Chemistry & Biochemistry
University of Colorado at Boulder
USA*

Dr. Melissa Nursey-Bray

*Australian Maritime College, Tasmania,
Australia*

Parvez Rana

*Department of Forestry and Environmental Science
Shahjalal University of Science and Technology
Bangladesh*

Mirza Hasanuzzaman

*Faculty of Agriculture, Sher-e-Bangla Agricultural
University
Sher-e-Bangla Nagar, Dhaka-1207,
Bangladesh*

Dr. Giri Kattel

*Murray Darling Freshwater Research Centre, La Trobe
University
471 Benetook Avenue, Mildura, Victoria 3500,
Australia*

Dr. M. Rufus Kitto

*Faculty of Marine Science-Obhur station,
King Abdulaziz University,
Jeddah 21589, Saudi Arabia*

Dr. Özge Zencir

*Kemah Vocational Training School,
Erzincan University, Kemah,
Erzincan, Turkey.*

Dr. Sahadev Sharma

*Laboratory of Ecology and Systematics,
Graduate School of Engineering and Science,
University of the Ryukyus, Senbaru 59,
Nishihara, Okinawa-903-0213 Japan*

Dr. Hasan Kalyoncu

*University of Süleyman Demirel,
Faculty of Art and Science,
Department of Biology,
32100 Isparta/Turkey*

Hammad Khan

*Department of Zoology and Fisheries,
University of Agriculture,
Faisalabad, Pakistan*

Mirza Hasanuzzaman

*Faculty of Agriculture,
Sher-e-Bangla Agricultural University
Sher-e-Bangla Nagar, Dhaka-1207,
Bangladesh*

Abdurrahman Dunder

*Siirt University, Science and Arts Faculty,
Department of Biology,
56000, Siirt, Turkey*

Meire Cristina Nogueira de Andrade

*College of Agronomic Sciences,
São Paulo State University, Brazil.*

Imran Ahmad Dar

*Dept. of Industries and Earth Sciences,
The Tamil University,
Ocean and Atmospheric Sciences & Technology Cell,
(A Unit of Ministry of Earth Sciences, Govt. of
India).*

S. Jayakumar

*Department of Ecology and Environmental
Sciences,
School of Life Sciences,
Pondicherry University,
Puducherry - 605 014, India*

Umer Farooq

*University of Veterinary & Animal Sciences
Lahore, Pakistan*

Instructions for Author

Electronic submission of manuscripts is strongly encouraged, provided that the text, tables, and figures are included in a single Microsoft Word file (preferably in Arial font).

The **cover letter** should include the corresponding author's full address and telephone/fax numbers and should be in an e-mail message sent to the Editor, with the file, whose name should begin with the first author's surname, as an attachment.

Article Types

Three types of manuscripts may be submitted:

Regular articles: These should describe new and carefully confirmed findings, and experimental procedures should be given in sufficient detail for others to verify the work. The length of a full paper should be the minimum required to describe and interpret the work clearly.

Short Communications: A Short Communication is suitable for recording the results of complete small investigations or giving details of new models or hypotheses, innovative methods, techniques or apparatus. The style of main sections need not conform to that of full-length papers. Short communications are 2 to 4 printed pages (about 6 to 12 manuscript pages) in length.

Reviews: Submissions of reviews and perspectives covering topics of current interest are welcome and encouraged. Reviews should be concise and no longer than 4-6 printed pages (about 12 to 18 manuscript pages). Reviews are also peer-reviewed.

Review Process

All manuscripts are reviewed by an editor and members of the Editorial Board or qualified outside reviewers. Authors cannot nominate reviewers. Only reviewers randomly selected from our database with specialization in the subject area will be contacted to evaluate the manuscripts. The process will be blind review.

Decisions will be made as rapidly as possible, and the journal strives to return reviewers' comments to authors as fast as possible. The editorial board will re-review manuscripts that are accepted pending revision. It is the goal of the AJFS to publish manuscripts within weeks after submission.

Regular articles

All portions of the manuscript must be typed double-spaced and all pages numbered starting from the title page.

The Title should be a brief phrase describing the contents of the paper. The Title Page should include the authors' full names and affiliations, the name of the corresponding author along with phone, fax and E-mail information. Present addresses of authors should appear as a footnote.

The Abstract should be informative and completely self-explanatory, briefly present the topic, state the scope of the experiments, indicate significant data, and point out major findings and conclusions. The Abstract should be 100 to 200 words in length. Complete sentences, active verbs, and the third person should be used, and the abstract should be written in the past tense. Standard nomenclature should be used and abbreviations should be avoided. No literature should be cited.

Following the abstract, about 3 to 10 key words that will provide indexing references should be listed.

A list of non-standard **Abbreviations** should be added. In general, non-standard abbreviations should be used only when the full term is very long and used often. Each abbreviation should be spelled out and introduced in parentheses the first time it is used in the text. Only recommended SI units should be used. Authors should use the solidus presentation (mg/ml). Standard abbreviations (such as ATP and DNA) need not be defined.

The Introduction should provide a clear statement of the problem, the relevant literature on the subject, and the proposed approach or solution. It should be understandable to colleagues from a broad range of scientific disciplines.

Materials and methods should be complete enough to allow experiments to be reproduced. However, only truly new procedures should be described in detail; previously published procedures should be cited, and important modifications of published procedures should be mentioned briefly. Capitalize trade names and include the manufacturer's name and address. Subheadings should be used. Methods in general use need not be described in detail.

Results should be presented with clarity and precision. The results should be written in the past tense when describing findings in the authors' experiments. Previously published findings should be written in the present tense. Results should be explained, but largely without referring to the literature. Discussion, speculation and detailed interpretation of data should not be included in the Results but should be put into the Discussion section.

The Discussion should interpret the findings in view of the results obtained in this and in past studies on this topic. State the conclusions in a few sentences at the end of the paper. The Results and Discussion sections can include subheadings, and when appropriate, both sections can be combined.

The Acknowledgments of people, grants, funds, etc should be brief.

Tables should be kept to a minimum and be designed to be as simple as possible. Tables are to be typed double-spaced throughout, including headings and footnotes. Each table should be on a separate page, numbered consecutively in Arabic numerals and supplied with a heading and a legend. Tables should be self-explanatory without reference to the text. The details of the methods used in the experiments should preferably be described in the legend instead of in the text. The same data should not be presented in both table and graph form or repeated in the text.

Figure legends should be typed in numerical order on a separate sheet. Graphics should be prepared using applications capable of generating high resolution GIF, TIFF, JPEG or Powerpoint before pasting in the Microsoft Word manuscript file. Tables should be prepared in Microsoft Word. Use Arabic numerals to designate figures and upper case letters for their parts (Figure 1). Begin each legend with a title and include sufficient description so that the figure is understandable without reading the text of the manuscript. Information given in legends should not be repeated in the text.

References: In the text, a reference identified by means of an author's name should be followed by the date of the reference in parentheses. When there are more than two authors, only the first author's name should be mentioned, followed by 'et al'. In the event that an author cited has had two or more works published during the same year, the reference, both in the text and in the reference list, should be identified by a lower case letter like 'a' and 'b' after the date to distinguish the works.

Examples:

Abayomi (2000), Agindotan et al. (2003), (Kelebeni, 1983), (Usman and Smith, 1992), (Chege, 1998;

1987a,b; Tijani, 1993,1995), (Kumasi et al., 2001) References should be listed at the end of the paper in alphabetical order. Articles in preparation or articles submitted for publication, unpublished observations, personal communications, etc. should not be included in the reference list but should only be mentioned in the article text (e.g., A. Kingori, University of Nairobi, Kenya, personal communication). Journal names are abbreviated according to Chemical Abstracts. Authors are fully responsible for the accuracy of the references.

Examples:

Chikere CB, Omoni VT and Chikere BO (2008). Distribution of potential nosocomial pathogens in a hospital environment. *Afr. J. Biotechnol.* 7: 3535-3539.

Moran GJ, Amii RN, Abrahamian FM, Talan DA (2005). Methicillinresistant *Staphylococcus aureus* in community-acquired skin infections. *Emerg. Infect. Dis.* 11: 928-930.

Pitout JDD, Church DL, Gregson DB, Chow BL, McCracken M, Mulvey M, Laupland KB (2007). Molecular epidemiology of CTXM-producing *Escherichia coli* in the Calgary Health Region: emergence of CTX-M-15-producing isolates. *Antimicrob. Agents Chemother.* 51: 1281-1286.

Pelczar JR, Harley JP, Klein DA (1993). *Microbiology: Concepts and Applications.* McGraw-Hill Inc., New York, pp. 591-603.

Short Communications

Short Communications are limited to a maximum of two figures and one table. They should present a complete study that is more limited in scope than is found in full-length papers. The items of manuscript preparation listed above apply to Short Communications with the following differences: (1) Abstracts are limited to 100 words; (2) instead of a separate Materials and Methods section, experimental procedures may be incorporated into Figure Legends and Table footnotes; (3) Results and Discussion should be combined into a single section.

Proofs and Reprints: Electronic proofs will be sent (e-mail attachment) to the corresponding author as a PDF file. Page proofs are considered to be the final version of the manuscript. With the exception of typographical or minor clerical errors, no changes will be made in the manuscript at the proof stage.

Fees and Charges: Authors are required to pay a \$550 handling fee. Publication of an article in the Journal of Ecology and the Natural Environment is not contingent upon the author's ability to pay the charges. Neither is acceptance to pay the handling fee a guarantee that the paper will be accepted for publication. Authors may still request (in advance) that the editorial office waive some of the handling fee under special circumstances

Copyright: © 2014, Academic Journals.

All rights Reserved. In accessing this journal, you agree that you will access the contents for your own personal use but not for any commercial use. Any use and or copies of this Journal in whole or in part must include the customary bibliographic citation, including author attribution, date and article title.

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract or as part of a published lecture, or thesis) that it is not under consideration for publication elsewhere; that if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher.

Disclaimer of Warranties

In no event shall Academic Journals be liable for any special, incidental, indirect, or consequential damages of any kind arising out of or in connection with the use of the articles or other material derived from the JENE, whether or not advised of the possibility of damage, and on any theory of liability.

This publication is provided "as is" without warranty of any kind, either expressed or implied, including, but not limited to, the implied warranties of merchantability, fitness for a particular purpose, or non-infringement. Descriptions of, or references to, products or publications does not imply endorsement of that product or publication. While every effort is made by Academic Journals to see that no inaccurate or misleading data, opinion or statements appear in this publication, they wish to make it clear that the data and opinions appearing in the articles and advertisements herein are the responsibility of the contributor or advertiser concerned. Academic Journals makes no warranty of any kind, either express or implied, regarding the quality, accuracy, availability, or validity of the data or information in this publication or of any other publication to which it may be linked.

Journal of Ecology and the Natural Environment

Table of Contents: Volume 7 Number 9, September 2015

ARTICLES

An integrated approach to delineation of the eco-climatic zones in Northern Nigeria

AbdulKadir, A., Usman, M. T. and Shaba, A. H.

Full Length Research Paper

An integrated approach to delineation of the eco-climatic zones in Northern Nigeria

AbdulKadir, A.^{1*}, Usman, M. T.¹ and Shaba, A. H.²

¹Department of Geography, Federal University of Technology, PMB 65, Minna, Bosso Campus Niger, Nigeria.

²Strategic Space Applications, National Space Research and Development Agency (NASRDA), PMB 437, Garki Abuja, Nigeria.

Received 18 June, 2015; Accepted 22 July, 2015

This study derived eco-climatic indices; a moisture quality index, onset and cessation dates, hydrologic growing season and Aridity Index all for the period (1950-2006), in addition to using National Oceanographic and Atmospheric Administration an Advanced Very High Resolution Radiometer images to derive a Soil Adjusted Vegetation Index map. These derived eco-climatic parameters were integrated using Geographic Information System (GIS) techniques. The overlay was hinged on a theoretical basis which holds that the variations of the Aridity index, cessation dates, length of the hydrologic growing season and vegetation index are directly proportional to eco-climatic zone changes while those for onset and MQI are inversely proportional. The quantitatively derived eco-climatic index and related maps identified five eco-climatic zones; wet, humid, sub-humid, dry sub-humid and semi-arid as against the three classic regional climatic zones. The eco-climatic map further unveils progressive transformation or southwards shifts in the boundaries of the regional climatic zones. Thus, the accurate delineation of the current eco-climatic zones would be essential for providing adequate information needed to achieve food security and sustainability of the physical environment that is fundamental to disaster risk reduction in sub-Saharan Africa.

Key words: Northern Nigeria, climate change, rainfall variability, southwards shifts, eco-climatic zones, food security.

INTRODUCTION

The potential impact of climate change, rainfall variability patterns and the dynamic hydrologic regimes have continue to escalate land degradation and make it imperative that the broad eco-climatic zones could have changed. Brian and Ed (1992) revealed that not only there are major shift in the ecological boundaries but also the character of the broadly distributed ecosystem and Abdulkadir et al (2013a) confirmed rapid advancement of

desert condition in the region. Similarly, Claire et al. (2008) concluded that climate change is inducing range shifts for many species as they follow their suitable climate space and further shifts are projected. Land degradation and it impacts on relevant socio-economic activities are typical phenomenon across northern Nigeria. The risks associated with these changes are real but highly uncertain (Ann et al., 2013). Specifically, the

*Corresponding author. E-mail: abuzaishatu@futminna.edu.ng. Tel: 08053020742 / 920003.

wide spread crop failure, hunger, famine and poverty, typical of region in recent times have necessitated identification of proactive measures for Disaster Risk Reduction (DRR) and disaster preparedness. Climatic and ecological changes threaten human health globally (Colin and David, 2010). Moreover, the existing climatic and eco-climatic parameter maps have been mapped on the basis of vegetation and annual rainfall totals using traditional vector maps overlay. It has been evident for a while that these parameters alone cannot explain conditions pertinent to environmental sustainability in dry and dry sub-humid environments (Muhammad et al., 2005).

These existing climatic and eco-climatic parameter maps have been mapped long ago. There is large consensus that in West Africa one of the major climate change impacts will be on rainfall, making it more variable and less reliable (Benoît, 2012). With time since they were first mapped over 40 years ago, changes have been occurring as a result of various land degradation processes resulting to likely shifts in boundaries and conditions of the mapped zones. Several studies have suggested that land-cover changes alter the weather (Mechi, 1994; Foodym 2001; Bounoua et al., 2002; and Zhang et al., 2006). Also, Julius (2015), asserts that changes in land use, particularly when joined with climate change and variability, could impact on ecosystems and natural resources in complex ways. Generally, there is the politics of the actual land area that is desertified; is it the Sudan/Sahel/Sudano-sahelian or the sub-humid zone? Understanding local disturbance regimes is fundamental to understanding changes in ecosystem properties and stability (Halpin, 1997). Similarly, Fraser et al. (2011) makes an empirical contribution to our understanding of how social-ecological systems are vulnerable to climate change. Also, the recurring crop failures experienced in this zone may be a response to inadequate information inherent in the use of the classic eco-climatic zones. With the power of Geographic Information System (GIS) technology and the need to track such changes, studies are now required that employ GIS tools in mapping eco-climatic zones in northern Nigeria.

Generally, drought occurrence, erratic rainfall distribution and shift in economic fortunes, all combine to impact on the environment. The most significant climatic change that has occurred has been a long-term reduction in rainfall in the semi-arid regions of West Africa (Nicholson, 2001). Apparently, aridity has been drastically altering the traditional eco-climatic zones resulting to recurring crop failure. Conversely, lack of information on the existing eco-climatic zone and changes that have taken place is an apparent problem. No researches have been carried out to investigate and map changes in the spatial extent of the traditional eco-climatic zones to establish new zones. Indeed, most research up till now has focused on analogues between present and future climates (Martin, 2014). These are gaps in the achievement of food

security and sustainability of the physical environment that the study intends to fill. Quantitative representativeness can characterize borders between eco-regions (Williams and Forrest, 2005). Delineation of current eco-climatic zones using modern geospatial techniques will provide accurate information on the state of the environment and could enhance agricultural productivity, quality of life and boost food security. As a country whose economy is heavily dependent on low-productivity rainfed agriculture, rainfall trends are often cited as one of the more important factors in explaining various socio-economic problems such as food insecurity (Wing et al., 2008).

Despite these apparent changes across the region and country in general, North/South climatic description and other broad regional climatic designation still dominate decision making. The two sub-climatic zones in Northern Nigeria are tropical continental north, tropical hinterland and the third zone is in response to relief; high plateaux. In addition, the eco-climatic parameter maps (onset, amount, cessation etc) produced independently is used in place of eco-climatic map. Hydrological, landscape and eco-regions were equally effective at defining regions interns of climate, landcover and water quality (David et al., 2004). Consequently, the integration of derived ecological and climatic parameters using Geographic Information System (GIS) to derive the eco-climatic map that will depict the eco-climatic condition in quantitative terms is crucial for sustainability of natural resources.

Study area

Northern Nigeria is located between Longitudes 3° to 15° East of the Greenwich meridian and Latitudes 9° and 14° north of the Equator, this region stretches from the Sokoto plains through the northern section of Hausa land to the Chad Basin. The extreme Northern part of the region share boundaries with the semi-arid and arid zone of the Niger Republic. The States located in this zone are Sokoto, Kebbi, Zamfara, Kastina, Kano, Jigawa, Yobe, Borno, Gombe, Bauchi, Adamawa, Kwara, Plateau, Taraba, Niger, Abuja (FCT), Nassarawa, Kogi and Benue States. The northern region, due to its latitudinal location is characterized by tropical continental dry climate. Rainfall amount is low, erratic and characterized by spatial and temporal variability. Agriculture is the most dominant economic activity consequently; the region has suffered great degradation from the hands of man and his livestock. The region is characterised by high temperature records and range, and dominated by savannah vegetation; Guinea to the south, Sudan and Sahel savanna to the north.

MATERIALS AND METHODS

The derived eco-climatic (Onset, Cessation, Hydrological Growing Season, Aridity Index and Moisture Quality Index) parameter

records (1950-2006) for the twelve selected meteorological stations in northern Nigeria were used (Abdulkadir et al., 2013b). The spatial spread of the globally referenced selected meteorological stations; Minna, Zaria, Yola, Maiduguri, Bauchi, Sokoto, Yelwa, Kano, Jos, Gausau, Nguru, and Katsina is taken as a sufficient basis for spatial interpolation of the non-observed locations in the zone. All the derived data are location specific except for vegetation. The location coordinates of the 12 data points were therefore used to extract Soil Adjusted Vegetation Index (SAVI) value. The derived eco-climatic parameters were defined numerically; 5, 4, 3, 2 and 1 indicating varying level of moisture effectiveness and the point data was transformed to spatial data X, Y, Z (longitude, latitude and factor) using krigging geo-statistical defaults in SURFER 8 to produce an accurate grid of the base map.

National Oceanographic Atmospheric Administration (NOAA) Advanced Very High Resolution Radiometer (AVHRR) satellite images of 1.1km spatial resolution over northern Nigeria was acquired from the National Remote Sensing Centre Jos, Nigeria. The atmospheric corrections were made to remove the effect of aerosols and obtain more accurate data from the satellite image since these particles; smoke, dust, and water vapour, strongly affect all spectral vegetation indices. They affect the contrast between the infrared (IR) and near infrared (NIR) values and lead to lower index values, whether they are based on the difference between the two ranges of NIR-Red or the ratio between the two (Dewidar and Al Rehili, 2013). Thereby optimizing visual interpretation and preparing the Images for SAVI transformation, ground control points were extracted from geo-referenced drainage map of Nigeria (rivers / streams intersection and shoreline features) and corresponding positions on the image were used to resample the image. The band 2 and 3 (0.72 - 1.1 μm and 3.55 - 3.93 μm) were transformed to vegetation data using Soil Adjusted Vegetation Index (SAVI) to quantify vegetation biomass in the region, where soil background is a significant component of the signal detected. SAVI has shown to be very sensitive to discriminate vegetation amount in sparsely vegetated areas (Chuvieco et al., 2002). Similarly Malingreau et al. (1989) selected AVHRR channel three to be preferable to channels four and five. The derived rainfall characteristics; onset (\emptyset), cessation (C), hydrological growing season (HGS), Aridity Index (AI), Moisture Quality index (MQI) and derived vegetation index (VI) were used to derive the eco-climatic indices for integration and delineation of the eco-climatic zones.

A Geospatial database was developed for the analysis and integration of the derived eco-climatic indices. These parameters were resampled using the spatial resolution of the image and imported into GIS environment (ArcGIS 9.3). These were input into an overlay module which superimposes the whole factor in a GIS environment for further analysis using equation below. Efficient and cost-effective methods are needed for delineating sub-field productivity zones to improve soil and crop site-specific management (Kitchen et al., 2005). The overlay was hinged on a theoretical basis which holds the values of the aridity index; cessation, hydrologic growing season and vegetation index are directly proportional to eco-climatic zones while those for onset and MQI are inversely proportional to the quantitative character of the eco-climatic zones. This is expressed as:

$$ECZ = AI * VI * HGS * C / \emptyset * MQI \quad (1)$$

By implication the highest value will be 625 for the best Eco-climatic Zone (ECZ) class and the least value was 0.04 for the lowest class. However, in reality the highest class for this analysis is 477.242 while the least is 0.202. As a result, the eco-climatic zones were ranked and interpreted as Wet Humid, Humid, Sub-humid, Dry Sub-humid and Semi-Arid/Arid (Wet Humid, for the best class and Semi-Arid/Arid for the worst) Table 1. These were used to delineate the final map showing the Eco-climatic zones.

Table 1. Eco-climatic Class Interpretation Scheme.

Eco-climatic class	Eco-climatic zones	Interpretation scheme
5	Wet	$350 \leq ECZ \leq 477$
4	Humid	$250 \leq ECZ \leq 350$
3	Sub-humid	$150 \leq ECZ \leq 250$
2	Dry Sub-humid	$50 \leq ECZ \leq 150$
1	Semi-Arid /Arid	$ECZ \leq 50$

The derived eco-climate zones features were extracted using contour lines to demarcate eco-climatic zones limit, and this was overlaid on Iloeje (1976)'s scanned and digitized climatic region map to visualize the differences and similarities that exist between the two.

RESULTS

The integration of the entire factors using the empirical relationship reflects the moisture feature typical of northern Nigeria. Eco-climatic index values decreased gradually from the south northwards as Maselli (et al., 1996) concludes that the substantial agreement between these sources testifies to the eco-climatic value of the product obtained. Signifying the accuracy of the index, the Eco-climatic Index (ECI) value ranged from 0.202 depicting extreme moisture stress conditions in the extreme north to 477.242 indicting the best eco-climatic condition in the southern belt of the region (Figure 1).

The derived eco-climatic index was reclassified quantitatively using Table 1; consequently five eco-climatic classes were identified (Figure 2); semi-arid/ arid, dry sub-humid, sub-humid, humid and wet eco-climatic zones. Climatic regions provide useful information for ecologist, natural resource managers and economic planners (Brian and Ed, 1992; Maselli et al., 1996; Yan et al., 2013). Arid/Semi-Arid eco-climatic zone is most wide spread in the northern Nigeria; this stretches from northwest about (Lat. 10°20'N) through the central portion (Lat. 11°12'N) to about Lat. 9°27'N along the northeast as portray on Figure 1. This eco-climatic condition covers the entire Sokoto, Zamfara, Katsina, Kano, Jigawa, Gombe, Yobe, Borno, and northern parts of Bauchi and Adamawa States. Moisture stress is homogeneous in this zone as evident in the eco-climatic index as it has the least eco-climatic index value ($ECZ < 50$), since all the eco-climatic factors have the poorest moisture condition in this zone. Southwards, eco-climatic zones are heterogeneous reflecting different levels of moisture effectiveness.

Dry Sub-humid climate extends from southern border of semi-arid/arid zone to Long.4° E 10° Lat. 8°N 40' through Long.8° E 9°, Lat.10°N 16' to Long.11° E 46' Lat.7°N 8'. It spread across northern Niger, Kaduna, Plateau and Taraba States, in addition to southern parts of Bauchi,




Figure 1. Eco-climatic index.




Figure 2. Eco-climatic zones.




Figure 3. Eco-climatic and Iloeje (1976) climatic zones comparison.

and Adamawa States. This zone is characterized by low eco-climatic index ($50 \leq ECZ \leq 150$) and are classified as dry sub-humid climate

Three varying strips of humid zones are identified; Sub-humid consists of a narrow zone stretching from the southern margin of Sudan climate to Lon.4°E 53 lat.8°N 7' through Lon.8°E 30 lat. 9°N 32' to Lon.10° E 35 lat.7° N 12'. This is found across the central portion of Kwara, Niger, Kaduna, Plateau and Taraba States with eco-climatic values $150 \leq ECZ \leq 250$. Humid ($250 \leq ECZ \leq 350$) spreads from the southern boundary of the Sub-humid zone to Lon.6°E 31 Lat.8°N26' and Lon. 9°E 7 Lat.7° N 53 covering southern parts of Kwara, Niger, Kaduna, Plateau and Taraba States as well as, northern parts of FCT and Nassarawa State. However, despite the fact large proportion of the region is dry, sub humid and humid there is small area that is wetter than humid condition that is been classified as the Wet eco-climatic zone which is found mainly in parts of FCT, Nassarawa and the extreme fringe of Niger and Taraba south. The highest eco-climatic index ($350 \leq ECZ \leq 448$) is recorded in this zone and is classified as Wet eco-climatic zone as apparent in Figure 2.

Comparison of eco-climatic zones and climatic regions

Iloeje (1976) identified four climatic regions in Nigeria; subequatorial south, tropical hinterland, tropical continental north and high Plateaux climate (Figure 3). This climatic classification was based on precipitation, temperature and relative humidity annual means values and totals. In contrast the current eco-climatic zone map is based on the integration of derived precipitation and vegetation data using Geospatial technique and empirical relationships to distinguish eco-climatic sub-units. As against Iloeje's classification that was based on broad regional levels and traditional vector map overlay, which does not identify micro variations in the ecological and climatic zones for planning purposes across the region. This study identified five eco-climatic zones; semi-arid/arid, dry sub-humid, sub-humid, humid and wet eco-climatic zones with underlying implication of drastic transformation of the regional climatic zone. The comparison established that the tropical continental north has been transformed to semi-arid/arid and dry sub-humid. The classical tropical hinterland revealed actual

characterization of varying humidity levels; dry sub-humid, sub-humid, humid and wet eco-climatic zones as evident from the on Figure 3. Consequently, comparison of the regional climatic and eco-climatic maps revealed significant transformation of the regional climatic zones.

Similarly, Centre for Climate change and Fresh water Resources Eco-climatic Atlas only plotted States and regional eco-climatic parameter maps for the six geo-political zones other than climatic or eco-climatic map. Furthermore, Nigeria Meteorological Agency produce annual rainfall prediction maps using derived rainfall parameter. However, all these do not integrate ecological parameter (vegetation biomass) with derived climatic parameter to depict the state of the environment in quantitative terms.

DISCUSSION

The mathematical combination of various layers to derive eco-climatic index and map of the northern Nigeria illustrate moisture variability typical of the region. Andrade et al. (2005) observed that spatial and temporal rainfall variability scenario is more compatible to the physical reality. The derived climatic and ecological variables are the building block for defining the zones in quantitative terms. The derived maps show clearly the vulnerability of the entire belt to drought, aridity and crop failure. Maps are needed to display ecosystem distribution and hierarchy (Tim and John, 1996). In addition, it identified the fundamental moisture condition patterns that could fill the gap of inaccurate and out-dated information evident of the developmental processes in the country. To be effective, management responses must now be rigorously assessed with focused and practical ecological analysis (Halpin, 1997). Reliable and sufficient geo-spatial information is shown here to be vital for sustainable development of any nation, since adequate, accurate maps and geo-spatial data base can provide basis for any developmental decision. As a result, these eco-climatic indicators and map are crucial environmental indicators and fundamental tool for natural resource management in this belt.

The homogeneous moisture stress identified in the Arid/ Semi-Arid zone beyond Latitude 10°N is an indication of aridity in northern Nigeria. Desertification is a serious threat to arid and semiarid environments-which cover 40% of the global land surface and are populated by approximately 1 billion humans (Veron e al., 2006). This trend is disastrous to the agricultural sector, rural livelihood and can escalate food insecurity. Similarly, Ann et al., (2013) concludes that Societal vulnerability to the risks associated with climate change may exacerbate ongoing social and economic challenges, particularly for those parts of society which dependent on resources that are sensitive to changes in climate.

These derived eco-climatic indices provide detailed and

accurate information on the eco climatic zones. Furthermore, the derived eco-climatic map depicts the present State of eco- climatic zones in the region. Designation of tension zones is an important prerequisite for formulating national polices that address land degradation and desertification (Eswaran and Reich, 1998; Alex, 2013). The derived eco-climatic map reveals that the eco-climatic index is fundamental for identifying, delimiting, and describing the major types of eco-climatic conditions in quantitative terms; moreover, it is sensitive to climatic variation across the region.

The most important point which emerges from the study is the need to identify the heterogeneity in the eco-climatic zones across the belt in the national planning policy. In addition, designs a tool for adaptation to the changing environment, enhance agricultural productivity, poverty alleviation and environmental sustainability. The sustainable management of environmental resources is not possible until their worth is appreciated and an appropriate price is paid for the benefits received (Soussan, 1992). The identified variability in the derived eco-climatic zone is of tremendous importance for farmers and, indeed, all the inhabitants, because agricultural productivity to large extent is a function of eco-climatic condition. Therefore, identification of these zones could enhance agricultural suitability planning across northern Nigeria. It is inevitable from the resultant map to promote water control and conservation particularly, in areas beyond Latitude 10°N . The development of irrigation scheme in this arid and semi-arid zone will minimize the effect of drought, augment all year round cultivation, ensure enough pasture and water for livestock and lessen the dangers of flooding downstream as already evident in the region. Moreover, in severe drought years that onset dates and HGS are insignificant (especially around Nguru and Maiduguri) irrigation agriculture will minimise the effect and reduce southward migration that usually characterized such years.

The apparent and widespread desertification across the northern region could result to meteorological changes will reduce rainfall effectiveness. By inference, the identified changes imposed localised impacts on regional circulation system and super-imposed global variability pattern, indicating drought and crop failure are inevitable in the region. The spatio-temporal trend indicated that the belt will become drier as such that the dwindling availability of the vital natural resources required for survival will be aggravated. Suggesting, food insecurity, famine, loss of biodiversity, poverty, women and children labour and socio-economic instability will be severe in the region. Accurate assessment of the status, change, and trend of desertification will be instrumental in developing global actions to prevent and eradicate the problem (Yang et al., 2005). The majority of the population in this belt live in rural areas where the primary occupation is agriculture (rainfed), so the identified trend is a threat to economic growth. Consequently, identification of the

current eco-climatic zones provide accurate and adequate information fundamental for the achievement of food security and sustainability of the physical environment in the belt where most economies are mostly agro-based.

Iloeje (1976)'s climatic map identified four climatic zones over the whole of Nigeria; high plateaux, sub-equatorial south, tropical hinterland and tropical continental north. This classification was based on quantitative definition of temperature and precipitation values (Koppen, 1918; Strahler, 1969). Generally, Köppen's limits are basically simple averages of the directly observed values of the climatic elements. It has been evident for a while (Muhammad et al., 2005) that these parameters alone cannot explain conditions pertinent to environmental sustainability in dry and dry sub-humid environments. The challenge therefore, is to use accurate and reliable methods to compute the derived parameters (onset, cessation, HGS and MQI) with little or no generalization. For example, NIMET (2008) prediction delineated the north eastern zone as a zone with 80 – 120 days of HGS and CCCFR (2005) also classified north western zone with 100- 120 HGS days. How these days of potential plant growth and development will be distributed throughout the season and how the rainfall receipt could be translated into actual moisture available to plants is not considered. This has remained the single reason why despite apparent seasonal improvements in the total rainfall receipt, crop failure has remained an issue across the belt. Consequently, the present study captured the moisture stress across the belt, better identifying the extreme northeast as a zone with less than 60 days of HGS, the northwest extending to the northeast with less than 90 HGS days signifying that crop failure is inevitable across the zone.

Generally, the country's climatic classification is based on broad regional levels which do not identify sub-units for the purpose of land use planning and other practical purposes. In addition these classifications are based on methods that are about two centuries' old or cumulative values despite the growing level of knowledge and technology. Above all, researches have attested to the fact that mean and total rainfall figures are irrelevant for plant growth and development. Thornthwaite (1948), Denton and Barnes (1988), Andrade et al. (2005) and Dantas et al. (2007) identified precipitation effectiveness and plant growth as fundamental in climatic classification. Consequently, climatic classification on the basis of mean temperature and precipitation values is not crucial for plant requirement as does the classifications based on precipitation effectiveness and vegetation biomass. It identifies the remarkable evidence of progressive transformation of tropical continental north to Arid/ Semi-Arid eco-climatic condition and Dry Sub-Humid eco-climatic condition is the transition zone between the dry north and the humid south. Tropical hinterland shows diversified level of humidity; Sub-humid, Humid and wet eco-climatic zones. Ecosystems are liable to undergo sudden

discontinuous transitions from a vegetated to a desert State (Sonia et al., 2007), these necessitate the re-appraisal of the broad regional classification that has been in use and the need to capture the variation within the topical areas for developmental policy.

The integration of the derived eco-climatic parameters are crucial for identifying the state of the environment in quantitative term, at micro / pixel level and this will enhance decision making regarding crop suitability and sustainability of the natural environment. The quantitative description of the eco-climatic condition provides an indicator of natural environment sensitivity to climate change and climatic vulnerability that will enhance management of natural resources and lead to the attainment of sustainable development. David and Chasca (2005) considered climate change as 'special' amongst livelihood disturbing factors in the developing world. The eco-climatic index and map have depicts shift of the climatic zones to lower latitudes, necessitating the adaptation of high latitude species in the lower latitude for enhanced livelihood. Furthermore, all measures should be towards reducing this southward shift and the impacts of the shift by proper resource management strategies. Thus, the derived eco climatic map will enhance crop sustainability decision, increase food production and sustainability of the physical environment.

Homogenous moisture stress identified north of latitude 10° (Sudano-sahelian states) reveals that drought severity is mainly a function of the changes that are evident in the region. Consequently the derive map confirms that Sudano-sahelian states (sokoto, Kebbi, Katsina, Kano, Jigawa, BOrno, Yobe Bauchi and Gombe has now been transformed to semi-arid and arid environment and this is important for any planning on natural resource management. Climate forecasting is not the panacea to all our problems in agriculture; instead, it is one of many risk management tools that sometimes play an important role in decision-making Holger and Roger (2005). Consequently, the derived eco-climatic map constitutes proactive approach in the region sustainability decision particularly, those regarding resource deficits. As a result the current eco-climatic condition and map of northern Nigeria is crucial for enhance food security and sustainability of physical environment.

Conclusion

Integration of meteorological and remote sensing data using empirical relationship in GIS environment has the capability of providing pointers to be the best possible sustainable solutions to the critical environmental issues. The derived eco-climatic index and map integrated climatic and vegetation indices and therefore better related to sustainable agricultural productivity and food security as against climatic classification based on mean

temperature and rainfall values. Consequently, sustainability of agricultural productivity and other socio-economic activities in the region is primarily a function of accurate, up-to-date and adequate eco-climatic information on the state of the environment.

Conflict of interests

The author(s) did not declare any conflict of interest.

REFERENCES

- Abdulkadir A, Usman MT, Shaba AH, Saidu S (2013). An appraisal of the eco-climatic characteristics in Northern Nigeria. *Afr. J. Environ. Sci. Technol.* 7(8): 748-757.
- Abdulkadir A, Usman MT, Shaba AH, Shaba (2013). Climate Change, Aridity Trend and Agricultural Sustainability of the Sudano-sahelian Belt of Nigeria. Special Issue: Development and Sustainability in Africa – Part 1. *Int. J. Dev. Sustain.* 2 (2):1436-1456.
- Alex DS (2013). Climate change hotspots mapping: what have we learned? *Clim. Change* 123(1):23-37.
- Andrade JAS de, Bastos EA, Barros AHC, Silva CO da, Gomes AAN(2005) Climatic Classification and semiarid zone regionalization of the Piauí State, Brazil, under distinct rainfall scenarios. *Rev. Ciênc. Agron.* 36 (2): 143-151.
- Ann O, Ndubuisi EL, Wilfred U (2013) Risk Management and Challenges of Climate Change in Nigeria. *J. Hum. Ecol.* 41(3):221-235.
- Benoit S (2012). Present and future climate change in the semi-arid region of West Africa: a crucial input for practical adaptation in agriculture. *Atmospheric science letters* Published online in Wiley Online Library. <http://dx.doi.org/10.1002/asl.368>
- Bounoua L, Defries R, Collatz GJ, Seller P, Khan H (2002). Effects of Landcover on Surface Climate. *Clim. Change* 52: 29-64. <http://dx.doi.org/10.1023/A:1013051420309>
- Brian R, Ed N (1992) Assessing the Sensitivity of Canada's Ecosystem to Climatic Change. *Clim. Change* 21:37-55. <http://dx.doi.org/10.1007/BF00143252>
- Centre for Climate change and Fresh Water Resources (CCFR) (2005). *Eco-climatic Atlas*. MINNA FUTMIN Consult.
- Chuvieco E, Martí N MP, Palacios A (2002) Assessment of di Verent spectral indices in the red-near-infrared spectral domain for burned land discrimination. *Int. J. Remote Sens.* 23 (23):5103-5110. <http://dx.doi.org/10.1080/01431160210153129>
- Claire CV, Pam B, Paul O, Hans B, Bianca N, Jesse O, Claire B, Kuipers H (2008). Adapting landscapes to climate change: examples of climate-proof ecosystem networks and priority adaptation zones. *J. Appl. Ecol.* 45:1722-173. <http://dx.doi.org/10.1111/j.1365-2664.2008.01569.x>
- Colin DB, David H (2010). Primary, secondary and tertiary effects of eco-climatic change: the medical response. *Postgrad. Med. J.* 86:230-234. <http://dx.doi.org/10.1136/pgmj.2009.082727>
- Dantas AAA, Carvalho LG de, Ferreira E (2007). Climatic classification and tendencies in Lavras region, MG. *Ciênc. agrotec.* 31(6): 1862-1866. <http://dx.doi.org/10.1590/S1413-70542007000600039>
- David MW, Thomas LW, Gerard M (2004). Delineation and evaluation of Hydrologic- landscape Regions in United States Using GIS Tools and Multivariate Statistical Analysis. *Environ. Manage.* 32 (1):871-888.
- David SGT, Chasca T (2005). Adaptation to Climate Change: Perspectives Across Scales. *Glob. Environ. Change Part A* 15(2): 115-124.
- Denton SR, Barnes BV (1988). An Ecological Climatic Classification of Michigan: A Quantitative Approach. *For. Sci.* 34 (1):119 – 138.
- Dewidar KH, Al Rehili B (2013). Assessment of vegetation indices for estimating plant coverage and plant density Mountains, Saudi Arabia. *Merit Res. J. Agric. Sci. Soil Sci.* 1(2):19-32.
- Eswaran H, Reich P (1998). Desertification: A global assessment of land quality. *Proceedings of 16th International Congress of Soil Science*, Montpellier, France.
- Fraser EDG, Dougill AJ, Hubacek K, Quinn CH, Sendzimir J, Termansen M (2011). Assessing vulnerability to climate change in dryland livelihood systems: conceptual challenges and interdisciplinary solutions. *Ecol. Soc.* 16(3):3. <http://dx.doi.org/10.5751/ES-03402-160303>
- Halpin PN (1997). Global Climate Change and Natural-area Protection: Management Responses and Research Directions. *Ecol. Appl.* 7(3): 828-843. [http://dx.doi.org/10.1890/1051-0761\(1997\)007\[0828:GCCANA\]2.0.CO;2](http://dx.doi.org/10.1890/1051-0761(1997)007[0828:GCCANA]2.0.CO;2)
- Holger M, Roger CS (2005). Seasonal and Inter-Annual Climate Forecasting: The New Tool for Increasing Preparedness to Climate Variability and Change in Agricultural Planning and Operations. *Clim. Change* 70 (1-2):221-253. <http://dx.doi.org/10.1007/s10584-005-5948-6>
- Iloeje NP (1976). *A New Geography of Nigeria*. Nigeria, Longman Limited ISBN 0 582 65510 2
- Julius GT (2015). The Extent of Spatial Temporal Variations in Land Use and its Implications on Forest Ecosystems: A Case of Nandi Hills Forests Kenya. *Dev. Ctry. Stud.* 5(8):129 -139.
- Kitchen NR, Myers DB, Drummonde ST, Hong SY (2005). Delineating productivity zones on claypan soil fields using apparent soil electrical conductivity. *Comput. Electron. Agric.* 46 (1-3):285-308. <http://dx.doi.org/10.1016/j.compag.2004.11.012>
- Malingreau JP, Tucker CJ, Laporte N (1989). AVHRR for monitoring global tropical deforestation. *Int. J. Remote Sens.* 10: 855-867. <http://dx.doi.org/10.1080/01431168908903926>
- Martin B (2014). European Isotherms Move Northwards by up to 15 Km/Year: Using Climate Analogues for Awareness-Raising. *Eur. Sci. J. /SPECIAL/ edition ISSN: 1857 – 7881.* pp. 547-556.
- Maselli F, Petkov L, Maracchi G, Conese C (1996). Eco-climatic classification of Tuscany through NOAA-AVHRR data. *Int. J. Remote Sens.* 17(12): 2369-2384. <http://dx.doi.org/10.1080/01431169608948778>
- Mechi GA (1994). Influence of the Land Surface in the Asian Summer Monsoon: Extend Conditions Versus Internal Feedbacks. *J. Clim.* 7: 1033-1049. [http://dx.doi.org/10.1175/1520-0442\(1994\)007<1033:IOTLSI>2.0.CO;2](http://dx.doi.org/10.1175/1520-0442(1994)007<1033:IOTLSI>2.0.CO;2)
- Muhammad TU, Emma A, Peter J, Mark T (2005). A conceptual framework for enhancing the utility of rainfall hazard for agriculture in marginal environment. *Nat. Hazards* 34: 111-129. <http://dx.doi.org/10.1007/s11069-004-4349-x>
- Nicholson SE (2001). Climatic and environmental change in Africa during the last two centuries. *Clim. Res.* 17:123-144. <http://dx.doi.org/10.3354/cr017123>
- NIMET (2008). Year 2008 Rainfall Prediction and Socio-economic implication for all parts of Nigeria. *Annual Bulletin*, Abuja Nigeria Meteorological Services.
- Sonia K, Max R, Minus VB, Michel L (2007). Local facilitation, bistability and transitions in arid ecosystems. *Theor. Popul. Biol.* 71(3): 367-379. <http://dx.doi.org/10.1016/j.tpb.2006.09.003>
- Soussan JG (1992). Sustainable development. Environmental issues in the 1990s. John Wiley and Sons. pp. 20-33.
- Strahler A (1969). *Physical Geography*, Third Edition. New York, John Wiley and Sons, Inc.
- Thornthwaite CW (1948). An approach toward a rational classification of climate. *Geogr. Rev.* 38:55-94. <http://dx.doi.org/10.2307/210739>
- Tim H, John Y (1996). Spatial Prediction of Climatic state factor Regions in Alaska. *Ecoscience* 3(4): 490-500.
- Veron SR, Paruelo JM, Oesterheld M (2006). Assessing desertification. *J. Arid Environ.* 66(4): 751-763. <http://dx.doi.org/10.1016/j.jaridenv.2006.01.021>
- Williams WH, Forrest MH (2005). Delineation and Visualization of ecoregions. *Environ. Manage.* 32 (1) 839-860.
- Wing HC, Gabriel BS, Ashbindu S (2008). Trends and spatial distribution of annual and seasonal rainfall in Ethiopia. *Int. J. Climatol.* 28(13):1723-1734. <http://dx.doi.org/10.1002/joc.1623>
- Yan B, Sylvie G, David RG, Héloïse LG, Patrick L, Jacques M (2013). Fire regime zonation under current and future climate over eastern Canada. *Ecol. Appl.* 23(4):904-923. <http://dx.doi.org/10.1890/12->

0698.1

Yang X, Zhang K, Jai B, Ci L (2005). Desertification assessment in China: An overview. *J. Arid Environ.* 63(2):517-531. <http://dx.doi.org/10.1016/j.jaridenv.2005.03.032>

Zhang K, Li X, Zhou W, Zhang D, Yu Z (2006). Land resource degradation in China: Analysis of status, trends and strategy. *Int. J. Sustain. Dev. World Ecol.* 13(5):397-408. <http://dx.doi.org/10.1080/13504500609469689>

A herd of zebras is shown in a savanna landscape. In the foreground, a waterhole is visible, with the zebras' heads and necks reflected in the water. The background consists of green grass and trees under a bright sky.

Journal of Ecology and The Natural Environment

Related Journals Published by Academic Journals

- *African Journal of Environmental Science and Technology*
- *International Journal of Biodiversity and Conservation*
- *Journal of Yeast and Fungal Research*
- *Journal of Entomology and Nematology*
- *African Journal of Biology and Environment*
- *Journal of Evolutionary Biology Research*

academicJournals