


African Journal of Biochemistry Research

Volume 9 Number 11 , December 2015

ISSN 1996-0778



*Academic
Journals*

ABOUT AJBR

The **African Journal of Biochemistry Research (AJBR)** (ISSN 1996-0778) is published Monthly (one volume per year) by Academic Journals.

African Journal of Biochemistry Research (AJBR) provides rapid publication (monthly) of articles in all areas of Biochemistry such as Nutritional biochemistry, Analytical biochemistry, Clinical Biochemistry, Human and Plant Genetics, Molecular and Cell Biology, Enzymology, Toxicology, Plant Biochemistry, Biochemistry Education etc. The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles are peer-reviewed.

Submission of Manuscript

Please read the **Instructions for Authors** before submitting your manuscript. The manuscript files should be given the last name of the first author

[Click here to Submit manuscripts online](#)

If you have any difficulty using the online submission system, kindly submit via this email ajbr@academicjournals.org.

With questions or concerns, please contact the Editorial Office at ajbr@academicjournals.org.

Editor

Prof. Johnson Lin

*School of Biochemistry, Genetics, Microbiology and
Plant Pathology
University of KwaZulu-Natal (Westville)
Private Bag X 54001, Durban
Republic of South Africa*

Associate Editors

Gregory Lloyd Blatch

*Dept Biochemistry Microbiology & Biotechnology
Rhodes University Grahamstown 6140
South Africa*

Dr. Serap Yalin

*Mersin University,
Faculty of Pharmacy,
Department of Biochemistry,
Yenisehir Kampusu,
Mezitli 33161
Mersin/Turkey*

Dr. Om Prakash Gupta

*Directorate of Wheat Research (ICAR)
Post Box-158, A
grasain Marg, Karnal-132001, Haryana, India*

Editorial Board

Dr. Desouky A.M. Abd-El-Haleem

*Biological Sciences Department,
College of Arts and Sciences,
Qatar University, Doha,
Qatar*

Dr. S.K. Trigun

*Biochemistry and Molecular Biology Section,
Banaras Hindu University
Varanasi-221005,
India*

Dr. Imed Gallouzi

*McGill University,
Biochemistry Department,
3655 Promenade Sir William Osler Montreal,
Quebec, H3G 1Y6,
Canada*

Dr. Ashraf A Khalil

*Protein Technology Lab, Mubarak City for Science,
New Borg Elarab,
Alexandria,
Egypt.*

Dr. Stanley Mukanganyama

*Department of Biochemistry,
University of Zimbabwe, Box MP 167,
Mount Pleasant, Harare,
Zimbabwe*

Prof. Salah A. Sheweita

*Taibah University, Faculty of Medicine,
Department of Biochemistry, PO Box 30001,
Madinah,
Saudi Arabia*

Dr Oluwafemi O Oguntibeju

*Department of Clinical Biochemistry,
School of Medicine,
Spartan Health Sciences University,
P.O. Box 324, Vieux Fort, St Lucia,
West Indies*

Dr. Robert L. Brown

*USDA ARS,
Southern Regional Research Center
1100 Robert E. Lee Blvd.,
New Orleans, LA 70124*

Dr. Edward Eteshola

*Biomedical Engineering Center
Davis Heart and Lung Research Institute
Ohio State University
473 W. 12th Avenue
Columbus, OH 43210*

G. Suresh Kumar

*Senior Scientist and Head
Biophysical Chemistry Laboratory
Indian Institute of Chemical Biology
Council of Scientific and Industrial Research
Jadavpur,
Kolkata 700 032,
India*

Xu Lu

*Department of Biochemistry and Molecular Biology
Colorado State University
Fort Collins,
CO 80523-1870
USA*

Mohammed A.A Sarhan

*Dept. Biological Sciences
Faculty of Science
King Khalid University
Saudi Arabia*

Mehrdad Behmanesh

*Department Of Genetics
School Of Science
P.O.Box 114-175 Tehran Iran
Iran*

Hans Verhagen

*Po Box 1 3720 Ba Biltoven
The Netherlands
Netherlands*

P.K. Sumodan

*Post Graduate Department Of Zoology
Government College Madappally India
India*

Baleseng Moseki

*University Of Botswana
Botswana*

Bhaskar C. Behera

*Agharkar Research Institute
Plant Science Division India
India*

Luiz Claudio Miletto

*Universidade Do Estado De Santa Catarina
Brasil*

Oladipo Gabriel Sunday

*University Of Port Harcourt
Port Harcourt-Nigeria
Nigeria*

Basiouny Ahmed El-Gamal

*Biochemistry Department
Faculty Of Science
Alexandria University
Egypt*

Aminigo Ebiokpo Rebecca

*University Of Port Harcourt
Portharcourt-Nigeria
Nigeria*

Jia Zeng

*Department Of Bioengineering
Central South University
Changsha Hunan 410083 P.R.China
China*

Adenike Kuku

*Obafemi Awolowo University
Department Of Biochemistry
Nigeria*

Elsayed Hafez

*Genetic Engineering and Biotechnology Research
Institute
Egypt*

Gabriella Castoria

*Via L. De Crecchio 7 -80138 Naples
Department Of General Pathology
Italy*

Salwa Seddik Abdel-Latif

*21 Elbatal Ahmed Abdel Aziz
Elmohandesien Giza
Egypt*

Erasto Vitus Mbugi

*Muhimbili University
Biochemistry Department
School Of Medicine
India*

Mohamed Rholam

*Université Paris7 - Denis-Diderot
France*

Hooi Ling Foo

*Universiti Putra Malaysia
Malaysia*

Jayanth Rao

*Biochemistry And Nutrition
Cftri Mysore
India*

Maznah Ismail

*Universiti Putra
Malaysia*

Svetlana Lutsenko

*Oregon Health & Science University
USA*

Gabriel Ugwem

*Rivers State University Of Science And Technology
P.M.B. 5080 Port Harcourt
Nigeria*

Hari Chhatpar

*Dept. Of Microbiology & Biotechnology Centre
Faculty Of Science
M.S.University Of Baroda
Vadodara 390 002
Baroda India*

Mahiuddin Alamgir

*The University Of New South Wales
Sydney Nsw-2052
Australia*

Sheeja Samuel Edwin

*B.R Nahata College of Pharmacy & Research Centre
India*

William Cho

*Room 1305 13/F Block R Department of Clinical
Oncology
Queen Elizabeth Hospital
30 Gascoigne Road Kowloon
Hong Kong*

Dr Suraini Abd-Aziz

*Universiti Putra Malaysia
Malaysia*

Dr. Mustafa Numan Bucak

*Lalahan Livestock Central Research Institute
Lalahan
Ankara Turkey*

Alparslan Kadir Devrim

*Department Of Biochemistry
Faculty of Veterinary Medicine
Kafkas University 36040 Kars
Turkey*

Vasudev R. Thakkar

*Sardar Patel University
Brd School of Biosciences
Sardar Patel University
Nagar*

Prof. Emmanuel Anosike

*Department Of Biochemistry
University Of Port Harcourt
Nigeria*

Dr. Usama Beshay

*New Bourg El-Arab City, Research Area
Alexandria 21934
Egypt*

Dr. Ramar Perumal Samy

*Department of Anatomy
Yong Loo Lin School of Medicine
National University of Singapore
Singapore*

Dr. Shin-ichi ONO

*Laboratory of Clinical Pharmacy
College of Pharmacy, Nihon University
Japan*

Prof. Lawal Bilbis

*Biochemistry Department
Usmanu Danfodiyo University Sokoto
Nigeria*

Dr. Adriana G. Chicco

*Department of Biochemistry
University of Litoral, Santa Fe
Argentina*

Prof. Zia-Ur-Rahman

*Department Of Physiology and Pharmacology
University Of Agriculture
Faisalabad
Pakistan*

Dr. Oluwole Ariyo

*Allen University
USA*

Prof. Francisco Torrens

*Institut Universitari de Ciència Molecular
Universitat de València
Spain*

Prof. Belkhodja Moulay

*University of Senia Oran
Algeria*

Dr. Hossam M Ashour

*Department of Microbiology and Immunology
Faculty of Pharmacy, Cairo University
Egypt*

Dr. Fidelis Ocloo

*Biotechnology and Nuclear Agriculture Research
Institute/GAEC
Ghana*

Ass. Prof. Alfonso Baldi

*Dept. Biochemistry, Sect. Pathology
Second University of Naples,
Italy*

Dr. Anandh Babu Pon Velayutham

*Department of Human Nutrition
Foods and Exercise 253 Wallace Hall Virginia Tech
Blacksburg VA 24061
USA*

Dr. Tapan K. Chaudhuri

*Department of Biochemical Engineering and
Biotechnology
Indian Institute of Technology Delhi, Hauz Khas
New Delhi-110016, India.*

Dr. Rong Zhang

*Shenyang Pharmaceutical University
China*

Ass. Prof. Tzong-Jih Cheng

*Department of Bio-Industrial Mechatronics
National Taiwan University
Taiwan*

Dr. Zuyong Xia

*Department of Radiology,
1201 Welch Rd, Room P089, Stanford, CA 94301
USA*

Dr. Pratap Kumar Das

*Indian Institute of Chemical Biology
India*

Dr. Vasudeo Pandharinath Zambare

*Advanced Enzyme Technologies Ltd
India*

Dr. A M Mujumdar

*Agharkar Research Institute
India*

Prof. Christine Clayton

*ZMBH
Im Neuenheimer Feld 282
69120 Heidelberg
Germany*

Prof. Rekik Boulbaba

*ESA Mateur
Département des sciences et techniques de
productions animales
Tanzania*

Dr. Farhad Mirzaei

*National Dairy Research Institute, NDRI
Karnal
India*

Dr. ROUABHI Rachid

*Biology Department
Tebessa University.
Algeria*

Prof. Vaclav Vetvicka

*University of Louisville
USA*

Dr. Ramesh Putheti, Ph.D

*Research scientist
Actavis Pharmaceuticals
10065 red run blvd,owings mills
Blvd,Maryland.USA.21030
USA*

Prof. Dr. Mustafa NAZIROGLU

*Head of Department of Biophysics
Medical (TIP) Faculty, Suleyman Demirel
University
Cunur, TR-32260 Isparta
TURKEY*

Dr. José Luis Arias Mediano

*Grupo Investigación Farmacia Práctica (CTS-205)
Dept. Farmacia y Tecnología Farmacéutica
Facultad de Farmacia
Campus Universitario de Cartuja, s/n Universidad
de Granada
18071 Granada.*

Ahmed Malki, PhD

*Lecturer of Biochemistry and Molecular Biology
Biochemistry Department
Faculty Of Science
Alexandria University
Alexandria,
Egypt*

Dr. Alireza Seidavi (PhD)

*Assistant Professor of Animal and Poultry
Nutrition,
Department of Animal Science,
College of Agriculture,
Islamic Azad University, Rasht Branch,
Rasht, Iran*

Amani S. Awaad

*Professor of pharmacognosy, Chemistry
Department
Faculty of Sciences, King Saud University .
Riyadh. KSA. P.O. Box 22452, Riyadh 11495.
Saudi Arabia*

Dr. Abdel-Tawab Mossa

*Environmental Toxicology Research Unit (ETRU),
Pesticide Chemistry Department,
National Research Centre,
Dokki,
Egypt*

Dr. Amal A. Mohamed

*Plant Biochemistry Department,
Agriculture Division - National Research Center,
31-El-Tahrir St.,
Dokki,
Cairo – Egypt*

Dr. Anabella Gaspar

*Department of Biochemistry,
University of Pretoria,
South Africa*

Dr. Anna Janecka

*Department of Biomolecular Chemistry,
Medical University of Lodz,
Mazowiecka 6/8,
92-215 Lodz,
Poland*

Dr. Caser Abdel

*Horticulture Department,
Dohuk University,
Iraq*

Dr. David Sheehan

*Dept Biochemistry,
University College Cork,
Ireland*

Dr. Dayananda Chandrappa

*Center for Bioenergy,
Department of Life and Physical
Sciences,
Cooperative Research,
Lincoln University,
Jefferson City,
USA*

Dr. Elsayed Abdelaal

*Special Graduate Faculty,
University of Guelph,
Onatrio,
Canada*

Dr. Etienne Marbaix

*CELL Unit,
de Duve Institute,
UCL-75.41, 75 avenue
Hippocrate,
B-1200 Bruxelles,
Belgium*

Dr. Gary L. Firestone

*Department of Molecular and Cell Biology,
University of California,
Berkeley,
CA, 94720,
USA*

Dr. Henryk Zielinski

*Institute of Animal Reproduction and Food
Research,
Polish Academy of Sciences,
Poland*

Dr. Irshad A. Nawchoo

*Department of Botany,
University of Kashmir,
India*

Dr. Luchai Butkhup

*Department of Biotechnology,
Faculty of Technology,
Mahasarakham University,
Mahasarakham 44000,
Thailand*

Dr. Luminita Vladescu

*Department of Analytical Chemistry,
Faculty of Chemistry,
University of Bucharest,
Romania*

Dr. Mira Debnath

*School of Biochemical Engineering,
Institute of Technology - Banaras Hindu University,
Varanasi,
India*

Dr. Nilesh S. Panchal

*Department of Biosciences,
Saurashtra University,
Rajkot-360005,
Gujarat.
India*

Dr. Rayappa A. Balikai

*University of Agricultural Sciences,
Dharwad,
Karnataka- 580 005,
India*

Dr. Saad Tayyab

*Institute of Biological Sciences,
University of Malaya,
50603 Kuala Lumpur,
Malaysia*

Dr. Shijun Fu

*Institute of Health Sciences,
Shanghai Institutes for Biological Sciences,
Chinese Academy of Sciences and Shanghai Jiao
Tong University School of Medicine, Shanghai,
P. R. China*

Dr. Shiming Zhang

*Weis Center for Research,
Geisinger Clinic,
Danville, Pennsylvania,
USA*

Dr. Thomas Efferth

*Department of Pharmaceutical Biology,
Institute of Pharmacy and Biochemistry,
University of Mainz, Heidelberg,
55128 Mainz,
Germany*

Instructions for Author

Electronic submission of manuscripts is strongly encouraged, provided that the text, tables, and figures are included in a single Microsoft Word file (preferably in Arial font).

The **cover letter** should include the corresponding author's full address and telephone/fax numbers and should be in an e-mail message sent to the Editor, with the file, whose name should begin with the first author's surname, as an attachment.

Article Types

Three types of manuscripts may be submitted:

Regular articles: These should describe new and carefully confirmed findings, and experimental procedures should be given in sufficient detail for others to verify the work. The length of a full paper should be the minimum required to describe and interpret the work clearly.

Short Communications: A Short Communication is suitable for recording the results of complete small investigations or giving details of new models or hypotheses, innovative methods, techniques or apparatus. The style of main sections need not conform to that of full-length papers. Short communications are 2 to 4 printed pages (about 6 to 12 manuscript pages) in length.

Reviews: Submissions of reviews and perspectives covering topics of current interest are welcome and encouraged. Reviews should be concise and no longer than 4-6 printed pages (about 12 to 18 manuscript pages). Reviews are also peer-reviewed.

Review Process

All manuscripts are reviewed by an editor and members of the Editorial Board or qualified outside reviewers. Authors cannot nominate reviewers. Only reviewers randomly selected from our database with specialization in the subject area will be contacted to evaluate the manuscripts. The process will be blind review.

Decisions will be made as rapidly as possible, and the journal strives to return reviewers' comments to authors as fast as possible. The editorial board will re-review manuscripts that are accepted pending revision. It is the goal of the AJFS to publish manuscripts within weeks after submission.

Regular articles

All portions of the manuscript must be typed double-spaced and all pages numbered starting from the title page.

The Title should be a brief phrase describing the contents of the paper. The Title Page should include the authors' full names and affiliations, the name of the corresponding author along with phone, fax and E-mail information. Present addresses of authors should appear as a footnote.

The Abstract should be informative and completely self-explanatory, briefly present the topic, state the scope of the experiments, indicate significant data, and point out major findings and conclusions. The Abstract should be 100 to 200 words in length. Complete sentences, active verbs, and the third person should be used, and the abstract should be written in the past tense. Standard nomenclature should be used and abbreviations should be avoided. No literature should be cited.

Following the abstract, about 3 to 10 key words that will provide indexing references should be listed.

A list of non-standard **Abbreviations** should be added. In general, non-standard abbreviations should be used only when the full term is very long and used often. Each abbreviation should be spelled out and introduced in parentheses the first time it is used in the text. Only recommended SI units should be used. Authors should use the solidus presentation (mg/ml). Standard abbreviations (such as ATP and DNA) need not be defined.

The Introduction should provide a clear statement of the problem, the relevant literature on the subject, and the proposed approach or solution. It should be understandable to colleagues from a broad range of scientific disciplines.

Materials and methods should be complete enough to allow experiments to be reproduced. However, only truly new procedures should be described in detail; previously published procedures should be cited, and important modifications of published procedures should be mentioned briefly. Capitalize trade names and include the manufacturer's name and address. Subheadings should be used. Methods in general use need not be described in detail.

Results should be presented with clarity and precision. The results should be written in the past tense when describing findings in the authors' experiments. Previously published findings should be written in the present tense. Results should be explained, but largely without referring to the literature. Discussion, speculation and detailed interpretation of data should not be included in the Results but should be put into the Discussion section.

The Discussion should interpret the findings in view of the results obtained in this and in past studies on this topic. State the conclusions in a few sentences at the end of the paper. The Results and Discussion sections can include subheadings, and when appropriate, both sections can be combined.

The Acknowledgments of people, grants, funds, etc should be brief.

Tables should be kept to a minimum and be designed to be as simple as possible. Tables are to be typed double-spaced throughout, including headings and footnotes. Each table should be on a separate page, numbered consecutively in Arabic numerals and supplied with a heading and a legend. Tables should be self-explanatory without reference to the text. The details of the methods used in the experiments should preferably be described in the legend instead of in the text. The same data should not be presented in both table and graph form or repeated in the text.

Figure legends should be typed in numerical order on a separate sheet. Graphics should be prepared using applications capable of generating high resolution GIF, TIFF, JPEG or Powerpoint before pasting in the Microsoft Word manuscript file. Tables should be prepared in Microsoft Word. Use Arabic numerals to designate figures and upper case letters for their parts (Figure 1). Begin each legend with a title and include sufficient description so that the figure is understandable without reading the text of the manuscript. Information given in legends should not be repeated in the text.

References: In the text, a reference identified by means of an author's name should be followed by the date of the reference in parentheses. When there are more than two authors, only the first author's name should be mentioned, followed by 'et al'. In the event that an author cited has had two or more works published during the same year, the reference, both in the text and in the reference list, should be identified by a lower case letter like 'a' and 'b' after the date to distinguish the works.

Examples:

Abayomi (2000), Agindotan et al. (2003), (Kelebeni, 1983), (Usman and Smith, 1992), (Chege, 1998;

1987a,b; Tijani, 1993,1995), (Kumasi et al., 2001) References should be listed at the end of the paper in alphabetical order. Articles in preparation or articles submitted for publication, unpublished observations, personal communications, etc. should not be included in the reference list but should only be mentioned in the article text (e.g., A. Kingori, University of Nairobi, Kenya, personal communication). Journal names are abbreviated according to Chemical Abstracts. Authors are fully responsible for the accuracy of the references.

Examples:

Chikere CB, Omoni VT and Chikere BO (2008). Distribution of potential nosocomial pathogens in a hospital environment. *Afr. J. Biotechnol.* 7: 3535-3539.

Moran GJ, Amii RN, Abrahamian FM, Talan DA (2005). Methicillinresistant *Staphylococcus aureus* in community-acquired skin infections. *Emerg. Infect. Dis.* 11: 928-930.

Pitout JDD, Church DL, Gregson DB, Chow BL, McCracken M, Mulvey M, Laupland KB (2007). Molecular epidemiology of CTXM-producing *Escherichia coli* in the Calgary Health Region: emergence of CTX-M-15-producing isolates. *Antimicrob. Agents Chemother.* 51: 1281-1286.

Pelczar JR, Harley JP, Klein DA (1993). *Microbiology: Concepts and Applications.* McGraw-Hill Inc., New York, pp. 591-603.

Short Communications

Short Communications are limited to a maximum of two figures and one table. They should present a complete study that is more limited in scope than is found in full-length papers. The items of manuscript preparation listed above apply to Short Communications with the following differences: (1) Abstracts are limited to 100 words; (2) instead of a separate Materials and Methods section, experimental procedures may be incorporated into Figure Legends and Table footnotes; (3) Results and Discussion should be combined into a single section.

Proofs and Reprints: Electronic proofs will be sent (e-mail attachment) to the corresponding author as a PDF file. Page proofs are considered to be the final version of the manuscript. With the exception of typographical or minor clerical errors, no changes will be made in the manuscript at the proof stage.

Fees and Charges: Authors are required to pay a \$550 handling fee. Publication of an article in the African Journal of Biochemistry Research is not contingent upon the author's ability to pay the charges. Neither is acceptance to pay the handling fee a guarantee that the paper will be accepted for publication. Authors may still request (in advance) that the editorial office waive some of the handling fee under special circumstances

Copyright: © 2014, Academic Journals.

All rights Reserved. In accessing this journal, you agree that you will access the contents for your own personal use but not for any commercial use. Any use and or copies of this Journal in whole or in part must include the customary bibliographic citation, including author attribution, date and article title.

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract or as part of a published lecture, or thesis) that it is not under consideration for publication elsewhere; that if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher.

Disclaimer of Warranties

In no event shall Academic Journals be liable for any special, incidental, indirect, or consequential damages of any kind arising out of or in connection with the use of the articles or other material derived from the AJBR, whether or not advised of the possibility of damage, and on any theory of liability.

This publication is provided "as is" without warranty of any kind, either expressed or implied, including, but not limited to, the implied warranties of merchantability, fitness for a particular purpose, or non-infringement. Descriptions of, or references to, products or publications does not imply endorsement of that product or publication. While every effort is made by Academic Journals to see that no inaccurate or misleading data, opinion or statements appear in this publication, they wish to make it clear that the data and opinions appearing in the articles and advertisements herein are the responsibility of the contributor or advertiser concerned. Academic Journals makes no warranty of any kind, either express or implied, regarding the quality, accuracy, availability, or validity of the data or information in this publication or of any other publication to which it may be linked.

ARTICLES

- A Production and antioxidant activity of alcoholic beverages made from various colored rice and wild rice Rats** 130
Ryoichi TAKESHITA, Noriaki SAIGUSA and Yuji TERAMOTO

Full Length Research Paper

Production and antioxidant activity of alcoholic beverages made from various colored rice and wild rice

Ryoichi TAKESHITA, Noriaki SAIGUSA and Yuji TERAMOTO*

Department of Applied Microbial Technology, Faculty of Biotechnology and Life Science, Sojo University, Ikeda 4-22-1, Nishi Ward, Kumamoto 860-0082, Japan.

Received 22 August 2015; Accepted 2 October 2015

In order to develop a new type of alcoholic beverage with antioxidant activity, the authors carried out this research. In this study, alcoholic beverages were made from unpolished black rice, unpolished red rice, unpolished green rice, polished white rice and unpolished wild rice using conventional cooked fermentation and economical uncooked fermentation system. The resulting alcoholic beverages made from cooked and uncooked rice grains had an ethanol concentration of approximately 11.0 to 13.8% (v/v). Alcoholic beverages made from various grains had antioxidant activity. Especially, the 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging activity of the alcoholic beverages made from black rice, red rice and wild rice were relatively higher as compared to other alcoholic beverages. On the other hand, the inhibitory activity of lipid peroxidation of the alcoholic beverages made from wild rice using cooked and uncooked fermentation system was higher than that of the other alcoholic beverages. From these results, the possibility of new types of alcoholic beverages with antioxidant activity was suggested.

Key words: Antioxidant activity, fermentation, colored rice, wild rice.

INTRODUCTION

As progress is made in fermentation technology and the brewing industry, many kinds of alcoholic beverages are being produced each year to catch the interest of consumers. In Japanese sake brewing, the cooking of rice grains is an important process. Rice bran, which contains proteins, lipids, vitamins and minerals, is usually removed from the rice grains by polishing before fermentation to produce an alcoholic beverage of high quality. Black rice (*Oryza sativa* var. *Japocica* cv. *Shiun*) contains anthocyanin pigments, such as cyanidin and

peonidin glucosides, in the bran layer. Anthocyanin having antioxidant activity is known to have physiological functions and contributes to the prevention of diseases such as arterial sclerosis, and is a type of functional food. Many reports were published concerning antioxidant activity of various foods (Chen et al., 1998; Hamasaka et al., 2004; Saigusa and Teramoto, 2014).

In this study, the authors tried to brew novel alcoholic beverages from various unpolished colored rice and wild rice. The characteristics of the alcoholic beverage made

* Corresponding author. E-mail: yuji@bio.sojo-u.ac.jp.



Figure 1. Picture of various rice and wild rice.

from various grains were determined to develop the functional alcoholic beverage, which has antioxidant activity (Yuwa-amornpitak et al., 2012). For a comparative study, alcoholic beverages are brewed from nonglutinous rice as well.

MATERIALS AND METHODS

Yeast strain

Industrial sake yeast *Saccharomyces cerevisiae* K7 purchased from the Brewing Society of Japan (Tokyo, Japan) were used for ethanol fermentation. K7 yeast is maintained on the plates of an agar-solidified YPD medium (yeast extract, 10 g; peptone, 20 g; glucose, 20 g; tap water, 1,000 ml).

Rice grains and wild rice

Black rice (*O. sativa* var. *Japonica* cv. *Shiun*), red rice (*O. sativa* var. *Japonica* cv. *Engimai*), green rice (*O. sativa* var. *Japonica* cv. *Midorinoka*) purchased from Kajiwara Beikoku Co. Ltd. (Kyoto Japan) and wild rice (*Zizania aquatica*) purchased from Suzusho Ltd. (Tokyo, Japan) were used for ethanol fermentation (Figure 1). Wild rice is belonging to genus *Zizania* and actually, it is not rice.

For comparative study, commercial polished nonglutinous white rice grains (*O. sativa* var. *Japonica* cv. *Hinohikari*) were used as material for ethanol fermentation. These cereals were ground to particles of 2 to 3 mm in diameter with an electric grinder and were used for ethanol fermentation.

Saccharifying agent

A glucoamylase preparation, Sumizyme, kindly donated by Shin Nihon Kagaku Kogyo Co., Ltd. (Anjo, Japan), was used as the saccharifying agent. Sumizyme is known to have raw starch digestibility and digest cooked rice and uncooked rice as well. Sumizyme is applicable to uncooked fermentation procedure

(Teramoto et al., 1994).

Chemicals

1,1-diphenyl-2-picrylhydrazyl (DPPH) was purchased from Nacalai Tesque (Kyoto, Japan). Trolox (6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid) was purchased from Sigma-Aldrich, Inc. (St. Louis, Mo, USA). BHT (2,6-di-tert-butyl-p-cresol) was purchased from Tokyo Kasei Co., Ltd. (Tokyo, Japan).

Ethanol fermentation procedure

Conventional fermentation with cooking was performed according to the procedure below. Thirty grams of grains and 50 ml of deionized water were dispensed into a 300-ml Erlenmeyer flask and autoclaved at 121°C for 15 min. Various grains such as unpolished black rice, unpolished red rice, unpolished green rice, unpolished wild rice and polished white rice were used as material for ethanol fermentation. After cooling, the cooked rice was mixed with 0.2 g of Sumizyme as the saccharifying agent, 40 ml of deionized water and 10 ml of a yeast suspension, which readily brought the population of yeast in the initial mash to 3.0×10^7 cells/ml. Ethanol fermentation was conducted at 25°C in the dark (Figure 2). Ethanol fermentation without cooking was done as follows. Thirty grams of raw grains, 90 ml of deionized water, 0.2 g of Sumizyme, and 10 ml of a yeast suspension were dispensed into a 300-ml Erlenmeyer flask. The population of yeast in the initial mash was adjusted to 3.0×10^7 cells/ml, and fermentation was conducted in the same manner as for ethanol fermentation with cooking (Figure 3). The decrease in weight of the Erlenmeyer flask and its contents as a result of the evolution of CO₂ gas was measured every 24 h.

General analytical methods

Fermented mash made from nonglutinous rice grains was centrifuged at 3,000 rpm for 15 min and filtered through No. 101 filter paper (Advantec Toyo Co., Ltd., Tokyo, Japan), and the resulting alcoholic beverage was analyzed.




Figure 2. Procedure for ethanol fermentation with cooking.




Figure 3. Procedure for ethanol fermentation without cooking.

Acidity was measured by titrating 10 ml of alcoholic beverage with 0.1 N NaOH. Reducing sugar as glucose was determined according to the methods of Somogyi and Nelson (Somogyi, 1952; Nelson, 1944). The amount of total phenolic compounds, expressed as gallic acid, was determined according to the Folin-Ciocalteu method (Singleton et al., 1999; Hamasaka et al., 2004).

The ethanol concentration of alcoholic beverage was determined with a gas chromatograph (model GC-14A; Shimadzu Co., Kyoto, Japan) equipped with a 3.1-m PEG-HT column (Gasukuro Kogyo, Inc., Tokyo, Japan).

Determination of antioxidant activity

The DPPH radical scavenging activity as the Trolox equivalent was measured on the basis of the method of Yamaguchi et al. (1998). The lipid peroxidation inhibitory activity as the BHT equivalent was determined using β -carotene (Hamasaka et al., 2004).

RESULTS AND DISCUSSION

The fermentation curves of the mashes made from various grains such as unpolished black rice, unpolished red rice, unpolished green rice, unpolished wild rice, and polished white rice are shown in Figure 4. In the case of cooked fermentation, fermentation of the mash containing wild rice proceeded quickly in one day; while, fermentation of the mash containing black rice proceeded slowly in one day. Final CO₂ output of each mashes are ~11.2 to 12.2 (Table 1). In the case of uncooked fermentation, fermentation of the mash containing wild rice proceeded quickly until two day. Wild rice grain is relatively easy to be digested comparing with other rice grains belonging to *O. sativa*.

Picture of resulting alcoholic beverages are shown in Figure 5. Alcoholic beverage made from uncooked black

rice shows brilliant red color just like grape wine. While the color of alcoholic beverage made from cooked black rice is faded. During cooking process, a part of anthocyanin pigment might be denatured. Red rice, green rice, and wild rice shows pinky color, green color, and black color, however, resulting alcoholic beverages have pale yellow or pale brown color.

The characteristics of alcoholic beverage made from various grains are shown in Table 1. The ethanol concentration of the alcoholic beverages was approximately 11.0 to 13.8% (v/v). The total amount of phenolic compounds of alcoholic beverage made from uncooked grains were higher than that of the beverage made from cooked grains (Table 1 and Figure 6). Alcoholic beverages made from black rice contained anthocyanin. From the results of Table 1, it was confirmed that only alcoholic beverage made from black rice contains anthocyanin. Min-Kyoung Kim et al. (2008) reported that anthocyanin have been identified in hydrochloric acid - methanol extract of black rice and wild rice, respectively. And both main anthocyanin was a cyaniding-3-glucoside. Though it is a small amount, it may be appeared that there is a possibility that contain anthocyanin also in the alcoholic beverage made from wild rice, too.

The antioxidant activity of various alcoholic beverages was determined. The DPPH radical scavenging activities of alcoholic beverages made from colored rice and wild rice are relatively higher than that of alcoholic beverage made from polished white rice (Figure 7). A bran fraction of black rice or red rice was taken by changing the rice milling rate, to examine the DPPH radical scavenging activities. As a results, black rice had high activity in bran




Figure 4. Time courses of fermentation of the mashes made from various grains with cooking [left] and without cooking [right] using K7 yeast. ○, polished white rice; ●, black rice; △, red rice; ▲, green rice; ◆, wild rice. Values are the mean of triplicates.

Table 1. Composition of initial mash and analysis of resulting alcoholic beverages.

Rice	Polished white rice		Black rice		Red rice		Green rice		Wild rice	
	With cooking	Without cooking	With cooking	Without cooking	With cooking	Without cooking	With cooking	Without cooking	With cooking	Without cooking
Fermentation										
Initial pH	4.7	4.8	4.9	4.9	4.9	4.9	5.0	5.0	5.0	5.1
Final pH	4.5	4.5	4.6	4.9	4.7	4.8	4.6	4.9	4.9	5.0
CO ₂ output (g)	11.8	11.5	11.8	11.3	11.9	12.2	11.9	11.2	11.9	11.8
Filtrate (ml)	76	83	67	75	66	76	73	73	48	63
Acidity (ml)	2.2	2.4	3.3	2.9	3.6	3.3	3.2	2.8	2.8	2.5
Ethanol concentration (% w/v)	12.1	12.6	12.1	12.6	13.8	11.0	12.5	11.4	13.0	13.3
Reducing sugar content (µg/mL)	540	590	660	780	770	630	670	820	800	720
Total phenolic compound (µg/mL)	380	560	470	580	500	640	400	570	600	680
Anthocyanin content (µg/mL)	ND	ND	61.8	94.6	ND	ND	ND	ND	ND	ND

Values are the mean of triplicates. ND, not detected.




Figure 5. Pictures of alcoholic beverage made from various grains using K7 yeast.




Figure 6. Total phenolic compound of alcoholic beverages made from various grains using K7 yeast. Open bar shows alcoholic beverage made by fermentation with cooking and closed bar shows alcoholic beverage made by fermentation without cooking.

obtained from 90-100 and 80-90%, and red rice had high activity in bran obtained from 90-100% (Fujita et al., 2010). In both of black rice and red rice, there is a high anti-oxidant substance on the outside of unpolished rice, and it was thought that these components have been extracted, in this time.

The inhibitory activity of lipid peroxidation of the alcoholic beverages is shown in Figure 8. The inhibitory

activity of lipid peroxidation of the alcoholic beverage made from wild rice was much higher than that of other alcoholic beverage (Figure 8). It has been reported that the inhibitory activity of lipid peroxidation of 21 colored rice and 1 colorless rice water-ethanol mixed extract.

As a result, almost colored rice was higher antioxidant activity (Seok et al., 2006). However, from our results, other than the wild rice, there was no difference in the




Figure 7. DPPH radical scavenging activity of alcoholic beverages made from various grains using K7 yeast. Open bar shows alcoholic beverage made by fermentation with cooking and closed bar shows alcoholic beverage made by fermentation without cooking.




Figure 8. Inhibitory activity of lipid peroxidation of alcoholic beverage made from various grains using K7 yeast. Open bar shows alcoholic beverage made by fermentation with cooking and closed bar shows alcoholic beverage made by fermentation without cooking.

colored rice and polished white rice. In order to extract the component having the inhibitory activity of lipid peroxidation, it was thought to be necessary to increase the ethanol concentration.

It has been suggested that the compounds related with DPPH radical scavenging activity and the compounds related with inhibitory activity of lipid peroxidation are different. Chen et al. (1998) reported that peptides with lipid peroxidation inhibitory activity did not have DPPH radical-scavenging activity. It is also possible that the antioxidant substances in alcoholic beverages made from colored rice need to be identified, as they may be different substances.

DPPH radical scavenging activity and inhibitory activity of lipid peroxidation of the alcoholic beverage made from cooked and uncooked wild rice were relatively higher. The compounds having antioxidant activity contained in wild rice might be resistant against cooking. We would like to determine the compounds related with DPPH radical scavenging activity and the inhibitory activity of lipid peroxidation.

In this study, the colored rice pigment other than black rice could not be applied to alcoholic beverages. In the future, it is necessary to study the extraction technology of colored rice pigment in the alcoholic fermentation period. However, the alcoholic beverages made from black rice, red rice and wild rice showed high DPPH radical scavenging activity compared with polished white rice. In particular, alcoholic beverage made from wild rice showed high antioxidant activity regardless with or without cooking. In production of alcoholic beverages having functionality, usefulness of wild rice was suggested. From these results, the possibility of new types of alcoholic beverages with antioxidant activity was suggested.

Conflict of interest

The authors have not declared any conflict of interests.

REFERENCES

- Chen HM, Muramoto K, Yamauchi F, Fujimoto K, Nokihara K (1998). Antioxidative properties of histidine-containing peptides designed from peptide fragments found in the digests of a soybean protein. *J. Agric. Food Chem.* 46:49-53.
- Fujita A, Fujitake H, Kawakami K, Nomura M (2010). Antioxidant activity of colored rice bran obtained at different milling yields. *J. Oleo Sci.* 59:563-568.
- Hamasaka T, Kumazawa S, Fujimoto T, Nakayama T (2004). Antioxidant activity and constituents of propolis collected in various areas of Japan. *Food Sci. Technol. Res.* 10:86-92.
- Kim MK, Kim HA, Koh K, Kim HS, Lee YS, Kim YH (2008). Identification and quantification of anthocyanin pigments in colored rice. *Nutr. Res. Pract.* 2:46-49.
- Nelson N (1944). A photometric adaptation of the Somogyi method for the determination of glucose. *J. Biol. Chem.* 153:375-380.
- Saigusa N, Teramoto Y (2014). Effects of distillation process on antioxidant activity of Japanese traditional spirits rice-*shochu*. *International J. Biomass Renewab.* 3:17-23.
- Seok HN, Sun PC, Mi YK, Hee JK, Nobuyuki K, Mendel F (2006). Antioxidative activities of bran extracts from twenty one pigmented rice cultivars. *Food Chem.* 94:613-620.
- Singleton VL, Orthofer R, Lamuela-Raventos RM (1999). Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods Enzymol.* 299:152-178.
- Somogyi M (1952). Notes on sugar determination. *J. Biol. Chem.* 195:19-23.
- Teramoto Y, Saigusa N, Ueda S (1994). Effects of the cooking process on the characteristics of aromatic red rice wine. *J. Inst. Brew.* 100:155-157.
- Yamaguchi T, Takamura H, Matoba T, Terao J (1998). HPLC method for evaluation of the free radical-scavenging activity of foods by using 1,1-diphenyl-1-2-picrylhydrazyl. *Biosci. Biotechnol. Biochem.* 62:1201-1204.
- Yuwa-amornpitak T, Koguchi M, Teramoto Y (2012). Antioxidant activity of herbal wine made from cassava starch. *World Appl. Sci. J.* 16:874-878.

African Journal of Biochemistry Research

Related Journals Published by Academic Journals

- - *International Journal of Plant Physiology and Biochemistry*
 - *Current Research in Biochemistry and Molecular Biology*
 - *African Journal of Biotechnology*
 - *Journal of Developmental Biology and Tissue Engineering*
 - *Journal of Enzymology and Metabolism*

academicJournals