



**Journal of
Public Health and
Epidemiology**

Volume 8 Number 1 January, 2016

ISSN 2141-2316



*Academic
Journals*

ABOUT JPHE

The **Journal of Public Health and Epidemiology (JPHE)** is published monthly (one volume per year) by Academic Journals.

Journal of Public Health and Epidemiology (JPHE) is an open access journal that provides rapid publication (monthly) of articles in all areas of the subject such as health observatory, biostatistics, occupational health, behavioral medicine etc. The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in JPHE are peer-reviewed.

Submission of Manuscript

Submit manuscripts as e-mail attachment to the Editorial Office at: jphe@academicjournals.org. A manuscript number will be mailed to the corresponding author shortly after submission.

The Journal of Public Health and Epidemiology will only accept manuscripts submitted as e-mail attachments.

Please read the **Instructions for Authors** before submitting your manuscript. The manuscript files should be given the last name of the first author.

Editors

Professor Mostafa A. Abolfotouh

*Professor of Family & Community Medicine
Head of Medical Team - Biobanking Section.
King Abdullah International Medical Research
Center, King Saud Bin-Abdulaziz University for
Health Sciences, National Guard Health Affairs,
Saudi Arabia*

Editorial Board

Dr. Guolian Kang

*The University of Alabama at Birmingham/1665
University Blvd, Ryals 443
Guolian
USA*

Dr. Mohammed Danlami Salihu

*Public Health Department
Faculty of Veterinary Medicine
Usmanu Danfodiyo University, Sokoto.
Nigeria.*

Prof. Jahanfar Jahanban

*Oral Pathology Dept. Dental faculty of Tehran Islamic
Azad University/
Address: B 107 Pezeshkan-Farabi Build No 67 Javanshir
St. Hosseinabad Pasdaran St. Tehran
Iran*

Okonko, Iheanyi Omezuruike

*University of Ibadan, Ibadan, Nigeria
Nigeria*

Dr. Afroditi K Boutou

*Respiratory Failure Unit, Aristotle University of
Thessaloniki, "G. Papanikolaou", Hospital, 57010,
Exohi.
Greece*

Dr. Anil K. Philip

*Rajiv Academy for Pharmacy/ delhi-Mathura Highway,
NH#2, Mathura-281001, Uttar Pradesh, India
India*

Dr. Bijan Mohammad hosseini

*Ayatollah Kashani Social Security Hospital
P.O Box: 14515 - 799 Tehran - Iran
Iran*

Dr. Brajadulal Chattopadhyay

*Department of Physics, Jadavpur University, Kolkata-
700032, India
India*

Dr. Carlos H Orces

*Laredo Medical Center, 1700 East Saunders, Laredo
Texas 78041
USA*

Mrs Iscah A. Moth

*Ministry of Public Health and Sanitation
P.O. Box 1210-40100 Kisumu
Kenya*

Prof. Tariq Javed

*Department of Pathology, Faculty of Veterinary Science,
University of Agriculture, Faisalabad-38040.
Pakistan.*

Dr. María Elena Dávila L

*Universidad Centroccidental "Lisandro Alvarado".
School of Medicine/ School of Health Science . Av.
Andrés Bello C/ Av. Libertador. Barquisimeto, Lara,
Venezuela, SA*

Dr. Lay Ching Chai

*Centre of Excellence for Food Safety Research, Faculty of
Food Science and Technology, Universiti Putra Malaysia,
43400 UPM Serdang, Selangor,
Malaysia*

Dr. Liting Song

*Appointment pending, Public Health Agency of
Canada/Health Canada
809-50 Riddington Drive,
Toronto, ON M2K 2J8
Canada*

Dr. Joaquim Xavier Sousa Jr

*Laboratory Immunodermatology of Clinics Hospital -
Av Dr Eneas Carvalho Aguiar, 255 3th floor Room 3016
05403-000 Sao Paulo, Brazil
Brazil*

Dr. K.K.I.U. Arunakumara

*Institution/address - Dept. of Crop Science, Faculty of
Agriculture, University of Ruhuna, Mapalana,
Kamburupitiya, Sri Lanka
Sri Lanka*

Dr. Keya Chaudhuri

*Indian Institute of Chemical Biology
Raja S C Mullick Road, Kolkata-700032, India
India*

Belchiolina Beatriz Fonseca

*Universidade Federal de Uberlândia, Rua Ceará s/n,
bloco 2D. saça 43, Campus Umuarama, Uberlândia MG,
Brazil. Brazil*

Dr. Charles R. Doarn

*Associate Professor of Public Health and Biomedical
Engineering
Director, Telemedicine Program
Department of Public Health Sciences
University of Cincinnati
USA*

Instructions for Author

Electronic submission of manuscripts is strongly encouraged, provided that the text, tables, and figures are included in a single Microsoft Word file (preferably in Arial font).

The **cover letter** should include the corresponding author's full address and telephone/fax numbers and should be in an e-mail message sent to the Editor, with the file, whose name should begin with the first author's surname, as an attachment.

Article Types

Three types of manuscripts may be submitted:

Regular articles: These should describe new and carefully confirmed findings, and experimental procedures should be given in sufficient detail for others to verify the work. The length of a full paper should be the minimum required to describe and interpret the work clearly.

Short Communications: A Short Communication is suitable for recording the results of complete small investigations or giving details of new models or hypotheses, innovative methods, techniques or apparatus. The style of main sections need not conform to that of full-length papers. Short communications are 2 to 4 printed pages (about 6 to 12 manuscript pages) in length.

Reviews: Submissions of reviews and perspectives covering topics of current interest are welcome and encouraged. Reviews should be concise and no longer than 4-6 printed pages (about 12 to 18 manuscript pages). Reviews are also peer-reviewed.

Review Process

All manuscripts are reviewed by an editor and members of the Editorial Board or qualified outside reviewers. Authors cannot nominate reviewers. Only reviewers randomly selected from our database with specialization in the subject area will be contacted to evaluate the manuscripts. The process will be blind review.

Decisions will be made as rapidly as possible, and the journal strives to return reviewers' comments to authors as fast as possible. The editorial board will re-review manuscripts that are accepted pending revision. It is the goal of the JPHE to publish manuscripts within weeks after submission.

Regular articles

All portions of the manuscript must be typed double-spaced and all pages numbered starting from the title page.

The Title should be a brief phrase describing the contents of the paper. The Title Page should include the authors' full names and affiliations, the name of the corresponding author along with phone, fax and E-mail information. Present addresses of authors should appear as a footnote.

The Abstract should be informative and completely self-explanatory, briefly present the topic, state the scope of the experiments, indicate significant data, and point out major findings and conclusions. The Abstract should be 100 to 200 words in length. Complete sentences, active verbs, and the third person should be used, and the abstract should be written in the past tense. Standard nomenclature should be used and abbreviations should be avoided. No literature should be cited.

Following the abstract, about 3 to 10 key words that will provide indexing references should be listed.

A list of non-standard **Abbreviations** should be added. In general, non-standard abbreviations should be used only when the full term is very long and used often. Each abbreviation should be spelled out and introduced in parentheses the first time it is used in the text. Only recommended SI units should be used. Authors should use the solidus presentation (mg/ml). Standard abbreviations (such as ATP and DNA) need not be defined.

The Introduction should provide a clear statement of the problem, the relevant literature on the subject, and the proposed approach or solution. It should be understandable to colleagues from a broad range of scientific disciplines.

Materials and methods should be complete enough to allow experiments to be reproduced. However, only truly new procedures should be described in detail; previously published procedures should be cited, and important modifications of published procedures should be mentioned briefly. Capitalize trade names and include the manufacturer's name and address. Subheadings should be used. Methods in general use need not be described in detail.

Results should be presented with clarity and precision. The results should be written in the past tense when describing findings in the authors' experiments. Previously published findings should be written in the present tense. Results should be explained, but largely without referring to the literature. Discussion, speculation and detailed interpretation of data should not be included in the Results but should be put into the Discussion section.

The Discussion should interpret the findings in view of the results obtained in this and in past studies on this topic. State the conclusions in a few sentences at the end of the paper. The Results and Discussion sections can include subheadings, and when appropriate, both sections can be combined.

The Acknowledgments of people, grants, funds, etc should be brief.

Tables should be kept to a minimum and be designed to be as simple as possible. Tables are to be typed double-spaced throughout, including headings and footnotes. Each table should be on a separate page, numbered consecutively in Arabic numerals and supplied with a heading and a legend. Tables should be self-explanatory without reference to the text. The details of the methods used in the experiments should preferably be described in the legend instead of in the text. The same data should not be presented in both table and graph form or repeated in the text.

Figure legends should be typed in numerical order on a separate sheet. Graphics should be prepared using applications capable of generating high resolution GIF, TIFF, JPEG or Powerpoint before pasting in the Microsoft Word manuscript file. Tables should be prepared in Microsoft Word. Use Arabic numerals to designate figures and upper case letters for their parts (Figure 1). Begin each legend with a title and include sufficient description so that the figure is understandable without reading the text of the manuscript. Information given in legends should not be repeated in the text.

References: In the text, a reference identified by means of an author's name should be followed by the date of the reference in parentheses. When there are more than two authors, only the first author's name should be mentioned, followed by 'et al'. In the event that an author cited has had two or more works published during the same year, the reference, both in the text and in the reference list, should be identified by a lower case letter like 'a' and 'b' after the date to distinguish the works.

Examples:

Abayomi (2000), Agindotan et al. (2003), (Kelebeni,

1987a,b; Tijani, 1993,1995), (Kumasi et al., 2001) References should be listed at the end of the paper in alphabetical order. Articles in preparation or articles submitted for publication, unpublished observations, personal communications, etc. should not be included in the reference list but should only be mentioned in the article text (e.g., A. Kingori, University of Nairobi, Kenya, personal communication). Journal names are abbreviated according to Chemical Abstracts. Authors are fully responsible for the accuracy of the references.

Examples:

Chikere CB, Omoni VT and Chikere BO (2008). Distribution of potential nosocomial pathogens in a hospital environment. *Afr. J. Biotechnol.* 7: 3535-3539.

Moran GJ, Amii RN, Abrahamian FM, Talan DA (2005). Methicillinresistant *Staphylococcus aureus* in community-acquired skin infections. *Emerg. Infect. Dis.* 11: 928-930.

Pitout JDD, Church DL, Gregson DB, Chow BL, McCracken M, Mulvey M, Laupland KB (2007). Molecular epidemiology of CTXM-producing *Escherichia coli* in the Calgary Health Region: emergence of CTX-M-15-producing isolates. *Antimicrob. Agents Chemother.* 51: 1281-1286.

Pelczar JR, Harley JP, Klein DA (1993). *Microbiology: Concepts and Applications*. McGraw-Hill Inc., New York, pp. 591-603.

Short Communications

Short Communications are limited to a maximum of two figures and one table. They should present a complete study that is more limited in scope than is found in full-length papers. The items of manuscript preparation listed above apply to Short Communications with the following differences: (1) Abstracts are limited to 100 words; (2) instead of a separate Materials and Methods section, experimental procedures may be incorporated into Figure Legends and Table footnotes; (3) Results and Discussion should be combined into a single section. Proofs and Reprints: Electronic proofs will be sent (e-mail attachment) to the corresponding author as a PDF file. Page proofs are considered to be the final version of the manuscript. With the exception of typographical or minor clerical errors, no changes will be made in the manuscript at the proof stage.

Fees and Charges: Authors are required to pay a \$650 handling fee. Publication of an article in the Journal of Public Health and Epidemiology is not contingent upon the author's ability to pay the charges. Neither is acceptance to pay the handling fee a guarantee that the paper will be accepted for publication. Authors may still request (in advance) that the editorial office waive some of the handling fee under special circumstances.

Copyright: © 2015, Academic Journals.

All rights Reserved. In accessing this journal, you agree that you will access the contents for your own personal use but not for any commercial use. Any use and or copies of this Journal in whole or in part must include the customary bibliographic citation, including author attribution, date and article title.

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract or as part of a published lecture, or thesis) that it is not under consideration for publication elsewhere; that if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher.

Disclaimer of Warranties

In no event shall Academic Journals be liable for any special, incidental, indirect, or consequential damages of any kind arising out of or in connection with the use of the articles or other material derived from the JPHE, whether or not advised of the possibility of damage, and on any theory of liability.

This publication is provided "as is" without warranty of any kind, either expressed or implied, including, but not limited to, the implied warranties of merchantability, fitness for a particular purpose, or non-infringement. Descriptions of, or references to, products or publications does not imply endorsement of that product or publication. While every effort is made by Academic Journals to see that no inaccurate or misleading data, opinion or statements appear in this publication, they wish to make it clear that the data and opinions appearing in the articles and advertisements herein are the responsibility of the contributor or advertiser concerned. Academic Journals makes no warranty of any kind, either express or implied, regarding the quality, accuracy, availability, or validity of the data or information in this publication or of any other publication to which it may be linked.

ARTICLE

**Perception of obstetric danger signs among women living on the
coastline of the Atlantic Ocean in rural Lagos, Nigeria**

1

Bangboye M. Afolabi, Emmanuel N. U. Ezedinachi, Sidney Opara,
Iwara Arikpo and Abiodun Ogunwale

Full Length Research Paper

Perception of obstetric danger signs among women living on the coastline of the Atlantic Ocean in rural Lagos, Nigeria

Bangboye M. Afolabi^{1*}, Emmanuel N. U. Ezedinachi², Sidney Opara², Iwara Arikpo² and Abiodun Ogunwale³

¹Health, Environment and Development Foundation, 34 Montgomery Road, Yaba, Lagos, Nigeria.

²Institute of Tropical Disease and Research, University of Calabar, Calabar, Nigeria.

³Project Hope, Washington DC, USA.

Received 23 July, 2015; Accepted 29 September, 2015

Implementation of safe motherhood programs in sub-Saharan African nations requires recognition of signs of potential emergency obstetric cases to facilitate immediate care seeking and urgent medical intervention. This questionnaire-based survey aimed to describe the perception of obstetric danger signs among women of reproductive age, living on the Atlantic coastline of rural Lagos. This cross-sectional, descriptive study took place in two non-congruent communities of Lagos State, Nigeria, between October 2012 and March 2013. Epi Info 7 was used to calculate sample size for the projected population in each community. Statistical analysis was carried out using STATA 13 software. A total of 829 females (20% single, 80% married) participated in the study. Among these, about 76 and 79% disagreed that bleeding and high blood pressure were obstetric danger signs, respectively. Multivariate analysis showed a positive and significant association between awareness of bleeding as obstetric danger sign as the dependent variable and both knowledge of antenatal care ($t=6.53$, $p=0.000$) and knowing a woman who died in pregnancy ($t=3.34$, $p=0.001$) as independent variables. Rural women on Atlantic Ocean coastline of Lagos had low perception of obstetric danger signs indicating promotion of acceptable maternal health in this environment.

Key words: Perception, danger signs, obstetrics, reproductive age, Atlantic ocean coastline, rural.

INTRODUCTION

Globally, about 287,000 maternal deaths occurred in 2010, a drop of 47% from the figure given in 1990. Sub-Saharan Africa (sSA) (58%) and Southern Asia (30%) provided 88% of deaths among the Countdown countries in 2010 and within these regions, India (20%) and Nigeria (14.5%) alone accounted for more than one-third of all

maternal deaths (Adams and Franz-Vasdeki, 2013). Since the end of the last century however, great progresses have been recorded on the status of maternal health and safe motherhood as more years have been added to their life-expectancy, and fertility rates have substantially reduced not only in the developed

*Corresponding author. E-mail: bmafolabi@heendef.org. Tel: +234(0)803 049 0729; +234 (1) 291 6562.

Author(s) agree that this article remain permanently open access under the terms of the [Creative Commons Attribution License 4.0 International License](http://creativecommons.org/licenses/by/4.0/)

nations, but also in the developing countries (UN, 2003; Grown et al., 2005). These strides have led to substantial decline in burdens associated with childbirth and child rearing (Grown et al., 2005). Despite these advances, the level of maternal mortality in sSA could be regarded as a global crisis; in almost all countries and in all income groups. Women in child-bearing age are five times more likely at risk of these diseases (WHO, 2015a). The Millennium Development (MDG) Goal 5a, "Improve maternal health" stipulated that by 2015, maternal mortality should be reduced by a ratio of three-quarters (WHO, 2015). Progress on MDG-5 for maternal health has been slower than for MDG-4, with deaths declining from around 400,000 to around 275,000 (Lozano et al., 2011; UNFPA, 2012). The major causes of maternal deaths are hemorrhage (22.9%), hypertensive disorders (18.5%), abortion (14.6%), sepsis (8.6%), and other direct and indirect causes (35.5%) (Lozano et al., 2012).

Over the past few decades, great strides have been made in improvement of women's health status; more than a decade has been added to life expectancy, and fertility rates in both developed and developing countries have declined substantially, helping to reduce burdens associated with childbirth and child rearing (Grown et al., 2005). Despite this progress, more than half a million women, 99% of whom live in the developing world continue to die every year in pregnancy and childbirth due to entirely preventable reasons (WHO, 2015b). Maternal mortality of 630 deaths per 100,000 live births in Nigeria is unacceptably high (Hofmeyr et al., 2003). According to the World Health Organization (WHO), maternal deaths are higher in women living in rural areas and among poorer communities; young adolescents face a higher risk of complications and death as a result of pregnancy than older women; and between 1990 and 2013, global maternal mortality declined by only 2.6% instead of a decline of 5.5% required to achieve MDG-5 (WHO, 2015b). Part of the leading complications responsible for nearly 75% of all maternal deaths include; severe bleeding and high blood pressure (pre-eclampsia and eclampsia) (Say et al., 2014). Common modes of presentation of eclampsia in Nigeria are elevated blood pressure, convulsion, previous history of hypertension, epigastric pain and pedal oedema (Jimoh et al., 2007). Another Nigerian study reported a prevalence of 0.9% for eclampsia which occurred mostly (93.3%) among unregistered patients and mostly (78.5%) among women below the age of 30 (Orji et al., 2007). Incidentally, elevated blood pressure or hypertension which is the most common medical problem encountered in pregnancy is one of the criteria for the diagnosis of eclampsia which also includes convulsion and proteinuria (Nelson-Piercy, 2007). Hypertension is also a leading cause of perinatal and maternal morbidity and mortality, and pregnant women with this condition are more likely to develop placental abruption, Disseminated Intravascular Coagulopathy (DIC), cerebral haemorrhage, hepatic

failure and acute renal failure (Report, 2000). A large proportion of maternal deaths, especially in Northern Nigeria, are associated with obstetric complications including haemorrhage and eclampsia (UNICEF, 2008). In Ethiopia, the level of knowledge of Obstetric Danger Signs (ODSs) during pregnancy, childbirth and postpartum period was low and in Uganda, the prevalence of recently delivered women who had knowledge of key danger signs, or those who were birth prepared was also very low (Bogale and Markos, 2015). While findings indicate a poor perception of common and serious ODSs in Pakistan, an Indian study reported that 90.5, 80.0 and 78.4% of pregnant women were able to recognize bleeding/leaking per vagina, decreased/absent fetal movements and seizures/fits during pregnancy as ODSs, respectively (Hasan and Nisar, 2002; Sangal et al., 2013).

However, most studies were health facility based, conducted mainly among those who were already pregnant or recently delivered and did not take into consideration non-pregnant and nulliparous women. Moreover, almost all the studies were conducted in the hinterland, far away from densely-populated Atlantic Ocean coastline. Furthermore, all women in these studies were lumped together as one category and not differentiated into various reproductive age groups. In addition, some of these studies focused mainly on medical modification of the damage already done, instead of addressing the root causes of the problem. Finally, very few studies in Africa consider a wide range of ODSs, but focused on either one or two. Data on perception of danger signs associated with pregnancy among coastal populations is very scanty leading to little knowledge of the exact figure of maternal morbidity and mortality in this geographical location. This study therefore aimed to explore the risk factors for perception of obstetric complications, assess the level of perception of women in various reproductive age groups and determine the respective roles of educational level and parity on the perception of identified ODSs. The test hypotheses were that the age, marital status, educational level, previous miscarriage and knowing a woman who died in pregnancy were significant risk factors in the perception of ODSs among the study group.

MATERIALS AND METHODS

This descriptive study compared perception of seven selected ODSs; vaginal bleeding, convulsion, high blood pressure, severe malaria, cessation of fetal movement, swollen legs, pale hands and feet among women in different childbearing age groups, taking into consideration their marital status, educational level and parity. Literature search revealed no similar study having been conducted in this geographical location in the country and extremely few in other parts similar location in sSA. The study population was defined as the homogenous group of people living together for at least 10 years at Elegushi Community in Ibeju-Lekki Local Government Area (LGA) and Ijede Community at Ikorodu LGA of Lagos State, Nigeria (Afolabi et al., 2015). In general there was a

high literacy rate of 88.4% in Lagos State and 91.6% among those aged 15 to 49 years (NPC, 2006). Although the 2006 population census of Ibeju-Lekki and of Ikorodu LGAs were recorded as 99,540 and 689,045, respectively by 2013 when this study took place, there was an increase in the population due to natural annual increase and rural-urban drift (Lagos State Government, 2005). The approximate population of Elegushi community and that of Ijede were 30,350 and 88,000, respectively, of which approximately 25% were women of child-bearing age, based on the census. Therefore, the sampling fraction (sample size 845; and population size 25%) to be selected for the study was calculated to be approximately 1 out of every 35 women in reproductive age. Each of these two communities is further subdivided into wards and further still, into different families living in either rented or owned houses. The populations were mostly traders with some who were government workers and farmers. Respondents were divided into three age groups and categorized into three corresponding stages of reproductive age. As shown in Figure 1, respondents below the age of 25 were classified as early reproductive age (ERA), those between 25 and 34 years as mid reproductive age (MRA), and those 35 years and above as late reproductive age (LRA).

Ikorodu, located at approximately 30 km from metropolitan Lagos, is a semi-urban town sited close to the Atlantic Ocean. It is separated from Ibeju-Lekki LGA by a large body of inland water, the Lagos Lagoon. Ikorodu shares its northern boundary with Ogun State and its eastern and western borders with Epe and Kosofe LGAs, respectively. It is a traditional settlement of the Aworis, a sub-group of the Yorubas. A modern 10-lane expressway has been newly built to link metropolitan Lagos to Ikorodu, thus enlarging its population. There are thermal stations, mini ports, and institutions of higher learning at Ikorodu. Ibeju-Lekki sits next to the Atlantic Ocean. It is more of a resort center with sea-side hotels and tourist center. The climate is generally coastal, though the rainy and dry seasons follow their regular pattern. Luscious and verdant vegetation surround these communities that have minimal air pollution. Houses are made of bricks, roads are mostly tarred and the communities have private and public schools and hospital systems. Most houses are connected to the electricity grid, though electricity is not regular. Housing and sewage systems however need to be improved. The State government conducts regular indoor residual spraying and occasionally gives free Long-Lasting Insecticidal Nets (LLINs) and free malaria treatment to indigenes.

The target population was women in different stages of childbearing age residing in these communities. Measured independent variables in the study were marital status, highest level of education, religion, and ethnic group. Perception was assessed by asking if the respondents regard each of the medical conditions described above as a danger sign in pregnancy. Epi Info 7 was used to calculate the sample size for the projected WCBA population of each site of study, using 30% as an expected frequency of correct perception of obstetric danger signs in each of the communities, a confidence limit of 5%, and confidence level set at 95%. The calculated sample size was 641 (319 for Ibeju-Lekki and 322 for Ikorodu). For contingencies, especially the considerations of attrition, non-response and increase in representativeness, the sample size was increased by 30%, bringing it to 833 and rounding the figures to 845. Systematic random sampling was used to select streets at regular intervals from the list of streets in each community. From the selected streets, the starting point of the survey was the first house in a particular direction and by simple random sampling every fourth house was selected for the study. If a selected house did not have any WCBA, then the next house was visited to replace it for a respondent. All the WCBA (15 to 49 years) who agreed to participate were interviewed. Only women aged between 15 and 49 years, resident within the communities of study and not just visiting, of sound mind and not mentally compromised were included in the study. Those excluded were women hospitalized for any reason,

those moribund, visitors and tourists, Caucasians and commercial sex workers.

Marital status was defined as "single" if the participant had never had a conjugal relationship or "married" if there had been or still was a conjugal relationship. Educational level meant the stage of education where each respondent was or stopped at the time of the study. Parity meant the number of times the respondent has delivered a child. The information was collected using pretested semi-structured questionnaires which were electronically prepared on a Mobile Data Capturing Device (MDCD). The electronic questionnaire was divided into six parts consisting of administrative information; demographic information; knowledge of Human Immunodeficiency Virus (HIV)/tuberculosis/malaria; contraceptive use; self-assessment of individual health; and in the last part of the electronic questionnaire, the reports of testing for hemoglobin concentration, malaria parasitemia, and HIV. It took roughly 45 min to complete a questionnaire in a face-to-face interview at household level. Six field workers were trained in administering the electronic questionnaire. Once data collection was finalized, questionnaires were electronically sent to a central location where they were automatically recorded in an Excel spreadsheet. The collected data were analyzed using STATA 13. Age distribution was assumed to be approximately normal, following a Gaussian curve. Two outliers in age were detected which prompted revisiting the respondents and correcting the ages which were due to transpositions. Frequency tables were used for categorical variables. Cross-tabulations and chi-square tests were used for appropriate variables such as age group and ANC, marital status and awareness of ODS. Confidence level ($1-\alpha$) was set at 95% and power ($1-\beta$) at 80%. Appropriate test statistics were applied and the p-value (<0.05) was considered statistically significant. Ethical considerations included taking verbal, informed consent from all the participants. The confidentiality of the data was ensured and clearance was obtained from Lagos State Government ethics committee. The survey was anonymous in order to preserve the privacy of study subjects.

RESULTS

Initially, 859 open-ended, semi-structured questionnaires were prepared for the communities of study but 6 of these were technically incomplete and had to be excluded from further analysis. Five respondents travelled out of their residences during the study and their participation had to be terminated. Another 3 women rejected to respond further based on both religious or cultural beliefs and 2 refused to participate at all, giving a response rate of 98.1%. Table 1 shows that the 829 respondents included 249 (30.0%) women in ERA, 410 (49.5%) in MRA and 170 (20.5%) in LRA. Demographic characteristics of the respondents were skewed more towards 662 (79.9%) married women; 679 (81.9%) secondary education (679, 81.9%); 675 (81.4%) Yoruba ethnic group and 347 (41.85) parity of 1 to 2. Only 52 (6.3%) of the respondents had no formal education and 3 (0.4%) had other forms of education such as attending Koranic schools. There were 478 (57.7%) Christians, 346 (41.7%) Muslims and 5 (0.6%) Traditionalists. Figure 1 shows the distribution of parity among the respondents. In all, 384 (51.4%) women in MRA, 335 (44.8%) of those with parity 1 to 2 and 614 (82.1%) of those who had secondary education ever used Antenatal care (ANC). A

Table 1. Socio-demographic characteristics of respondents.

| Variable | Item | Frequency | Percent |
|-------------------|---------------------|-----------|---------|
| Age group (years) | <25 | 249 | 30.0 |
| | 25-34 | 410 | 49.5 |
| | ≥35 | 170 | 20.5 |
| Marital status | Single | 167 | 20.1 |
| | Married | 662 | 79.9 |
| Educational level | No formal education | 52 | 6.3 |
| | Secondary | 679 | 81.9 |
| | Undergraduate | 6 | 0.7 |
| | Higher degree | 50 | 6.0 |
| | Diploma | 39 | 4.7 |
| | Others | 3 | 0.4 |
| Religion | Christianity | 478 | 57.7 |
| | Islam | 346 | 41.7 |
| | Traditional | 5 | 0.6 |
| Ethnicity | Yoruba | 675 | 81.4 |
| | Ibo | 72 | 8.7 |
| | Hausa | 2 | 0.2 |
| | Others | 80 | 9.7 |
| Parity | 0 | 182 | 22.0 |
| | 1-2 | 347 | 41.8 |
| | 3-4 | 232 | 28.0 |
| | 5-6 | 59 | 7.1 |
| | 7-8 | 9 | 1.1 |

similar proportion among these groups was observed for those who considered that ANC as important in pregnancy (Table 2). As shown in Table 3, women in MRA were over three times more likely to visit ANC ($X^2=23.43$; $P=0.000002$, $OR=3.34$; 95% $CI=2.01, 5.57$) compared to those in ERA. Women in LRA were over four times more likely to visit ANC when matched with those in ERA. There was no statistically significant difference ($X^2=0.23$; $P=0.63$, $OR=1.21$; 95% $CI=0.56, 2.64$) in the proportion of women in LRA who visited ANC compared to those in MRA (Figure 2).

Figures 3 and 4 presents a graphical illustration of perception of ODSs by women in different age groups clearly showing that a large proportion of the women have low perception of these ODSs. As illustrated in Table 4, higher proportions of women in MRA had better perception of ODS than those in ERA or LRA. However, that proportion was statistically significant only for vaginal bleeding ($X^2=4.31$; $P=0.037$, $OR=1.40$; 95% $CI=1.02, 1.92$), severe malaria ($X^2=3.67$; $P=0.05$, $OR=1.38$; 95% $CI=0.99, 1.91$), pale hands and feet ($X^2=4.43$; $P=0.03$,

$OR=1.71$; 95% $CI=1.03, 2.85$) when MRA group was compared with ERA group; and only for convulsion ($X^2=4.75$, $p=0.03$, $OR=1.81$, 95% $CI=1.06, 3.10$) when MRA group was compared with LRA group. Also, single women were more than twice likely to disregard vaginal bleeding ($X^2=17.0$; $P=0.000043$, $OR=2.09$; 95% $CI=1.46, 2.98$) and about twice as likely to also disregard convulsion ($X^2=5.05$; $P=0.02$, $OR=1.80$; 95% $CI=1.10, 3.00$) as ODSs. In general, there was no significant difference in the risks of an ODS among the respondents. Nevertheless, among those who disagreed that bleeding is an ODS, risk of vaginal bleeding during pregnancy was slightly higher among women in ERA compared with MRA ($RR=1.17$, 95% $CI=1.0, 1.9$) and among those who were singles compared to those who were married ($RR=1.37$, 96% $CI=1.2, 1.7$).

As shown in Tables 5 and 6, there were significant associations (positive and negative, respectively) between knowing a woman who died in pregnancy and perception of bleeding as an ODS ($t=3.72$, $P=0.000$, 95% $CI=0.07, 0.22$) on one hand and perception of pale hands




Figure 1. Frequency distribution of number of deliveries among rural women in Lagos, Nigeria.

Table 2. Utilization and perceived importance of antenatal care by respondents relative to age group (years), parity and educational status.


| Variable | Item | Ever used ANC | | ANC is important in pregnancy | | |
|-------------------|---------------|---------------|-----------|-------------------------------|-----------|------------|
| | | Yes | No | Yes | No | Don't know |
| | | Freq. (%) | Freq. (%) | Freq. (%) | Freq. (%) | Freq. (%) |
| Age group | <25 | 203 (27.1) | 46 (56.8) | 205 (28.1) | 8 (44.4) | 36 (50.0) |
| | 25-34 | 384 (51.4) | 26 (32.1) | 384 (52.7) | 6 (33.3) | 20 (27.8) |
| | ≥35 | 161 (21.5) | 9 (11.1) | 150 (20.6) | 4 (22.2) | 16 (22.2) |
| Parity | 0 | 126 (16.8) | 56 (69.1) | 136 (18.4) | 8 (44.4) | 38 (52.8) |
| | 1-2 | 335 (44.8) | 12 (14.8) | 327 (44.3) | 6 (33.3) | 14 (19.4) |
| | 3-4 | 226 (30.2) | 6 (7.4) | 214 (29.0) | 4 (22.2) | 14 (19.4) |
| | 5-6 | 53 (7.1) | 6 (7.4) | 55 (7.4) | 0 (0.0) | 4 (5.6) |
| | 7-8 | 8 (1.1) | 1 (1.2) | 7 (0.9) | 0 (0.0) | 2 (2.8) |
| Educational level | No formal | 47 (6.3) | 5 (6.3) | 49 (6.6) | 1 (5.6) | 2 (2.8) |
| | Secondary | 614 (82.1) | 65 (80.2) | 600 (81.2) | 15 (83.3) | 64 (88.9) |
| | Undergraduate | 4 (0.5) | 2 (2.5) | 5 (0.7) | 0 (0.0) | 1 (1.4) |
| | Higher degree | 42 (5.6) | 8 (9.9) | 44 (6.0) | 2 (11.1) | 4 (5.6) |
| | Diploma | 38 (5.1) | 1 (1.2) | 38 (5.1) | 0 (0.0) | 1 (1.4) |
| | Others | 3 (0.4) | 0 (0.0) | 3 (0.4) | 0 (0.0) | 0 (0.0) |
| Total | - | 748 (90.2) | 81 (9.8) | 739 (89.4) | 18 (2.2) | 72 (8.7) |

and feet as an ODS ($t=-2.30$, P -value=0.02, 95% CI = -0.11, -0.01) on the other hand. This singular factor of knowing another woman who died in pregnancy was

responsible for 33% of the variation in perception of bleeding ($r=0.14$, $r^2=0.33$, $P=0.0001$) and 22% of the variation perception of pale hands and feet ($r = -0.06$, $r^2 =$

Table 3. Relationship between age and utilization of antenatal care.

| Age group (years) | Statistics | Age group (years) | | |
|-------------------|--------------------------|-------------------|-------------------|-------------------|
| | | <25 | 25-34 | ≥35 |
| <25 | X ² (P-value) | - | 23.43 (0.000002) | 15.39 (0.00009) |
| | OR (95% CI) | - | 3.34 (2.01, 5.57) | 4.05 (1.93, 8.53) |
| 25-34 | X ² (P-value) | 23.43 (0.00002) | - | 0.23 (0.63) |
| | OR (95% CI) | 0.30 (0.18, 0.50) | - | 1.21 (0.56, 2.64) |
| ≥35 | X ² (P-value) | 15.39 (0.00009) | 0.23 (0.63) | - |
| | OR (95% CI) | 0.25 (0.12, 0.52) | 0.83 (0.38, 1.80) | - |

**Figure 2.** Attitude of coastal women in various reproductive stages on antenatal care.

0.22, P-value=0.005) as ODSs. There was also a negative but significant association between age group and pale hands and feet ($t = -3.41$, P-value=0.001, 95% CI=-0.10, -0.03), accounting for a significant 22% of the variation in perception of pallor as an ODS ($r=-0.07$, $r^2=0.022$, P-value =0.005). Substantial but negative association was also observed between educational level and severe malaria ($t=-2.46$, P-value=0.14, 95% CI = -

0.06, -0.01) which explains an insignificant 14% of the variation in the said perception ($r= -0.04$, $r^2=0.014$, P=0.09). Marital status was significantly but negatively associated with cessation of fetal movement ($t=-2.86$, P-value =0.004, 95% CI=-0.09, -0.02) which explained a significant 15% of the variation in perception of cessation of fetal movement as an ODS ($r=-0.06$, $r^2=0.015$, P-value = 0.05).




Figure 3. Perception of obstetric danger signs according to age group of women in rural coastal lagos, Nigeria.




Figure 4. Perception of obstetric danger sign by marital status of women in coastal rural Lagos, Nigeria.

Table 4. Relative risks of the danger signs in pregnancy by age group (years) and marital status of women living in rural areas of Lagos.

| Age group | Vaginal bleeding | | Convulsion | | High Blood Pressure | | Severe malaria | | Cessation of fetal movement | | Swollen legs | | Pale hands and feet | |
|-----------------------|------------------|-----------------|------------------|-----------------|---------------------|-----------------|------------------|-----------------|-----------------------------|-----------------|------------------|-----------------|---------------------|-----------------|
| | Yes Freq. (%) | No Freq. (%) | Yes Freq. (%) | No Freq. (%) | Yes Freq. (%) | No Freq. (%) | Yes Freq. (%) | No Freq. (%) | Yes Freq. (%) | No Freq. (%) | Yes Freq. (%) | No Freq. (%) | Yes Freq. (%) | No Freq. (%) |
| <25 | 108 (26.9) | 141 (32.9) | 48 (33.6) | 201 (29.3) | 35 (27.3) | 214 (30.5) | 84(25.8) | 165(32.8) | 19 (26.4) | 230 (31.2) | 57 (25.9) | 192 (31.5) | 23 (20.0) | 226 (31.7) |
| 25-34 | 212 (52.9) | 198 (46.3) | 76 (53.1) | 334 (48.7) | 69 (53.9) | 341 (48.7) | 169 (52.0) | 241 (47.8) | 43 (59.7) | 367 (48.5) | 114 (51.8) | 296 (48.6) | 61 (53.0) | 349 (48.9) |
| χ^2 (p-value) | 4.31 (0.04) | | 0.06 (0.81) | | 0.90 (0.34) | | 3.67 (0.05) | | 1.48 (0.22) | | 1.94 (0.16) | | 4.43 (0.04) | |
| RR (95% CI) | 1.17 (1.0, 1.9) | | 0.99 (0.9, 1.1) | | 1.03 (1.0, 1.1) | | 1.13 (1.0, 1.3) | | 1.03 (1.0, 1.1) | | 1.07 (1.0, 1.2) | | 1.07 (1.0, 1.1) | |
| ≥ 35 | 81 (20.2) | 89 (20.8) | 19 (13.3) | 151 (22.0) | 24 (18.8) | 146 (20.8) | 72 (22.2) | 98 (19.4) | 10 (13.9) | 160 (21.7) | 49 (22.3) | 121 (19.9) | 31 (27.0) | 139 (19.4) |
| χ^2 (p-value) | 0.75 (0.39) | | 4.94 (0.03) | | 0.00 (0.98) | | 3.21 (0.07) | | 0.48 (0.49) | | 1.88 (0.17) | | 7.29 (0.007) | |
| RR (95% CI) | 1.08 (0.9, 1.3) | | 0.91 (0.8, 1.0) | | 1.00 (0.9, 1.1) | | 1.15 (1.0, 1.3) | | 0.98 (0.9, 1.0) | | 1.1 (1.0, 1.2) | | 1.1 (1.0, 1.2) | |
| χ^2 (p-value) | 0.79 (0.37) | | 4.75 (0.03) | | 0.66 (0.42) | | 0.06 (0.80) | | 3.07 (0.08) | | 0.06 (0.80) | | 1.01 (0.31) | |
| RR (95% CI) | 0.92 (0.8, 1.1) | | 0.92 (0.9, 1.0) | | 0.97 (0.9, 1.0) | | 1.02 (0.9, 1.2) | | 0.95 (0.9, 1.0) | | 1.01 (0.9, 1.1) | | 1.0 (1.0, 1.1) | |
| Marital status | | | | | | | | | | | | | | |
| Single | 57 (14.2) | 110 (25.7) | 19 (13.3) | 148 (21.6) | 23 (18.0) | 144 (20.5) | 54 (16.6) | 113 (22.4) | 12 (16.7) | 155 (20.5) | 42 (19.1) | 125 (20.5) | 26 (22.6) | 141 (19.8) |
| Married | 344 (85.8) | 318 (74.3) | 124 (86.7) | 538 (78.4) | 105 (82.0) | 557 (79.5) | 271 (83.4) | 391 (77.6) | 60 (83.3) | 602 (79.5) | 178 (80.9) | 484 (79.5) | 89 (77.4) | 573 (80.2) |
| χ^2 (p-value) | 17.0 (0.00004) | | 5.05 (0.02) | | 0.45 (0.50) | | 4.14 (0.04) | | 0.59 (0.44) | | 0.21(0.65) | | 0.50 (0.48) | |
| RR (95% CI) | 1.37 (1.2, 1.7) | | 1.09 (1.0, 1.2) | | 1.02 (1.0, 1.1) | | 1.15 (1.0, 1.3) | | 1.02 (1.0, 1.1) | | 1.02 (0.9, 1.1) | | 0.98 (0.9, 1.0) | |
| Total | 401 (48.4) | 428 (51.6) | 143 (17.3) | 686 (82.7) | 128 (15.4) | 701 (84.6) | 325 (39.2) | 504 (60.8) | 72 (8.7) | 757 (91.3) | 220 (26.5) | 609 (73.5) | 115 (13.9) | 714 (86.1) |

Freq: Frequency.

DISCUSSION

Keen perception of obstetric danger signs is the critical and indispensable initial stage in being compliant and accepting appropriate and timely referral to obstetric and newborn care. ODSs are global problems that have not been satisfactorily addressed, especially in developing countries. From the community perspective, both awareness of the danger signs of an obstetric complication and knowledge about their severity are important.

Community members, especially women should therefore know when to take applicable lifesaving action for each of the complications. About 90% of the respondents had utilized ANC at least once, a figure higher than the 69% reported for sSA (Lincetto et al., 2006).

For 2.2% of respondents, ANC was not important during pregnancy and another 8.7% were not sure of the importance of ANC. Women who did not attend ANC were significantly less likely to be aware of both critical and non-critical

pregnancy complications when compared with those who do not know any complications (Doctor et al., 2013). Furthermore, antenatal period clearly presents opportunities for reaching pregnant women with a number of interventions that may be vital to their health and well-being and that of their infants. It is a window of opportunity to inform women and families about danger signs and symptoms and about the risks of labour and delivery. It also provides the route for ensuring that pregnant women do in practice, deliver with

Table 5. Multiple regression analyses on associations between ODSs as dependent variables and other independent variables.

| | Observed | R ² | F | P |
|-----------------------------|----------|----------------|-------|--------|
| Bleeding | 829 | 0.033 | 4.59 | 0.0001 |
| Convulsion | 829 | 0.010 | 1.34 | 0.24 |
| High Blood Pressure | 829 | 0.006 | 0.79 | 0.58 |
| Severe malaria | 829 | 0.014 | 1.84 | 0.09 |
| Cessation of fetal movement | 829 | 0.015 | 2.03 | 0.05 |
| Swollen feet | 829 | 0.019 | 2.63 | 0.02 |
| Pale hands and feet | 829 | 0.022 | 3.140 | 0.005 |

Table 6. Correlation coefficient outcomes between ODSs as dependent variables and other independent variable.

| Parameter | ODS | Coefficient (r) | Standard error | t | P-value | 95% Confidence interval |
|-------------------------------------|-----------------------------|-----------------|----------------|-------|---------|-------------------------|
| Knows a woman who died in Pregnancy | Bleeding in pregnancy | 0.14 | 0.04 | 3.72 | 0.00 | 0.07, 0.22 |
| | Pale hands and feet | -0.06 | 0.03 | -2.30 | 0.02 | -0.11, -0.01 |
| Age group (years) | Convulsion | 0.05 | 0.02 | 2.29 | 0.023 | 0.01, 0.09 |
| | Pale hands and feet | -0.07 | 0.02 | -3.41 | 0.001 | -0.10, -0.03 |
| Educational level | Severe malaria | -0.04 | 0.01 | -2.46 | 0.014 | -0.06, -0.01 |
| | Swollen feet | -0.03 | 0.01 | -2.58 | 0.010 | -0.06, -0.01 |
| Marital status | Cessation of fetal movement | -0.06 | 0.02 | -2.86 | 0.004 | -0.09, -0.02 |

the assistance of a skilled health care provider (AbouZahr and Wardlaw, 2003). Factors that could be responsible for not seeing ANC as important might be related to previous experiences of respondents themselves, or of relatives and friends. These factors may include services provided at ANC, the environment where the health facility is located, difficulty in accessing the health facility, health staff unfriendly attitude, competing Traditional Birth Attendants and stock-outs of essential medicine. In addition, there have been reports that cost, in the form of transportation to and fro, as well as out-of-pocket spending may mitigate against appropriate utilization of ANC.

Similar to results from studies on women in Nigeria and in other parts of the world, respondents in this study demonstrated low perception of ODSs (Bogale and Markos, 2015; Kabakyenga et al., 2011; Hasan and Nisar, 2002; Nambala and Ngoma, 2013). For example, only 48.4% of respondents recognized vaginal bleeding as a danger sign in pregnancy, compared to 90.5% reported from India (Sangal et al., 2013). Multiple factors may contribute to this finding. First, this danger sign is relatively easy to recognize. Second, bleeding is common in women in reproductive age regardless of whether they are pregnant or not. Additionally, bleeding in pregnancy

may be provoked and not directly linked to the pregnancy itself such as post-coital bleeding of cervical origin which may be due to infection or malignancy. Finally, bleeding with tissues, especially in early pregnancy, may be as a result of spontaneous (or illegally induced) abortion (Jager, 2015). Vaginal bleeding after mid-pregnancy is associated with maternal and fetal risks. Bleeding at this time often results from placenta previa, abruptio placenta and vasa previa (Sakornbut et al., 2015).

Similar to the report from Pakistan, about 90% of respondents perceived cessation of fetal movement as not a danger sign (Hasan and Nisar, 2002). Fetal movement, a marker of fetal well-being, is a relatively simple indirect means of fetal assessment that indicates a lack of significant acidosis. With fetal monitoring, movement in healthy fetuses increases sympathetic response and accelerates heart rate. Fetuses with severe acid-base disorders cannot oxygenate their muscles adequately and do not move (Warren, 2009). Nearly 50% of late-pregnancy intra-uterine fetal deaths have no associated risk factors, though fetal demise may however be heralded by decreased fetal movement followed by cessation of movement at least 12 h before death (Institute for Clinical Systems Improvement, 2008). In view of this, Harrington et al. (1998) contend that maternal

perception of decreased fetal movement is not associated with poor fetal outcome.

Convulsion, high blood pressure and swollen feet, taken together are common precursors of pre-eclampsia, of which over 80% of respondents were not aware of it as danger signs in pregnancy. About 50% of women aged 25 to 34 years demonstrated low perception of this trio as ODSs. Eclampsia was found more in age group 20 to 29 years which accounted for 60.5% of eclamptic patients at Ilorin in Nigeria (Jimoh et al., 2007) and occurring more in primigravidae and more during wet season than dry season (Orji et al., 2007). Since there is currently no known measure to effectively and consistently prevent pre-eclampsia, early detection is critical in reducing the morbidity and mortality due to this condition (Duley, 2003; Hofmeyr et al., 2003; Vainia et al., 2002).

High blood pressure in pregnancy is definitely detrimental to the mother separately, the fetus separately and both the mother and the fetus collectively. Hypertension could predate pregnancy, be associated with pregnancy or occur at post-partum period. Pregnant women should be made aware of the need to seek immediate advice from health care professionals if they experience symptoms such as severe headache, problems with vision (blurring or flashing before the eyes, severe pain just below the ribs, vomiting and sudden swelling of the face, hands or feet (National Institute for Health, 2015).

Strengths and weaknesses

The sample size was large enough; thereby improving external validity and statistical inferences. Weaknesses of the study include the absence of qualitative methods, such as focused group discussions, which would have better addressed some questions raised. Some of the respondents may be experiencing pregnancy for the first time and may not be too familiar with the appropriate responses to the questions raised. Occasional rains and a community disturbance disrupted field works briefly.

CONCLUSION

Accurate perception of ODS is a critical step in minimizing the high levels of reproductive morbidity and mortality in Nigeria. Since knowledge of ODSs is largely absent among pregnant women, attempts must be made to increase their knowledge and improve their perception through mass media announcement, Information, Education and Communication (IEC) as well as Behavioral Change Communication (BCC), activities and regular visits by Community Health Workers (CHWs) and other healthcare providers. Women education, from secondary level, is meaningfully related with level of knowledge about ODSs and also with health seeking incase these danger signs appear (Sandal et al., 2012).

RECOMMENDATIONS

The relevance of ANC as well as of Community Health Workers (CHWs) in empowering women with adequate knowledge on ODSs cannot be over-emphasized. While ANC is one of the windows of opportunity for women in child-bearing age in Nigeria and other developing economies to be empowered on ODSs, other windows of opportunity are the home through radio jingles and television programs; market places where flyers with pictorial diagrams and hand bills can be distributed; schools where female students at secondary level can be taught about their reproductive system; and at religious gatherings, saloons, village meetings and peer-group meetings. CHWs, meanwhile can provide education on ODSs since community perception on illness and health care can lead to pathways for better health seeking behavior (Rashid, 2001; Watkins and Plant, 2004).

Monitoring and evaluation programs should pool awareness of pregnancy danger signs with other indicators measuring related aspects of knowledge and behavior to evaluate the real impact of any awareness-raising program. Matching indicators may comprise if necessary, the proportion of the population who can identify the site of emergency obstetric services and the proportion of the population who anticipate using these services in the event of emergency.

Further studies should assess similar research topic in other rural coastal areas of Atlantic and Indian Oceans and possible the Mediterranean Sea. Future studies should consider using quantitative and qualitative techniques and hospital records for triangulation.

ACKNOWLEDGEMENT

This study is one of the projects sponsored by the Federal Ministry of Science and Technology Education Post-Basic (STEP-B) of Nigeria, assisted by the World Bank.

Conflict of interest

Authors have none to declare.

REFERENCES

- AbouZahr C, Wardlaw T (2003). Antenatal Care in Developing Countries: Promises, Achievements and Missed Opportunities. WHO and UNICEF pp. 1-37.
- Adam T, Franz-Vasdeki J (2013). A Quantitative Mapping of Trends in Reductions of Maternal and Child Mortality in the High Mortality-burden Countdown to 2015 countries. Alliance for Health Policy and Systems Research and the Partnership for Maternal, Newborn and Child Health pp. 1-32.
- Afolabi BM, Ogunwale A, Ajibade AA (2015). Knowledge, non-use, use and source of information on contraceptive methods among women in various stages of reproductive age in rural Lagos, Southwest

- Nigeria. *Clin. Ophthalmol.* 9:885-887.
- Bogale D, Markos D (2015). Knowledge of obstetric danger signs among childbearing age women in Goba district, Ethiopia: a cross-sectional study. *BMC Pregnancy Childbirth* 15:77.
- Doctor HV, Findley SE, Cometto G (2013). Women's awareness of obstetric complications, preparations for delivery and utilization of skilled birth attendants in Northern Nigeria. Available at: <http://paa2011.princeton.edu/abstracts/110256>
- Duley L (2003). Pre-eclampsia and the hypertensive disorders in pregnancy. *Br. Med. Bull.* 67:161-76.
- Grown C, Gupta GR, Pande R (2005). Taking action to improve women's health through gender equality and women's empowerment. *Lancet* 365(9458):541-543.
- Harrington K, Thompson O, Jordan L, Page J, Carpenter RG, Campbell S (1998). Obstetric outcome in women who present with a reduction in fetal movements in the third trimester of pregnancy. *J. Perinat. Med.* 26(2):77-82.
- Hasan IJ, Nisar N (2002). Womens' Perceptions regarding Obstetric Complications and Care in a Poor Fishing Community in Karachi. *J. Pak. Med. Assoc.* 52:148
- Hofmeyr GJ, Roodt A, Atallah AN, Duley L (2003). Calcium supplementation to prevent pre-eclampsia: a systematic review. *S. Afr. Med. J.* 93(3):224-8.
- Institute for Clinical systems Improvement (2008). Routine Prenatal Care. 12th ed. August. Available at: http://www.icsi.org/prenatal_care_4/prenatal_care_routine__full_version__2.html
- Jager B (2015). Vaginal bleeding in the first trimester of pregnancy. Available at: https://www.glowm.com/pdf/Chap-02_Jager.pdf.
- Jimoh AAG, Akintade OA, Balogun OR, Aboyeji AP (2007). Eclampsia – A ten Year Review in a Nigerian Teaching Hospital. *Nig. Hosp. Pract.* 1(3):80-3.
- Kabakyenga JK, Östergren P, Turyakira E, Pettersson KO (2011). Knowledge of obstetric danger signs and birth preparedness practices among women in rural Uganda. *Reprod. Health* 8:33.
- Lagos State Government (2005). Lagos State Bureau of Statistics. Population. Available at: <http://www.lagosstate.gov.ng/pagelinks.php?p=6>.
- Lincetto O, Mothebesoane-Anoh S, Gomez P, Munjanja SP (2006). Antenatal care. *In Opportunities for Africa's newborns: Practical data, policy and programmatic support for newborn care in Africa.* Edited by Joy L, Kate K. PMNCH, Cape Town. Available at: <http://www.who.int/pmnch/media/publications/africannewborns/en/>
- Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, Abraham J, Adair T, Aggarwal R (2012). Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380:2095–2128.
- Lozano R, Wang H, Foreman KJ, Rajaratnam JK, Naghavi M, Marcus JR (2011). Progress towards Millennium Development Goals 4 and 5 on maternal and child mortality: an updated systematic analysis. *Lancet* pp. 378:9797, 1139-1165.
- Nambala BS, Ngoma C (2013). Knowledge and Perception of Women Towards Danger Signs in Pregnancy in Choma Rural District, Zambia. *M. J. Zambia* 40(2):43-7.
- National Institute for Health and Clinical Excellence (2015). Hypertension in pregnancy. NICE clinical guideline 107. Available at: <https://www.nice.org.uk/guidance/cg107>
- Nelson-Piercy C (2007). *Handbook of Obstetric Medicine.* 3rd ed: Informa Health Care.
- Orji EO, Olabode TO, Orji VO (2007). Making Motherhood Safer in Eclamptic Patients. *Ibom Med. J.* 2(1):1-7.
- Rashid SF (2001). Indigenous notions of the workings of the body: conflicts and dilemmas with Norplant use in rural Bangladesh. *Qual. Health Res.* 11(1):85-102.
- Report of the National High Blood Pressure (2000). Education Program Working Group on High Blood Pressure in Pregnancy. *Am. J. Obstet. Gynecol.* 183(1):S1-S22.
- Sakornbut E, Leeman L, Fontaine P (2015). Late Pregnancy Bleeding. Available at: www.aafp.org/afp/2007/0415/p1199.
- Sangal R, Srivastava R, Singh K, Srivastava DK, Khan H (2013). Knowledge on Components of ANC Care & their Utilization in Women attending ANC Clinic at BRD Medical College, Gorakhpur.. *JARBS.* 5(2): 159-163.
- Say L, Chou D, Gemmill A, Tunçalp Ö, Moller AB, Daniels J (2014). Global Causes of Maternal Death: A WHO Systematic Analysis. *Lancet Glob. Health* 2:e323-e333.
- UNICEF (2008). *The State of the World's Children 2009.* New York: United Nations Children's Fund.
- UNFPA (2012). UNICEF, WHO, World Bank. Trends in Maternal Mortality 1990-2010. Available at: <http://www.unfpa.org/public/home/publications/pid/10728>.
- United Nations (2003). *World fertility report 2003.* New York: UN Department of Economic and Social Affairs Population Division.
- Vainia M, Kujansu E, Iso-Mustajarvi M, Maenpaa J (2002). Low dose acetylsalicylic acid in prevention of pregnancy induced hypertension and intrauterine growth retardation in women with bilateral uterine artery notches. *Br. J. Obstet. Gynaecol.* 109(2):161-7
- Warren J (2009). Clinical commentary. *In: Darby-Stewart A, Strickland C, Jamieson B. Do abnormal fetal kick counts predict intrauterine death in average-risk pregnancies? Clin. Inquir.* 58(4):220a-220c.
- Watkins RE, Plant AJ (2004). Pathways to treatment for tuberculosis in Bali: patient perspectives. *Qual. Health Res.* 14(5):691-703.
- World Health Organization (2015b). Medical centre. Available at: www.who.int/mediacenter/factsheets/fs348/en/.
- World Health Organization (2015a). MDG 5: Improve Maternal Health. Available at: http://www.who.int/topics/millennium_development_goals/maternal_health/en/



Journal of Public Health and Epidemiology

Related Journals Published by Academic Journals

Journal of Diabetes and Endocrinology

Journal of Medical Genetics and Genomics

Journal of Medical Laboratory and Diagnosis

Journal of Physiology and Pathophysiology

Medical Practice and Reviews

Research in Pharmaceutical Biotechnology

academicJournals