

The background of the cover is a photograph of various mathematical manipulatives scattered on a light-colored wooden surface. These include white plastic blocks with numbers and symbols, colorful geometric shapes like triangles and circles, and a white ruler. The image is framed with rounded corners.

African Journal of Mathematics and Computer Science Research

Volume 9 Number 2 July 2016

ISSN 2006-9731



ABOUT AJMCSR

The African Journal of Mathematics and Computer Science Research (ISSN 2006-9731) is published bi-monthly (one volume per year) by Academic Journals.

The African Journal of Mathematics and Computer Science Research (AJMCSR) (ISSN:2006-9731) is an open access journal that publishes high-quality solicited and unsolicited articles, in all areas of the subject such as Mathematical model of ecological disturbances, properties of upper fuzzy order, Monte Carlo forecast of production using nonlinear econometric models, Mathematical model of homogenous tumour growth, Asymptotic behavior of solutions of nonlinear delay differential equations with impulse etc. All articles published in AJMCSR are peer-reviewed.

Contact Us

Editorial Office: ajmcsr@academicjournals.org

Help Desk: helpdesk@academicjournals.org

Website: <http://www.academicjournals.org/journal/AJMCSR>

Submit manuscript online <http://ms.academicjournals.me/>

Editors

Prof. Mohamed Ali Toumi

*Département de Mahtématiques
Faculté des Sciences de Bizerte
7021, Zarzouna, Bizerte
Tunisia.*

Associate Professor Kai-Long Hsiao,

*Department of Digital Entertainment,
and Game Design,
Taiwan Shoufu University,
Taiwan,
R. O. C.*

Dr. Marek Galewski

*Faculty of Mathematics and Computer Science,
Lodz University
Poland.*

Prof. Xianyi Li

*College of Mathematics and Computational Science
Shenzhen University
Shenzhen City
Guangdong Province
P. R. China.*

Editorial Board

Dr. Rauf, Kamilu

*Department of mathematics,
University of Ilorin,
Ilorin, Nigeria.*

Dr. Adewara, Adedayo Amos

*Department of Statistics,
University of Ilorin.
Ilorin.
Kwara State.
Nigeria.*

Dr. Johnson Oladele Fatokun,

*Department of Mathematical Sciences
Nasarawa State University, Keffi.
P. M. B. 1022, Keffi. Nigeria.*

Dr. János Toth

*Department of Mathematical Analysis,
Budapest University of Technology and
Economics.*

Professor Aparajita Ojha,

*Computer Science and Engineering,
PDPM Indian Institute of Information
Technology,
Design and Manufacturing, IT Building,
JEC Campus, Ranjhi, Jabalpur 482 011 (India).*

Dr. Elsayed Elrifai,

*Mathematics Department,
Faculty of Science,
Mansoura University,
Mansoura, 35516, Egypt.*

Prof. Reuben O. Ayeni,

*Department of Mathematics,
Ladoke Akintola University,
Ogbomoso,
Nigeria.*

Dr. B. O. Osu,

*Department of Mathematics,
Abia State University,
P. M. B. 2000,
Uturu,
Nigeria.*

Dr. Nwabueze Joy Chioma,

*Abia State University,
Department of Statistics,
Uturu,
Abia State,
Nigeria.*

Dr. Marchin Papzhytski,

*Systems Research Institute,
Polish Academy of Science,
ul. Newelska 6 0-60-66-121-66,
03-815 Warszawa,
POLAND.*

Amjad D. Al-Nasser,

*Department of Statistics,
Faculty of Science, Yarmouk University,
21163 Irbid,
Jordan.*

Prof. Mohammed A. Qazi,

*Department of Mathematics,
Tuskegee University,
Tuskegee,
Alabama 36088,
USA.*

Professor Gui Lu Long

*Dept. of Physics,
Tsinghua University,
Beijing 100084,
P. R. China.*

Prof. A. A. Mamun, Ph. D.

*Ruhr-Universitaet Bochum,
Institut fuer Theoretische Physik IV,
Fakultaet fuer Phyik und Astronomie,
Bochum-44780,
Germany.*

Prof. A. A. Mamun, Ph. D.

*Ruhr-Universitaet Bochum,
Institut fuer Theoretische Physik IV,
Fakultaet fuer Phyik und Astronomie,
Bochum-44780,
Germany.*

Short Communication

Properties of the Euler phi-function on pairs of positive integers $(6x - 1, 6x + 1)$

Mbakiso Fix Mothebe* and Sebaameng Samuel

Department of Mathematics, University of Botswana, Pvt Bag 00704, Gaborone, Botswana.

Received 25 April 2016; Accepted 20 July 2016

Let $n \geq 1$ be an integer. Define $\phi_2(n)$ to be the number of positive integers x , $1 \leq x \leq n$, for which both $6x-1$ and $6x+1$ are relatively prime to $6n$. The primary goal of this study is to show that ϕ_2 is a multiplicative function, that is, if $\gcd(m, n) = 1$, then $\phi_2(mn) = \phi_2(m)\phi_2(n)$.

Key words: Euler phi-function, multiplicative function.

THE EULER ϕ_2 FUNCTION

Let $n \geq 1$ be an integer and let $S = \{1, 7, 11, 13, 17, 19, 23, 29\}$, the set of integers which are both less than and relatively prime to 30. In Mothebe and Modise (2016) we define $\phi_3(n)$ to be the number of integers x , $0 \leq x \leq n - 1$, for which $\gcd(30n, 30x + i) = 1$ for all $i \in S$. We proved that this function is multiplicative and thereby obtained a formula for its evaluation.

For each $n \in \mathbb{N}$ let $\phi_2(n)$ denote the number of positive integers x , $1 \leq x \leq n$, for which both $6x - 1$ and $6x + 1$ are relatively prime to $6n$. In this note we draw analogy with our study of ϕ_3 and show that ϕ_2 is multiplicative.

For example if $n = 5 \in S$, then $\phi_2(n) = 3$ since the pairs $(11, 13)$, $(17, 19)$ and $(29, 31)$ are the only ones with components that are relatively prime to 30.

We now proceed to show that we can evaluate $\phi_2(n)$ from the prime factorization of n . Our arguments are based on those used by Burton (2002) to show that the Euler phi-function is multiplicative. We first note:

Theorem 1.1: Let k and s be nonnegative numbers and let $p \geq 5$ be a prime number. Then the following hold:

- (i) $\phi_2(2^k) = 2^k$.
- (ii) $\phi_2(3^s) = 3^s$.
- (iii) $\phi_2(p^k) = p^k - 2p^{k-1}$

Proof. (i) and (ii): For all nonnegative integers k and s and x :

$$\gcd(6x - 1, 6 \cdot 2^k) = \gcd(6x + 1, 6 \cdot 2^k) = 1$$

and

$$\gcd(6x - 1, 6 \cdot 3^s) = \gcd(6x + 1, 6 \cdot 3^s) = 1.$$

Proof. (iii): Clearly $\gcd(6x - 1, 6p^k) = 1$ if and only if p does not divide $6x - 1$ and $\gcd(6x + 1, 6p^k) = 1$ if and only if p does not divide $6x + 1$. There is one integer between 1 and p that satisfies the congruence relation $6x \equiv 1 \pmod{p}$. Hence there are p^{k-1} integers between 1 and p^k that satisfy $6x \equiv 1 \pmod{p}$. Similarly there are p^{k-1} integers of the form $6x + 1$ between 1 and p^k divisible by p . Thus the set $\{(6x - 1, 6x + 1) \mid 1 \leq x \leq p^k\}$

*Corresponding author. E-mail: mothebemf@mopipi.ub.bw.

p^k } contains exactly $p^k - 2p^{k-1}$ pairs corresponding to integers x for which both $\gcd(6x - 1, 6p^k) = 1$ and $\gcd(6x + 1, 6p^k) = 1$. Thus $\phi_2(p^k) = p^k - 2p^{k-1}$.

For example $\phi_2(5^2) = 5^2 - 2 \cdot 5 = 15$ and $\phi_2(5) = 5 - 2 \cdot 1 = 3$ as observed earlier.

We recall that:

Definition 1.2: A number theoretic function f is said to be multiplicative if $f(mn) = f(m)f(n)$ whenever $\gcd(m, n) = 1$.

From the proof of Theorem 1.1 it is clear that for all integers k and s : $\phi_2(2^k 3^s) = \phi_2(2^k)\phi_2(3^s)$.

We now show that the function ϕ_2 is multiplicative. This will enable us to obtain a formula for $\phi_2(n)$ based on a factorization of n as a product of primes. We require the following results:

Lemma 1.3: Given integers m, n , $\gcd(6x - 1, 6mn) = 1$ if and only if $\gcd(6x - 1, 6m) = 1$ and $\gcd(6x - 1, 6n) = 1$.

Similarly given integers m, n , $\gcd(6x + 1, 6mn) = 1$ if and only if $\gcd(6x + 1, 6m) = 1$ and $\gcd(6x + 1, 6n) = 1$. This is an immediate consequence of the following standard result.

Lemma 1.4: Given integers m, n, k , $\gcd(k, mn) = 1$ if and only if $\gcd(k, m) = 1$ and $\gcd(k, n) = 1$.

We note also the following standard result.

Lemma 1.5: If $a = bq + r$, then $\gcd(a, b) = \gcd(b, r)$.

Theorem 1.6: The function ϕ_2 is multiplicative, that is, if $\gcd(m, n) = 1$, then $\phi_2(mn) = \phi_2(m)\phi_2(n)$.

Proof: The result holds if either m or n equals 1. We shall therefore assume neither m nor n equals 1. Arrange the integer pairs $(6x - 1, 6x + 1)$, $1 \leq x \leq mn$, in an $n \times m$ array as follows:

$$\begin{bmatrix} (5,7) & \dots & (6m - 1, 6m + 1) \\ (6(m + 1) - 1, 6(m + 1) + 1) & \dots & (6(2m) - 1, 6(2m) + 1) \\ \vdots & \ddots & \vdots \\ (6((n - 1)m + 1) - 1, 6((n - 1)m + 1) + 1) & \dots & (6(mn) - 1, 6(mn) + 1) \end{bmatrix}$$

we know that $\phi_2(mn)$ is equal to the number of pairs $(6x - 1, 6x + 1)$ in this matrix for which both $6x - 1$ and $6x + 1$ are relatively prime to $6mn$. By virtue of Lemma 1.3 this is the same as the number of pairs $(6x - 1, 6x + 1)$ in the same matrix for which both $6x - 1$ and $6x + 1$ are relatively prime to each of $6m$ and $6n$.

We first note, by virtue of Lemma 1.5, that $\gcd(6(qm + x) - 1, 6m) = \gcd(6x - 1, 6m)$ and likewise

$\gcd(6(qm + x) + 1, 6m) = \gcd(6x + 1, 6m)$. Therefore the pairs $(6(qm + x) - 1, 6(qm + x) + 1)$ in the x^{th} column are both relatively prime to $6m$ if and only if both $6x - 1$ and $6x + 1$ are relatively prime to $6m$.

Therefore only $\phi_2(m)$ columns contain pairs $(6x - 1, 6x + 1)$ both relatively prime to $6m$ and every other pair in the column will constitute of integers both relatively prime to $6m$. The problem now is to show that in each of these $\phi_2(m)$ columns there are exactly $\phi_2(n)$ integer pairs $(6x - 1, 6x + 1)$ that are both relatively prime to $6n$, for then altogether there would be $\phi_2(m)\phi_2(n)$ pairs in the table that are relatively prime to both $6m$ and $6n$.

The entries that are in the x^{th} column (where it is assumed $\gcd(6x - 1, 6m) = \gcd(6x + 1, 6m) = 1$) are: $(6x - 1, 6x + 1)$, $(6(m + x) - 1, 6(m + x) + 1)$, \dots , $(6((n - 1)m + x) - 1, 6((n - 1)m + x) + 1)$.

There are n pairs in this sequence and for no two pairs: $(6(qm + x) - 1, 6(qm + x) + 1)$, $(6(jm + x) - 1, 6(jm + x) + 1)$ in the sequence do we have:

$$6(qm + x) - 1 \equiv 6(jm + x) - 1 \pmod{n}$$

and

$$6(qm + x) + 1 \equiv 6(jm + x) + 1 \pmod{n}$$

since otherwise we would arrive at a contradiction $q \equiv j \pmod{n}$. Thus the terms of the sequence, $x, m + x, 2m + x, \dots, (n - 1)m + x$ are congruent modulo n to $0, 1, 2, \dots, n - 1$ in some order.

Now suppose t is congruent modulo n to $qm + x$. Then the integers, $6(qm + x) - 1$ and $6(qm + x) + 1$ are both relatively prime to $6n$ if and only if both $6t - 1$ and $6t + 1$ are relatively prime to $6n$. The implication is that the x^{th} column contains as many pairs of integers that are relatively prime to $6n$ as does the set $\{(5, 7), (11, 13), \dots, (6n - 1, 6n + 1)\}$, namely $\phi_2(n)$ pairs. Thus the number of pairs of integers $(6x - 1, 6x + 1)$ in the matrix that are relatively prime to $6m$ and $6n$ is $\phi_2(m)\phi_2(n)$. This completes the proof of the theorem.

The following result immediately follows from Theorem 1.1 and Theorem 1.6:

Corollary 1.7: If the integer $n > 1$ has the prime factorization $n = 2^{k_1} 3^{k_2} p_3^{k_3} \dots p_r^{k_r}$ then $\phi_2(n) = 2^{k_1} 3^{k_2} (p_3^{k_3} - 2p_3^{k_3-1}) \dots (p_r^{k_r} - 2p_r^{k_r-1})$.

Conflict of Interests

The authors have not declared any conflict of interests.

REFERENCES

Burton DM (2002). Elementary Number Theory. McGraw-Hill, International Editions.

Mothebe MF, Modise BT (2016). An extension of the Euler Phi-function to Sets of Integers Relatively Prime to 30. J. Math. Res. 8(2):45-48. <http://dx.doi.org/10.5539/jmr.v8n2p45>.



African Journal of Mathematics and Computer Science Research

Related Journals Published by Academic Journals

- *African Journal of Pure and Applied Chemistry*
- *International Journal of Physical Sciences*
- *Journal of Geology and Mining Research*
- *Journal of Environmental Chemistry and Ecotoxicology*
- *Journal of Internet and Information Systems*
- *Journal of Oceanography and Marine Science*
- *Journal of Petroleum Technology and Alternative Fuels*

academicJournals