

A photograph of a person harvesting carrots in a field. The person is wearing a green shirt and a green rubber boot. They are holding a bunch of harvested carrots in their left hand and pulling a carrot from the soil with their right hand. The background shows a green field with other plants.

Journal of Stored Products and Postharvest Research

Volume 8 Number 4 April 2017

ISSN 2141-6567



*Academic
Journals*

ABOUT JSPPR

The Journal of Stored Products and Postharvest Research (JSPPR) is published monthly (one volume per year) by Academic Journals.

The Journal of Stored Products and Postharvest Research (JSPPR) is an open access journal that provides rapid publication (monthly) of articles in all areas of the subject such as Biological control of rot-inducing fungi, Post harvest losses of rice from harvesting to milling, Genetic variability studies between released varieties of cassava, Seed borne pathogens on farmer-saved sorghum etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in JSPPR are peer-reviewed.

Contact Us

Editorial Office: jsppr@academicjournals.org

Help Desk: helpdesk@academicjournals.org

Website: <http://www.academicjournals.org/journal/JSPPR>

Submit manuscript online <http://ms.academicjournals.me/>

Editors

Dr. Dalbir Singh Sogi

Department of Food Science and Technology
Guru Nanak Dev University Amritsar 143 005
India

Dr. Raquel Lima Miranda

397 O St SW Washington DC 20024
USA.

Dr. Héctor Eduardo Martínez Flores

Division of Graduate Studies of the Chemical Pharmacology
Faculty Universidad Michoacana de San Nicolás de Hidalgo
Mexico.

Dr. Naveen Kumar

University of Florida
Southwest Florida Research and Education Centre
USA.

Editorial Board

Prof. N.M. Nnam

Department of Home Science Nutrition and Dietetics
University of Nigeria Nsukka
Nigeria

Dr. Sunil Pareek

Department of Horticulture
Rajasthan College of Agriculture
Maharana Pratap University of Agriculture and Technology
Udaipur, Rajasthan, 313001
India

Dr. K.A. Raveesha

Department of Studies in Botany
University of Mysore
Manasagangotri
Mysore-570 006
India

Dr. M.M. Prasad

Central Institute of Fisheries Technology
Visakhapatnam Research Centre
India

Dr. Charles Fokunang

Department of Pharmacy
Faculty of Medicine and Biomedical Sciences
University of Yaounde
Cameroon.

Dr.Zachée Ngoko

Institute for Agricultural Research and Development (IRAD)
Bambui, Box 80 Bamenda
Cameroon

Dr.Zachée Ngoko

Institute for Agricultural Research and Development (IRAD)
Bambui, Box 80 Bamenda
Cameroon

Dr. Mahbub Hasan

Department of Zoology
Rajshahi University
Rajshahi-6205
Bangladesh

Dr. Mohamed A. Eltawil

Agric. Engineering Department
Kafrelsheikh University
Egypt

Dr. Younes Rezaee Danesh

Department of Plant Protection
Faculty of Agriculture
Urmia University
Iran

ARTICLE

- Determination of the appropriate doses of promising botanical powders against maize weevil, *Sitophilus zeamais* Mots (Coleoptera: Curculionidae) on maize grain** 49
Tadele Shiberu and Mulugeta Negeri

Full Length Research Paper

Determination of the appropriate doses of promising botanical powders against maize weevil, *Sitophilus zeamais* Mots (Coleoptera: Curculionidae) on maize grain

Tadele Shiberu* and Mulugeta Negeri

Department of Plant Sciences, College of Agriculture and Veterinary Sciences, Ambo University, Ethiopia.

Received 19 September, 2014; Accepted 29 October, 2014

The present laboratory study was carried out in the entomological laboratory the Department of plant science, Ambo University. The experiment was carried out using randomized complete block design with three replications to evaluate the efficacy of four locally available botanicals. The botanicals evaluated were: Neem (*Azadirachta indica*) leaf and seed, hop bush (*Dodonaea angustifolia*) fresh leaf, Pyrethrum (*Chrysanthemum cinerariaefolium*) flower and Lomen grass (*Cymbopogon citratus*) leaf powders at three different doses (3, 4 and 5 g)/100 g of grain maize against maize weevil, *Sitophilus zeamais* Motsch. Their effectiveness was determined using different parameter such as grain damage and weight losses. The results revealed that all test materials at 5 g doses exhibited mortality action against maize weevil while at 3 g doses it gave the lowest percentage efficacy. Powdered leaves and seeds of neem and pyrethrum flower at 4 and 5 g were showed statistical significant ($P < 0.01$) differences while powdered leaves of lemon grass at the similar rates were observed with moderately effectiveness and fresh leaves of hop bush gave the lowest mortality rate within 28 days of exposure in all tested doses. The result showed that the lower number of grains damaged in maize grains treated with botanical powders and the grain weight loss was also found low as compared to the untreated control. These findings suggest that botanical powders exert better mortality effect and hence reduced maize grain damage. It was also revealed that despite the high seed damage recorded in all botanical products, the *S. zeamays* numbers were relatively higher in untreated control while no grain seed damage was observed in standard check which was treated with Actellic dust. Maize grains treated with botanicals, indicated that insect reproduction and development were impaired in all botanical pesticides. However, all the doses tested for their insecticidal efficacy had an effect on the percentage weevil mortality and was found to be directly proportional to the amount of powder used. Therefore, The present study was suggested that Pyrethrum flower, neem leaf and seed powder can be used as good alternatives to chemical insecticides against *S. zeamays* due to their higher high mortality, lower grain damage and lower maize weight losses recorded as compared to untreated and synthetic insecticides.

Key words: Botanicals, powder, mortality, maize weevil, *Sitophilus zeamais*, grain damage, actellic dust.

INTRODUCTION

Maize is the third most important cereal crop following wheat and rice. It is also, one of the most popular crop

plants all over the world, grown in over 140 million hectares (Zaidi and Singh, 2005). In Ethiopia, Maize is one of the major staples consumed food second place in contribution which 17% of land and production, 205,472.37 ton in year of 2011/2012 (CSA, 2012).

Dankyi et al. (1995) reported maize is an important source of carbohydrate and forms about 90 to 95% of the total calories intake of the coastal Savanna people. Insect pest damage to stored grains results in major economic losses in Ethiopia where subsistence grain production supports the livelihood of majority of the population; grain loss caused by storage pests such as the maize weevil, *Sitophilus zeamais* (Mots.) threatens food security. Reduction of insect damage in stored grains is a serious problem in developing countries in the tropics due to favorable climatic conditions and poor storage structures (Bekele et al., 1997).

Damage of maize grain begins from the field just before harvest and the insects are carried into the store where the population builds up rapidly (Adedire and Lajide, 2003). However, during storage, maize seeds are very susceptible to attack by maize weevils among of these *S. zeamais* is very serious. Grain damaged by insect pests in storage often result in significant losses including loss of viability (ISTA, 1995). Post harvest storage of maize is highly constrained by insect pest of maize weevil. Yield losses due to *S. zeamais* reported from farmers interviews, storage loss of maize due to weevil was estimated to be about 25 to 30% in the western part of Ethiopia (Emana, 1998). There are however, major problem to the use of chemical insecticides including human risk, cost of procurement, availability of insecticides on the market, development of resistant strains and residue in the food crops which has led scientists to investigating plant products as alternatives (Lale, 2002). Moreover, there is a need to determine an appropriate dose of some botanicals to effectively control storage insect pests. Therefore, this study was carried out to determine the potential efficacy of different plant products at different doses in powder form to use in the management of *S. zeamais*, under storage conditions.

MATERIALS AND METHODS

Rearing of maize weevils

The study was carried out at Ambo University plant science Department under the laboratory of Entomology, having an altitude of 2287 m above sea level, latitude of 08°7'0" North and longitude of 38°7'0" East. The mean maximum and minimum temperature of the laboratory was 24.2±2°C and 16.8±2°C, respectively, during the

study period.

The insects were reared on whole maize in 5 L plastic jars after being treated for mites (Udo, 2000). Whole grains of local variety bought from local market were disinfested in an oven at 60°C for 1 h (Asmanizar and Idris, 2008) before using them as a substrate for insect rearing. The moisture content of maize grain was adjusted to 12 to 13%. After three weeks of oviposition, the parent weevils were sieved out after oviposition. Later the grain were kept in the incubator for adult emergence while the emerging generation of same age insects re-cultured at 24±2°C, 65±5% relative humidity (R.H.). The F1 generation was used for the experiment.

Collection and preparation of materials

Leaves and seeds of neem, *Azadirachta indica* were collected from Melka Werer Agricultural Research center, flower of Pyrethrum, *Chrysanthemum cinerariaefolium* was collected from Kulumsa Agricultural research center, leaves of hop bush, *Dodonaea angustifolia* and lomen grass, *Cymbopogon citrates* were collected from Ambo areas. All the plant materials were dried under shade, in a well-ventilated area in the Entomological Laboratory of Plant Science Department, before grinding into fine powders using hand machine blender and sieved using 80 µm Laboratory sieve. The Actellic dust 2% insecticide was purchased from private limited chemical company. The powders were tested individually for their potential in protecting maize grains during storage.

Treatment application

The experiment was laid out in Randomized Complete Block Design (RCBD) with three replications. Twenty pairs of *S. zeamais* were introduced into the plastic jar containing 100 g grain maize in 0.5 L plastic jars at 3, 4, and 5 g% (w/w), while in control treatment there was no any botanicals added and in the standard check Actellic 2% dust was added in 0.1 g/100 g of maize grain. Each botanical was weighed and added to the maize grain in each jar and shaken well for uniform coating. The jars were covered with muslin cloth and secured with rubber bands as a ventilated lid. The treated grains in the jar were kept for about 28 days and mortality rate assessments were performed regularly every 1, 7, 14, 21 and 28 days after exposure of botanical powders.

Seed germination

The effect of treatments and their interactions on seed germination and viability was examined after 28 days of grain storage period. Seed germination was tested using 50 randomly picked seeds from undamaged grains after separation of damaged and undamaged grains in each jar according to the methods described in Haines (1991). The 50 grain sub-samples were germinated on moisture filter paper (Whatman No. 1) in Petri dishes arranged in a RCBD with four replicates. The experiment was maintained under laboratory conditions. The number of germinated seedlings from each Petri dish was counted and recorded after 7 days. The percent germination was computed according to the methods of Zibokere (1994) as follows:

*Corresponding author. E-mail: tshiberu@yahoo.com.

Author(s) agree that this article remain permanently open access under the terms of [the Creative Commons Attribution License 4.0 International License](https://creativecommons.org/licenses/by/4.0/)

$$\text{Viability index (\%)} = \frac{NG \times 100}{TG}$$

Where NG = number of seeds that germinated, TG = total number of test seeds.

Data collection

Mortality rate were assessed for damage arising from natural insect infestations after 1, 7, 14, 21 and 28 days of treatment application. The assessment periods selected based on an earlier report by Dobie (1974) and Girma et al. (2008). Dead weevils were removed and counted during each assessment. Damage assessment was carried out on treated and untreated grains by taking samples of 50 grains from each jar. Each treatment was separated into undamaged and insect-damaged grains. The number of grains in each treatment was counted, weighed and the percent weight loss (percent insect damage) of maize grains in storage was computed according to the methods described in Haines (1991) as follows:

$$\text{Percent of weight loss} = \frac{UNd - DNu}{U(Nd + Nu)} \times 100$$

Where U = weight of undamaged grains, D = weight of insect-damaged grains, Nu = number of undamaged grains, Nd = number of insect-damaged grains.

Data analysis

The Analysis of Variance (ANOVA) was performed using Statistical Analysis Software (SAS, 2003), Data on % cumulative mortality as least significant difference (LSD) test was used to determine the differences among the means for different treatments, grain damage and yield loss parameters.

RESULTS AND DISCUSSION

The results revealed that significant ($P < 0.05$) differences among the treatments at all the doses tested for their insecticidal efficacy and showed an effect on the percentage weevil mortality and was found to be directly proportional to the amount of powder used. However, lower mortality was observed within one day after the exposure of weevils to botanical powders at all treatments. A very lower amount of mortality resulted within 28 days when the weevils were exposed to the lowest dose (3 g) of the botanical powders. Both leaf and seed powder of *A. indica* gave the efficacy of 86.33 and 83.33% at the dose of (5 g) against maize weevils after 7th days of exposure, respectively. Similarly, after 14th days of application the efficacy of the neem seed was showed similar results, while neem leaves gave 100% mortality. The other botanicals indicated moderate results at the highest dose after 14th day exposure except *C. cinerariaefolium* was showed 92.33% mortality rate and it gave 100% mortality rate within 21 days of exposure. Whereas *Cymbopogon citrates* gave a moderate percent

at 5 g dose 62.22% mortality when compared with other botanical treatments and it was showed significantly ($P < 0.01$) different results from that of untreated check. *Dodonaea angustifolia* leaf powder gave less mortality rate even at 5 g dose application within 28 days of exposure.

Results confirm that botanicals can be used to protect stored maize grains against *S. zeamays* infestations. The use of botanical products and other available materials to protect stored maize grains have been reported by other workers (Lale, 1995; Keita et al., 2000). Lale (2000) had advanced that oil products from the neem plant are particularly more effective against insects. *Sitophilus zeamays* lays eggs inside the maize grains therefore the present study revealed that the use of different botanical powders could be applied in the control of *S. zeamays*. In this study the live maize weevils *S. zeamays* showed least in maize grains that were treated with pyrethrum flower and neem leaf powder as compared with other plant materials. Marilei et al. (2010), also reported that 40 g of corn treated with 6g of the extracts from leaves and seeds extracts from neem can be considered as a viable alternative for controlling the *S. zeamais* in stored corn. The results of the current study was confirmed by the previous work of Wanyika et al. (2009), that they have reported *C. cinerariaefolium*, affected the survival rate of the adult weevils in treatments (5.0 and 7.5 g) with a mortality rate range from 76.66 to 100%. It is also supported by the findings of Maribet and Aurea (2008), who found that lower mortality of adult maize weevils from corn grains treated with *C. citrates* within 24 days after insect introduction.

Endersby and Morgan (1991) observed that *A. indica* derivations are most effective as feeding poisons for nymphs or larvae of phytophagous insects. Hence, mortality rate of maize storage weevils, *S. zeamays* that caused by some botanicals as reported in the present study is most likely due to these inherent properties. Moreover, the results indicated that all doses gave more than 30% deaths of weevils were dead after 28 days of exposure to the botanicals powder (Table 1). All tested botanicals performed well in the reduction of maize weevil during maize storage as compared to the untreated control.

Grain damage and weight losses

The grain damaged and weight losses concerned from the maize grains treated with neem seed and pyrethrum flower powders, the lowest values were recorded at the dose of 5 g. Similar numbers of weigh losses of maize grains were relatively recorded in all the treatments. This study suggests that neem leaf, neem seed and pyrethrum flower powder can be used as good alternatives to manage maize weevil *S. zeamays* from maize storage (Table 2).

Table 1. Effects of different botanicals at different doses on the survival of maize weevils, *S. zeamais*.

Treatments	Rate/100 g	Mortality percent after treatment application				
		1 st day	7 days	14 days	21 days	28 days
<i>Azadirachta indica</i> (seed)	3	10.00 ^c	25.67 ^d	40.00 ^d	40.0 ^c	46.67 ^d
	4	11.11 ^c	36.67 ^c	53.33 ^c	92.33 ^a	93.33 ^a
	5	13.33 ^c	83.33 ^b	83.33 ^a	95.55 ^a	95.55 ^a
<i>Azadirachta indica</i> (Leaf)	3	0.00 ^d	23.33 ^d	55.55 ^c	63.33 ^b	63.33 ^c
	4	10.00 ^c	36.67 ^c	62.22 ^c	100.0 ^a	100.0 ^a
	5	11.11 ^c	86.33 ^b	100.0 ^a	100.0 ^a	100.0 ^a
<i>Dodonae angustifolia</i> (fresh leaf)	3	0.00 ^d	10.00 ^g	13.33 ^{fg}	16.67 ^{ef}	16.67 ^f
	4	0.00 ^d	13.33 ^{ef}	20.0 ^{efg}	20.00 ^e	20.00 ^f
	5	0.00 ^d	20.00 ^{de}	30.0 ^{de}	30.77 ^d	31.78 ^e
<i>Chrysanthemum cinerariaefolium</i> (flower)	3	8.89 ^c	10.00 ^g	60.00 ^c	60.00 ^b	86.67 ^b
	4	32.33 ^b	13.33 ^{ef}	76.67 ^b	100.0 ^a	100.0 ^a
	5	25.56 ^b	20.00 ^{de}	92.33 ^a	100.0 ^a	100.0 ^a
<i>Cymbopogon citrates</i>	3	0.00 ^d	10.00 ^g	10.0 ^{gh}	10.00 ^g	34.44 ^e
	4	0.00 ^d	10.00 ^g	10.0 ^{gh}	10.00 ^g	47.78 ^d
	5	0.00 ^d	10.00 ^g	21.11 ^{ef}	40.00 ^c	62.22 ^c
Actellic dust 2%		100 ^a	100.0 ^a	100.0 ^a	100.0 ^a	100.0 ^a
Untreated check		0.00 ^d	1.11 ^g	1.11 ^h	3.33 ^g	5.56 ^g
MSE		2.45	4.11	4.96	3.74	4.29
LSD		5.50	9.17	11.08	8.37	9.60
CV (%)		17.00	13.75	10.03	6.50	6.61

Means with the same letter are not significantly different.

Table 2. Effects of botanicals and *S. zeamais* on grain weight loss and germination.

Treatments	Rate/100 g	Grain damage (%)	Weight losses (%)
<i>Azadirachta indica</i> (seed)	3	3.67 ^{cbd}	0.27 ^{bcde}
	4	2.00 ^{ed}	0.27 ^{bcde}
	5	1.67 ^e	0.17 ^{bcde}
<i>Azadirachta indica</i> (Leaf)	3	3.33 ^{cde}	0.70 ^{bcde}
	4	3.00 ^{cde}	0.70 ^{bcde}
	5	3.33 ^{cde}	0.20 ^{bcde}
<i>Dodonae angustifolia</i> (leaf)	3	3.00 ^{cde}	0.67 ^{bcde}
	4	2.00 ^e	0.63 ^{bcde}
	5	2.67 ^{cde}	0.47 ^{bcde}
<i>Chrysanthemum cinerariaefolium</i> (flower)	3	4.33 ^{bc}	0.10 ^{bcde}
	4	5.33 ^b	0.03 ^{de}
	5	2.33 ^{de}	0.07 ^{cde}
<i>Cymbopogon citrates</i>	3	4.33 ^{bc}	0.80 ^b
	4	2.50 ^{de}	0.60 ^{bcde}
	5	2.17 ^{de}	0.73 ^{bc}
Actellic dust 2%		0.00 ^f	0.00 ^e
Untreated check		12.67 ^a	5.33 ^a
MSE		0.76	0.31
LSD		1.67	0.7
CV (%)		22.09	20.19

Means with the same letter are not significantly different.

Conclusion

In general, this study has revealed the plant powders at 5 g dosage had significant effects on mortality rate against *S. zeamais* within 28 days of exposure on maize grains. In particular, *A. indica* (seed), *A. indica* (leaf) and *Chrysanthemum cinerariaefolium* (flower) were observed at the dosage of 5 g with good performance after 7 to 21 days of applications and gave of 83.33 to 100% of mortality and maize grain damage and weight losses. Finally, it is recommended that neem seed and leaf, and pyrethrum flowers were effective when used as at 4 g powder/100 g of maize grain against maize weevil. It is also economically important and environmentally safe.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

REFERENCES

- Adedire CO, Lajide L (2003). Ability of extracts of tropical plant species to protect maize against infestation by the maize weevil, *Sitophilus zeamais* during storage. *Nigeria J. Exp. Biol.* 2:175-179.
- Asmanizar DA, Idris AB (2008). Effect of selected plant extract on mortality of adult *Sitophilus zeamais* Motschulsky (Coleoptera: Curculionidae), pest of stored rice grains. *Malaysian Appl. Biol.* 37(2):41-46.
- Bekele AJ, Obeng-Ofori D, Hassanali A (1997). Evaluation of *Ocimum kenyense* (Ayobangira) as source of repellents, toxicants and protectants in storage against three major stored product insect pests. *J. Appl. Ent.* 121:169-173.
- CSA (2012). Report of Federal Democratic Republic of Ethiopia, Statistical Report on Socio-Economic Characteristics of the Population in Agricultural Households, Land Use, Area and Production of Crops. Addis Ababa, Ethiopia.
- Dankyi AA, Anchirinah VM, Apau AO (1995). Profitability and impact of extension test plots on maize production. *Newsletter* pp. 1-3.
- Dobie P (1974). The susceptibility of different type of maize to post harvest infestation by *Sitophilus zeamais* and *Sitotroga cerealella* and the importance of this factor at the small scale farm level CIMMYT No. 10:98-113.
- Emana G (1998). Use of botanical plants in the control of stored maize grain insect pests in Ethiopia. In: *Maize production technology for the future: Challenges and opportunities*. Proceedings of the sixth Eastern and Southern Africa Regional Maize Conference. pp. 105-108.
- Endersby NM and Morgan WC (1991). Alternatives to synthetic chemical insecticides for use in crucifer crops. *Biol. Agric. Hort.* 8:33-52.
- Girma D, Tadele T, Abreham T (2008). Management of the maize weevil *Sitophilus zeamais* Motsch, (Coleoptera: Curculionidae) using botanical insecticides on three maize genotypes. *Pest Manag. J. Ethiopia* 12:49-56.
- Haines CP (1991). *Insects and Arachnids of tropical stored products: Their biology and identification*. A Training Manual. 2nd edition. Natural Resources Institute.
- ISTA (1995). *Handbook on vigour test methods for maize*. 117 pages.
- Keita MS, Vincent C, Schmith JP, Ramaswamy S, Belanger A (2000). Effect of various essential oil on *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae). *J. Stored Prod. Res.* 36:335-364.
- Lale NES (1995). An overview of the use of plant products in the management of stored product coleoptera in the products. *Post Harvest News Inform*, 6:69N-75N.
- Lale NES (2000). *Stored Product Entomology and Acarology in tropical Africa*. Molepublication, Maiduguri, Nigeria. 204 p.
- Maribet LP, Aurea CR (2008). Insecticidal action of five plants against maize weevil, *Sitophilus zeamais* motsch. (Coleoptera: Curculionidae). *KMITL Sci. Tech. J.* 8:1.
- Marilei NS, Alberto CV, Augusto TAP, Heron OSL (2010). Leaves, stems and seeds powder of neem (*Azadirachta indica* A. Juss.) used as repellent of *Sitophilus zeamais* (Coleoptera: Curculionidae) in stored corn. *Open journal systems*.
- SAS Institute (2003). *The SAS System version 9.1 for windows*. SAS Institute, Cary, NC, USA.
- Udo IO (2000). Efficacy of Candlewood, *Zanthoxylum xanthoxyloides* Lam. For the control of three stored product pests. M. Phil Thesis, University of Ghana, Legon – Accra. P 81.
- Wanyika HN, Kareru PG, Keriko JM, Gachanja AN, Kenji NJ, Mukiira GM (2009). Contact toxicity of some fixed plant oils and stabilized natural pyrethrum extracts against adult maize weevils (*Sitophilus zeamais* Motsch.). *Afr. J. Pharm. Pharmacol.* 3(2):066-069.
- Zaidi G, Singh SKF (2005). Morphology and growth of maize – IITA/CIMMYT Research, Guide 12, IITA, Ibadan, Nigeria. pp. 15-18.
- Zibokere DS (1994). Insecticidal potency of red pepper (*Capsicum annuum*) on pulse beetle (*Callosobruchus maculatus*) infesting cowpea (*Vigna unguiculata*) seeds during storage. *Indian J. Agric. Sci.* 64:727-728.

A photograph of a person's hands holding a bunch of fresh, orange carrots with green tops. The background is a blurred green field, suggesting a farm or garden setting. The image is framed with rounded corners.

Journal of Stored Products and Postharvest Research

Related Journals Published by Academic Journals

- Journal of Plant Breeding and Crop Science
- African Journal of Agricultural Research
- Journal of Horticulture and Forestry
- International Journal of Livestock Production
- Journal of Soil Science and Environmental Management
- Journal of Cereals and Oilseeds

academicJournals