



International Journal of Genetics and Molecular Biology

Volume 9 Number 4, November 2017

ISSN 2006-9863



*Academic
Journals*

ABOUT IJGMB

The **International Journal of Genetics and Molecular Biology (IJGMB)** (ISSN 2006-9863) is published Monthly (one volume per year) by Academic Journals.

International Journal of Genetics and Molecular Biology (IJGMB) provides rapid publication (monthly) of articles in all areas of the subject such as DNA and RNA, Influence of risk factors on onset of hyperlipidemia in people with cerebrovascular insult, Study of proteinase activity of *Lactobacillus plantarum* etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in IJGMB are peer-reviewed.

Contact Us

Editorial Office: ijgmb@academicjournals.org

Help Desk: helpdesk@academicjournals.org

Website: <http://www.academicjournals.org/journal/IJGMB>

Submit manuscript online <http://ms.academicjournals.me/>.

Editor

Prof. Kasem Zaki Ahmed

*Department of Genetics,
Faculty of Agriculture,
Minia University, El-Minia,
Egypt, ET – 61517*

Prof. Evgeny N.ti Imyanitov N.N.

*Petrov Institute of Oncology,
Pesochny-2, 197758, St.-Petersburg,
Russia*

Dr. A. Muthusamy

*Department of Biotechnology
Manipal Life Sciences Centre
Manipal University
Planetarium Complex
Manipal – 576 104
Karnataka,
India*

Associate Editors

Dr. Chang-Gu Hyun

*Jeju Biodiversity Research Institute (JBRI) &
JeJu Hi-Tech Industry Development Institute (HiDI),
South Korea*

Santosh A. Khedkar

*Computational Medicinal Chemist
45 Aldrich St. Apartment-1
Somerville MA 02145,
USA*

Dr. Yehia Zakaria Gad

*Department of Medical Molecular Genetics, Division
of Human Genetics and Genome Research, National
Research Center, El-Behooth (ex-Tahrir) st., Dokki,
Giza, 12311
Egypt*

Editorial Board

Dr. Ibrahim Ilker Ozyigit

*Marmara University, Sciences and Arts Faculty,
Department of Biology, 34722, Goztepe,
Istanbul,
Turkey*

Dr. (Mrs.) Meena Misra

*Institute of Frontier Sciences & Biotechnology,
Baramunda,
Bhubaneswar-751004,
India*

Instructions for Author

Electronic submission of manuscripts is strongly encouraged, provided that the text, tables, and figures are included in a single Microsoft Word file (preferably in Arial font).

The **cover letter** should include the corresponding author's full address and telephone/fax numbers and should be in an e-mail message sent to the Editor, with the file, whose name should begin with the first author's surname, as an attachment.

Article Types

Three types of manuscripts may be submitted:

Regular articles: These should describe new and carefully confirmed findings, and experimental procedures should be given in sufficient detail for others to verify the work. The length of a full paper should be the minimum required to describe and interpret the work clearly.

Short Communications: A Short Communication is suitable for recording the results of complete small investigations or giving details of new models or hypotheses, innovative methods, techniques or apparatus. The style of main sections need not conform to that of full-length papers. Short communications are 2 to 4 printed pages (about 6 to 12 manuscript pages) in length.

Reviews: Submissions of reviews and perspectives covering topics of current interest are welcome and encouraged. Reviews should be concise and no longer than 4-6 printed pages (about 12 to 18 manuscript pages). Reviews are also peer-reviewed.

Review Process

All manuscripts are reviewed by an editor and members of the Editorial Board or qualified outside reviewers. Authors cannot nominate reviewers. Only reviewers randomly selected from our database with specialization in the subject area will be contacted to evaluate the manuscripts. The process will be blind review.

Decisions will be made as rapidly as possible, and the journal strives to return reviewers' comments to authors as fast as possible. The editorial board will re-review manuscripts that are accepted pending revision. It is the goal of the AJFS to publish manuscripts within weeks after submission.

Regular articles

All portions of the manuscript must be typed double-spaced and all pages numbered starting from the title page.

The Title should be a brief phrase describing the contents of the paper. The Title Page should include the authors' full names and affiliations, the name of the corresponding author along with phone, fax and E-mail information. Present addresses of authors should appear as a footnote.

The Abstract should be informative and completely self-explanatory, briefly present the topic, state the scope of the experiments, indicate significant data, and point out major findings and conclusions. The Abstract should be 100 to 200 words in length. Complete sentences, active verbs, and the third person should be used, and the abstract should be written in the past tense. Standard nomenclature should be used and abbreviations should be avoided. No literature should be cited.

Following the abstract, about 3 to 10 key words that will provide indexing references should be listed.

A list of non-standard **Abbreviations** should be added. In general, non-standard abbreviations should be used only when the full term is very long and used often. Each abbreviation should be spelled out and introduced in parentheses the first time it is used in the text. Only recommended SI units should be used. Authors should use the solidus presentation (mg/ml). Standard abbreviations (such as ATP and DNA) need not be defined.

The Introduction should provide a clear statement of the problem, the relevant literature on the subject, and the proposed approach or solution. It should be understandable to colleagues from a broad range of scientific disciplines.

Materials and methods should be complete enough to allow experiments to be reproduced. However, only truly new procedures should be described in detail; previously published procedures should be cited, and important modifications of published procedures should be mentioned briefly. Capitalize trade names and include the manufacturer's name and address. Subheadings should be used. Methods in general use need not be described in detail.

Results should be presented with clarity and precision. The results should be written in the past tense when describing findings in the authors' experiments. Previously published findings should be written in the present tense. Results should be explained, but largely without referring to the literature. Discussion, speculation and detailed interpretation of data should not be included in the Results but should be put into the Discussion section.

The Discussion should interpret the findings in view of the results obtained in this and in past studies on this topic. State the conclusions in a few sentences at the end of the paper. The Results and Discussion sections can include subheadings, and when appropriate, both sections can be combined.

The Acknowledgments of people, grants, funds, etc should be brief.

Tables should be kept to a minimum and be designed to be as simple as possible. Tables are to be typed double-spaced throughout, including headings and footnotes. Each table should be on a separate page, numbered consecutively in Arabic numerals and supplied with a heading and a legend. Tables should be self-explanatory without reference to the text. The details of the methods used in the experiments should preferably be described in the legend instead of in the text. The same data should not be presented in both table and graph form or repeated in the text.

Figure legends should be typed in numerical order on a separate sheet. Graphics should be prepared using applications capable of generating high resolution GIF, TIFF, JPEG or Powerpoint before pasting in the Microsoft Word manuscript file. Tables should be prepared in Microsoft Word. Use Arabic numerals to designate figures and upper case letters for their parts (Figure 1). Begin each legend with a title and include sufficient description so that the figure is understandable without reading the text of the manuscript. Information given in legends should not be repeated in the text.

References: In the text, a reference identified by means of an author's name should be followed by the date of the reference in parentheses. When there are more than two authors, only the first author's name should be mentioned, followed by 'et al'. In the event that an author cited has had two or more works published during the same year, the reference, both in the text and in the reference list, should be identified by a lower case letter like 'a' and 'b' after the date to distinguish the works.

Examples:

Abayomi (2000), Agindotan et al. (2003), (Kelebeni, 1983), (Usman and Smith, 1992), (Chege, 1998;

1987a,b; Tijani, 1993,1995), (Kumasi et al., 2001) References should be listed at the end of the paper in alphabetical order. Articles in preparation or articles submitted for publication, unpublished observations, personal communications, etc. should not be included in the reference list but should only be mentioned in the article text (e.g., A. Kingori, University of Nairobi, Kenya, personal communication). Journal names are abbreviated according to Chemical Abstracts. Authors are fully responsible for the accuracy of the references.

Examples:

Chikere CB, Omoni VT and Chikere BO (2008). Distribution of potential nosocomial pathogens in a hospital environment. *Afr. J. Biotechnol.* 7: 3535-3539.

Moran GJ, Amii RN, Abrahamian FM, Talan DA (2005). Methicillinresistant *Staphylococcus aureus* in community-acquired skin infections. *Emerg. Infect. Dis.* 11: 928-930.

Pitout JDD, Church DL, Gregson DB, Chow BL, McCracken M, Mulvey M, Laupland KB (2007). Molecular epidemiology of CTXM-producing *Escherichia coli* in the Calgary Health Region: emergence of CTX-M-15-producing isolates. *Antimicrob. Agents Chemother.* 51: 1281-1286.

Pelczar JR, Harley JP, Klein DA (1993). *Microbiology: Concepts and Applications*. McGraw-Hill Inc., New York, pp. 591-603.

Short Communications

Short Communications are limited to a maximum of two figures and one table. They should present a complete study that is more limited in scope than is found in full-length papers. The items of manuscript preparation listed above apply to Short Communications with the following differences: (1) Abstracts are limited to 100 words; (2) instead of a separate Materials and Methods section, experimental procedures may be incorporated into Figure Legends and Table footnotes; (3) Results and Discussion should be combined into a single section.

Proofs and Reprints: Electronic proofs will be sent (e-mail attachment) to the corresponding author as a PDF file. Page proofs are considered to be the final version of the manuscript. With the exception of typographical or minor clerical errors, no changes will be made in the manuscript at the proof stage.

Fees and Charges: Authors are required to pay a \$550 handling fee. Publication of an article in the International Journal of Genetics and Molecular Biology is not contingent upon the author's ability to pay the charges. Neither is acceptance to pay the handling fee a guarantee that the paper will be accepted for publication. Authors may still request (in advance) that the editorial office waive some of the handling fee under special circumstances

Copyright: © 2017, Academic Journals.

All rights Reserved. In accessing this journal, you agree that you will access the contents for your own personal use but not for any commercial use. Any use and or copies of this Journal in whole or in part must include the customary bibliographic citation, including author attribution, date and article title.

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract or as part of a published lecture, or thesis) that it is not under consideration for publication elsewhere; that if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher.

Disclaimer of Warranties

In no event shall Academic Journals be liable for any special, incidental, indirect, or consequential damages of any kind arising out of or in connection with the use of the articles or other material derived from the IJGMB, whether or not advised of the possibility of damage, and on any theory of liability.

This publication is provided "as is" without warranty of any kind, either expressed or implied, including, but not limited to, the implied warranties of merchantability, fitness for a particular purpose, or non-infringement. Descriptions of, or references to, products or publications does not imply endorsement of that product or publication. While every effort is made by Academic Journals to see that no inaccurate or misleading data, opinion or statements appear in this publication, they wish to make it clear that the data and opinions appearing in the articles and advertisements herein are the responsibility of the contributor or advertiser concerned. Academic Journals makes no warranty of any kind, either express or implied, regarding the quality, accuracy, availability, or validity of the data or information in this publication or of any other publication to which it may be linked.

ARTICLE

- Broad sense heritability and genetic advance for grain yield and yield components of chickpea (*Cicer arietinum* L.) genotypes in western Ethiopia** 16
Biru Alemu, Kassahun Tesfaye, Teklehaimanot Haileselassie
and Dagnachew Lule

Full Length Research Paper

Broad sense heritability and genetic advance for grain yield and yield components of chickpea (*Cicer arietinum* L.) genotypes in western EthiopiaBiru Alemu^{1,3*}, Kassahun Tesfaye^{2,3,4}, Teklehaimanot Haileselassie⁴ and Dagnachew Lule⁵¹Haro Sabu Agricultural Research Center, P. O. Box 10, Dale Sadi, Kellem Wollega, Ethiopia.²Ethiopian Institute of Biotechnology, Addis Ababa, Ethiopia.³Department of Microbial, Cellular, and Molecular Biology, Addis Ababa University, Addis Ababa, Ethiopia.⁴Institute of Biotechnology, Addis Ababa University, Addis Ababa, Ethiopia.⁵Oromia Agricultural Research Institute, Addis Ababa, Ethiopia.

Received 6 October, 2017; Accepted 21 November, 2017

Genetic variability is a prerequisite to crop improvement. The objective of the present study was to assess and quantify the genetic variability, estimate heritability and genetic advance for yield and yield contributing characters of released and pipeline chickpea varieties based on agro-morphological traits. A total of 16 chickpea materials (8 improved varieties, 7 advanced lines, and one local check) were planted in Randomized Complete Block Design (RCBD) with three replications at Shambu, Hawa Galan, Mata, Alaku Belle, and Badesso, in Western Ethiopia. Important agronomic data were collected and subjected to analysis using statistical analysis software (SAS). The combined analysis of variance (ANOVA) indicated highly significant differences ($P \leq 0.01$) among chickpea genotypes for grain yield and other agronomic traits except for number of seed per pod and branch per plant, indicating the existence of ample genetic variability among present chickpea genotypes. Genotypic coefficient of variation (GCV) and phenotypic coefficient of variation (PCV) for the character such as grain filling duration, days to flowering and hundred seed weight, showed very small difference demonstrating that the observed variations for these traits were mostly due to genetic factors with little impact of environment. The higher heritability coupled with higher genetic advance was noted for 100 seed weight, number of pods per plant, number of seed per pod and grain yield signifying the ease of phenotype-based selection for the improvement of those traits.

Key words: Chickpea (*Cicer arietinum* L.), genetic variability, phenotypic coefficient of variation (PCV), genotypic coefficient of variation (GCV), selection.

INTRODUCTION

Among pulses, chickpea (*Cicer arietinum* L.) is the fourth leading grain legume primarily grown for food and feed as well as soil fertility amendment in Ethiopia (CSA, 2015). Besides, Ethiopia is known as secondary center of origins for different crop in general and for chickpea in particular.

Regardless of its importance and presence of immense genetic variability, its yield is by far below its potential. This yield gap can be altered through proper manipulation of variable genotypes and thereby used in a breeding program. The main objective of any breeding

programs in the world is to produce high-yielding and better-quality lines for release as cultivars to farmers and commercial growers. The prerequisite to achieving this goal is to find sufficient amount of variability.

For long-time, plant breeders were aware that predominant portion of phenotypic variation were accounted for by environmental variation than from genotypes and genotypes by environmental interactions. Thus, in-depth knowledge of genetic variation has paramount importance for yield and its component improvement in any crop as the observed variability is a combined effect of genetic, environments and various interactions between genes and environments, of which only the former one is heritable. Genotypic variability is, therefore, the component of variation which is due to the genotypic differences among individuals within a population and is the main concern of plant breeders all over the world (Tadesse et al., 2016).

Selection would generate promising genotypes only if the source germplasms used in crossing are genetically diverse (Ali et al., 2008). When genetically diverse parents were chosen for crossing, there is a likelihood of producing heterotic progenies and advance required genetic recombination.

However, the amount of genetic variability present in any genotypes will not merely dictate the selection of superior genotypes. Therefore, in addition to the existence of genetic variability, the extent to which the characters in questions are transmitted from parents to offspring is very important to draw realistic conclusion vis-à-vis superior selections (Falconer and Mackay, 1996). Transmission of a quantitative trait from parents to offspring which by itself is highly influenced by genotype and environment interaction and traits with low heritability might not be improved through selection. Additionally, the estimates of heritability alone do not lonely deliver an idea about the expected gain in the next generation but should be considered in conjunction with estimates of genetic advance, the change in the mean value among successive generations (Johnson et al., 1955; Shukla et al., 2006).

Assessment of genetic variability using appropriate tools such as genetic coefficient of variation, heritability estimates, and genetic advance is, therefore, undeniably indispensable to boost yield as well as the quality of any crop and/or forage in any breeding program (Ali et al., 2008).

The objective of the present study was, therefore, to assess and quantify the genetic variability of chickpea germplasm grounded on agro-morphological traits and evaluating heritability and genetic advance among desi type chickpea genotypes for yield and yield contributing characters.

MATERIALS AND METHODS

Experimental sites, designs and experimental materials

A field experiment was conducted at five locations viz., Shambu (09° 32'N 037° 04'E), Hawa Galan (08° 38' N 034° 50'E), Mata (08° 34' N 034° 44'E), Alaku Belle (08° 37'N 034° 42'E) and Badesso (034° 42'E 034°47'E), Western Ethiopia, during the 2016/2017 main cropping season. A total of 16 Desi type chickpea varieties viz., 8 cultivars released over three decades, 1 local variety and 7 advanced lines collected from Debre Zeit Agricultural Research Center (DZARC) were used (Table 1).

The experiment was laid out in a randomized complete block design with three replicates and with plot size of 3 m length and 1.8 m width. All other crop management practices and recommendations were applied uniformly to all varieties as recommended for the crop.

Statistical analysis

Days to 50% flowering, days to 90% maturity, grain filling duration, number of pods per plant, number of seed per pod, plant height, number of branches per plant, hundred seed weight, and grain yield data were collected as suggested in IBPGR, ICRISAT, ICARDA (1993) and were subjected to analysis using appropriate software. Analysis of variance (ANOVA) using randomized complete block design was computed for all measured characters using SAS (SAS Institute, 2003). Genotypic (σ^2g), genotype by location interaction (σ^2gl) and phenotypic variances (σ^2p) were obtained from the analysis of variance (ANOVA) table according to Comstock and Robinson (1952) as follows,

$$\text{Genotypic variance } (\sigma^2g) = \frac{M_{sg} - M_{se}}{lr}$$

Where, M_{sg} is mean square of genotype, M_{se} is mean square of error, l is number of locations and r is replication.

Genotype by location interaction variance was estimated as:

$$\text{Genotype by location interaction variance } (\sigma^2gl) = \frac{M_{sgl} - M_{se}}{r}$$

Where, M_{sgl} is mean square of genotype by location interaction, M_{se} is mean square of error, and r is replication.

$$\text{Phenotypic variance } (\sigma^2p) = \delta^2g + \delta^2e + \sigma^2gl$$

Where, σ^2g is genotypic variance, σ^2e environmental variance, and σ^2gl is genotype by location interaction.

Heritability in the broad sense (H^2) was estimated according to (Falconer and Mackay, 1996) as follows:

$$H^2 = \frac{\delta^2g}{\delta^2p} \times 100$$

Where, σ^2g is genotypic variance and σ^2p is phenotypic variance.

The mean values for genetic analyses were used to determine phenotypic coefficient of variation (PCV) Environmental coefficient of variations (ECV), and genotypic coefficient of variation (GCV), as described by Singh and Choudhary (1985) as follows:

$$\text{GCV} = \frac{\sqrt{\delta^2g}}{\bar{x}} \times 100$$

*Corresponding author. E-mail: birrollee@gmail.com. Tel: +251917096447.

Table 1. Passport description of the test genotypes.

Genotype code	Genotype name	Status	Year of release
G1	Akaki	Released	1995
G2	Dalota	Released	2013
G3	Dimtu	Released	2012
G4	Dubie	Released	1978
G5	Local	Local variety	-
G6	Mariye	Released	1985
G7	Minjar	Released	2010
G8	Natoli	Released	2007
G9	Teketay	Released	2013
G10	DZ-2012-CK-0032	Advanced line	-
G11	DZ-2012-CK-0034	Advanced line	-
G12	DZ-2012-CK-0233	Advanced line	-
G13	DZ-2012-CK-0237	Advanced line	-
G14	DZ-2012-CK-0312	Advanced line	-
G15	DZ-2012-CK-0313	Advanced line	-
G16	DZ-2012-CK-20113-2-0042	Advanced line	-

Where, σ^2g is genotypic variance and \bar{x} is grand mean of a character

$$PCV = \frac{\sqrt{\delta^2p}}{\bar{x}} \times 100$$

Where, σ^2p is phenotypic variance and \bar{x} is grand mean of a character.

Environmental coefficient of variations (ECV) was estimated using:

$$ECV = \frac{\sqrt{\delta^2e}}{\bar{x}} \times 100$$

Where, σ^2e is error variance and \bar{x} is grand mean of a character.

Genetic advance (GA) was calculated with the method suggested by Allard (1960); Singh and Choudhary (1985) as follows:

$$GA = K \cdot \sqrt{\delta^2p} \times H^2$$

Where, K is 2.06 at 5% selection intensity, $\sqrt{\delta^2p}$ is square root of phenotypic variance and H^2 is heritability.

Genetic advance as a percentage of the mean (GAM) was computed as:

$$GAM (\%) = \frac{GA}{\bar{x}} \times 100$$

Where, GA is genetic advance and \bar{x} is grand mean of a character.

RESULTS AND DISCUSSION

The combined analysis of variance (ANOVA) for grain yield and other agronomic traits of 16 chickpea genotypes grown at five locations in 2016/2017 is shown in Table 2.

The analysis of variance (ANOVA) indicated highly

significant differences ($P \leq 0.01$) among chickpea genotypes for grain yield and other agronomic traits except for seed per pod and branch per plant demonstrating the presence of ample genetic variability among chickpea genotypes. Once the presence of genetic variability is secured, it is the task of the breeder to separate the heritable portion from the non-heritable part to plan for the proper breeding program. This might give an opportunity to plant breeders to improvement of characters with high heritability via selection.

Splitting the total phenotypic variance into the genotypic, genotype by environment interaction, and error variances were done on their expected mean squares (EMS) as per the fixed effects model. The genotypic variance (σ^2g) accounts for low proportion of the total phenotypic variance (σ^2p) as compared to the variances due to error (σ^2e) in most of the traits studied (Table 3). This shows that the observed phenotypic differences were not alone due to inherent genetic differences among the genotypes but also the substantial portion was due to environmental factor and the interplay between environment and genotype. Genotypic variance, phenotypic variance, genotypic coefficient of variability (GCV), environmental coefficient of variation (ECV), phenotypic coefficient of variability (PCV), broad sense heritability, and genetic advance for nine (9) agronomic characters are shown (Table 3).

Estimates of phenotypic coefficients of variation (PCV) and genotypic coefficients of variation (GCV) ranged from 2.24 (for days to maturity) to 33.60% (for number of pod per plant) and 6.71 (for days to maturity) to 20.67% (for 100 seed weight), respectively. Likewise, estimates of environmental coefficient of variation ranged from 1.83 (for days to maturity) to 28.89 (for number of pods per

Table 2. Combined analysis of variance for grain yield and other agronomic traits of sixteen chickpea genotypes grown at Shambu, Hawa Galan, Badesso, Mata, and Alaku Belle in 2016.

Source of variation	Mean square								
	D2F	D2M	GF	PPP	SPP	PH	BR	HSW	GYL
Rep (2)	10.4 ^{ns}	43.83 ^{**}	96.53 ^{**}	84.84 ^{ns}	0.22 [*]	4.74 ^{ns}	3.32 ^{**}	22.8 [*]	2685.7 ^{***}
Rep (Loc) (8)	1.29 ^{ns}	97.22 ^{***}	107.8 ^{***}	552.12 ^{***}	0.18 ^{**}	187.2 ^{***}	2.55 ^{**}	10.7 ^{ns}	655.5 ^{***}
Loc (4)	1806.8 ^{***}	1032.2 ^{***}	2017.1 ^{***}	1546.8 ^{***}	0.35 ^{**}	5303.8 ^{***}	12.45 ^{***}	671.2 ^{***}	19905.6 ^{***}
Geno (15)	115.9 ^{***}	21.03 ^{***}	193.2 ^{***}	443.41 ^{***}	0.96 ^{ns}	226.3 ^{***}	1.36 [*]	568.4 ^{***}	1059.2 ^{***}
G × Loc (60)	11.26 [*]	13.25 ^{**}	27.89 ^{**}	95.88 ^{ns}	0.05 ^{ns}	79.4 ^{***}	0.61 ^{ns}	12.8 ^{**}	292.7 ^{***}
Error (150)	6.67	6.39	13.02	81.76	0.06	30.94	0.69	5.9	131.7
Cv	4.09	1.82	4.78	28.88	20.6	10.73	19.98	8.2	21.76
R	0.90	0.86	0.87	0.6	0.43	0.869	0.56	0.9	0.86

***= Very highly significant at $P \leq 0.001$; ** = highly significant at $P \leq 0.01$; * = significant at $P \leq 0.05$; ns = not significant at $P = 0.05$; a Numbers in parentheses are degrees of freedom associated with the corresponding source of variation; Rep (Loc) = Replications within locations; D2F= Days to flowering; D2M= Days to maturity; GF= grain filling duration; PPP=pods per plant; SPP=seed per pod; PH=plant height BR=branch per plant; HSW=hundred seed weight; GYL= grain yield.

Table 3. Variance components, genotypic, environment, and phenotypic coefficients of variation, and heritability estimates of grain yield and other agronomic traits of sixteen chickpea genotypes grown at Shambu, Hawa Galan, Badesso, Mata and Alaku Belle in 2016/2017.

Trait	σ^2_g	σ^2_{gl}	σ^2_e	σ^2_p	GCV (%)	ECV	PCV (%)	H ² (%)	GA	GAM
Days to flowering	7.28	1.53	6.67	15.48	4.27	4.08	6.23	47.04	3.81	6.03
Days to maturity	0.98	2.29	6.39	9.65	0.71	1.83	2.24	10.11	0.65	0.47
Grain filling duration	12.01	4.96	13.02	29.99	4.6	4.79	7.27	40.05	4.52	5.99
Pods per plant	24.11	4.71	81.76	110.58	15.69	28.89	33.60	21.8	4.72	15.08
Seed per pod	0.06	0.00	0.06	0.12	19.91	19.91	27.77	51.43	0.37	29.84
Plant height	13.02	16.14	30.94	60.11	6.97	10.74	14.96	21.67	3.46	6.68
Branch per plant	0.04	0.03	0.69	0.71	5.07	19.92	20.18	6.31	0.11	2.64
Hundred seed weight	37.49	2.27	5.97	45.73	20.67	8.25	22.82	81.98	11.42	38.54
Grain yield	0.06	0.54	0.13	0.25	14.93	21.62	29.74	25.2	0.26	15.59

σ^2_g = genotypic variance, σ^2_{gl} = variance due to $g \times l$ interaction, σ^2_e = error variance, σ^2_p = phenotypic variance, GCV = genotypic coefficient of variation, ECV = environmental coefficient of variation, PCV = phenotypic coefficient of variation, H² = heritability in broad sense, GA = genetic advance, GAM = genetic advance as the percentage of the mean.

plant) (Table 3). Ali et al. (2010) also reported lowest phenotypic and environmental coefficient of variation for days to maturity in chickpea.

Deshmukh et al. (1986) classified PCV and GCV values as low (0-10%), moderate (10 - 20%) and high (>20%). Reliant on this suggestion, high GCV was observed for 100 seed weight, and moderate GCV was observed for number of seed per pod, number of pods per plant, and grain yield. The low value of a genotypic coefficient of variation (GCV) and phenotypic coefficient of variation (PCV) was noted for characters such as days to 50% flowering and days to maturity. Similar result was reported in numerous findings (Panwar and Singh, 2000; Bergale et al., 2001; Dwivedi et al., 2008; Khodadadi et al., 2011; Kumar et al., 2011).

The result of GCV expresses the true genetic potential of the genotypes. In the present study, the PCV values were higher than the corresponding GCV suggesting the

existence of substantial environmental variation. This result was completely in agreement with the result of combined analysis of variance (ANOVA) table. However, traits such as grain filling duration, days to flowering and 100 seed weight, which showed very small difference indicating the observed variations for these traits were mostly due to genetic factors with little environmental factors.

Estimation of broad sense heritability, genetic advance (at 5% selection intensity) and the corresponding genetic advance as a percent of the mean were estimated for each measured character. Heritability estimates are classified as low (5-10%), medium (10-30%) and high (30-60%) (Dabholkar, 1992). The broad sense heritability (H²) values more than 80% was observed for 100 seed weight. This implies that the expected gain from selection would be high if this trait is used as selection criteria in chickpea breeding.

Traits having high heritability combined with genetic advance could result in a better genetic gain through selection since the variation that prevails in such a trait is due to additive gene action (Johnson et al., 1955; Vimal and Vishwakarma, 1998). Hundred seed weight, number of pods per plant, number of seed per pod and grain yield showed high heritability combined with high genetic advance as a percentage of the mean which could be used as a powerful tool in phenotypic selection as such characters are controlled by the additive genes and less influenced by the environment.

Genetic gain (GAM) that could be expected from selecting the top 5% of the genotypes as a percent of the mean, varied from 0.47% for days to maturity to 38.54% for 100 seed weight. According to Johnson et al. (1955), genetic advance as a percent of the mean was categorized as high ($\geq 20\%$), moderate (10-20%) and lower (0-10%). Depending on this delineation, number of seed per pod and 100 seed weight showed high genetic advance as a percent of the mean, whereas number of pods per plant and grain yield showed moderate genetic advance as a percent of the mean. Therefore, selection based on traits with a high-level genetic advance as a percent of the mean will result in the improvement of the performance of the genotypes.

Conclusion

It can be concluded that direct selection can be done for most of the yield attributing traits since it exhibited high genetic variability and high range of variation. Traits with high heritability and genetic advance such as 100 seed weight, number of seed per pod and grain yield will grant superior genotypes through phenotypic based selection.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

ACKNOWLEDGMENTS

The authors are grateful for the chickpea peer grant scheme (USAID) and Oromia Agricultural Research Institute for financial support.

REFERENCES

Ali Q, Ahsan M, Farooq J (2010). Genetic variability and trait association in chickpea (*Cicer arietinum* L.) genotypes at seedling stage. *Electron. J. Plant Breed.* 1(3):334-341.

- Ali Y, Atta BM, Akhter J, Monneveux P, Lateef Z (2008). Genetic variability, association and diversity studies in wheat (*Triticum aestivum* L.) germplasm. *Pak. J. Bot.* 40(5):2087-2097.
- Allard I (1960). Principles of Plant Breeding, Chapter 6 through Chapter 9, University of California, Davis.
- Bergale S, Mridulla B, Holkar AS, Ruwali KN, Prasad SVS (2001). Genetic variability, diversity and association of quantitative traits with grain yield in bread wheat (*Triticum aestivum* L.). *Madras Agric. J.* 88(7):457-460.
- Comstock RE, Robinson HF (1952). Genetic parameters, their estimation and significance. In: Proceedings of the 6th International Grassland Congress 1:248-291.
- Central Statistical Agency (CSA) (2015). Crop and livestock product utilization. Statistical Bulletin. Addis Ababa, Ethiopia.
- Dabholkar AR (1992). Elements of Biometrical Genetics. Concept Publishing Company, New Delhi 110059. P 431.
- Deshmukh SN, Basu MS, Reddy PS (1986). Genetic variability, character association and path coefficients of quantitative traits in Virginia bunch varieties of groundnut. *Indian J. Agric. Sci.* 56:816-821.
- Dwivedi SL, Upadhyaya HD, Stalker HT, Blair MW, Bertoli DJ, Nielen S, Ortiz R (2008). Enhancing crop gene pools with beneficial traits using wild relatives. *Plant Breed. Rev.* 30:179-187.
- Falconer DS, Mackay TC (1996). Introduction to quantitative genetics. John Wiley and Sons. Inc. pp. 313-320.
- IBPGR, ICRISAT, ICARDA (1993). Descriptors for Chickpea (*Cicer arietinum* L.). International Board for Plant Genetic Resources, Rome, Italy; International Crops Research Institute for the Semi-Arid Tropics, Patancheru, India; International Center for Agricultural Research in the Dry Areas, Aleppo, Syria.
- Johnson HW, Robinson HF, Comstock RE (1955). Estimates of genetic and environmental variability in soybeans. *Agron. J.* 47(7):314-318.
- Khodadadi M, Fotokani MH, Miransari M (2011). Genetic diversity of wheat (*Triticum aestivum* L) genotypes based on clusters and principal component analyses for breeding strategies. *Aust. J. Crop Sci.* 5(1):17-22.
- Kumar S, Singh AK, Singh R (2011). Diversity analysis of wheat genotypes based on morpho-physiological and RAPD markers in relation to moisture stress. *Vegetos* 24(2):217-224.
- Panwar D, Singh I (2000). Genetic variability and character association of some yield components in winter x spring nursery of wheat. *Adv. Plant Sci.* 8(1):95-99.
- SAS Institute (2003). SAS/STAT guide for personal computers, version 9.1 edition. Cary, NC: SAS Institute Inc.
- Shukla S, Bhargava A, Chatterjee A, Srivastava A, Singh SP (2006). Genotypic variability in vegetable amaranth (*Amaranthus tricolor* L) for foliage yield and its contributing traits over successive cuttings and years. *Euphytica* 151(1):103-110.
- Singh RK, Chaudhary BD (1985). Analysis in biometrical genetics. Kalyani Publisher, New Delhi, India.
- Tadesse M, Fikre A, Eshete M, Girma N, Korbu L, Mohamed R, Bekele D, Funga A, Ojiewo CO (2016). Correlation and Path Coefficient Analysis for Various Quantitative Traits in Desi Chickpea Genotypes under Rainfed Conditions in Ethiopia. *J. Agric. Sci.* 8(12):112-118.
- Vimal SC, Vishwakarma SR (1998). Heritability and genetic advance in barley under partially reclaimed saline-sodic soil. *Rachis* (1-2): 56-57.



International Journal of Genetics and Molecular Biology

Related Journals Published by Academic Journals

- *Biotechnology and Molecular Biology Reviews*
- *African Journal of Biotechnology*
- *African Journal of Microbiology Research*
- *African Journal of Biochemistry Research*
- *Journal of Computational Biology and Bioinformatics Research*
- *International Journal for Biotechnology and Molecular Biology Research*
- *Journal of Developmental Biology and Tissue Engineering*

academicJournals