

OPEN ACCESS

Journal of **Yeast and Fungal Research**



November 2018
ISSN 2141-2413
DOI: 10.5897/JYFR
www.academicjournals.org



**ACADEMIC
JOURNALS**
expand your knowledge

ABOUT JYFR

The Journal of Yeast and Fungal Research (JYFR) (ISSN 2141-2413) is published Monthly (one volume per year) by Academic Journals.

Journal of Yeast and Fungal Research (JYFR), provides rapid publication (monthly) of articles in all areas of the subject such as Yeast physiology, fermentation and biotechnology, Bioremediation, Ethanol fermentation, economic importance of yeast etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in JYFR are peer-reviewed.

Contact Us

Editorial Office: jyfr@academicjournals.org

Help Desk: helpdesk@academicjournals.org

Website: <http://www.academicjournals.org/journal/JYFR>

Editor

Prof. Viroj Wiwanitkit, M.D.

*Wiwanitkit House, Bangkhae,
Bangkok Thailand 10160.
Visiting Prof. Tropical Medicine,
Hainan Medical College,
Hainan China.*

Associate Editors

Dr. Wolfram Siede,

*Department of Cell Biology and Anatomy
University of North Texas
Health Science Center.*

Dr. Mohsen Asker

*Microbial Biotechnology Dept.
National Research Centre
Cairo, Egypt.*

Prof. Chester R. Cooper, Jr.

*Youngstown State University
One University Plaza
Youngstown, Ohio.*

Prof. Fukai Bao

*Department of Microbiology and immunology,
Kunming Medical University
Yunnan, P. R. of China*

Dr. Raviraja N Seetharam

*Department of Oncology,
Montefiore Medical Center / Albert Einstein Cancer
Center,
Hofheimer Room No. 413,
E 210th St, Bronx, NY.*

Dr. Linghuo Jiang

*Tianjin Medical University,
Tianjin Research Center of Basic Medical Sciences,
China.*

Editorial Board

Dr. Jose Guedes de Sena Filho

*Federal University of Paraiba State Brazil/
University of Oklahoma.*

Dr. Fabien C.C. Hountondji

*Agriculture and Livestock Research/Ministry of
Agriculture,
Salalah, Oman.*

Dr. Zhenjia Chen

*Instituto de Tecnologia Química e Biológica (ITQB),
Universidade Nova de Lisboa, Oeiras,
Portugal.*

Dr. Bankole Munir Akinwale

*Ministry Of Health
Lagos State,
Nigeria.*

Dr. Yiguang Wang

*Institute of Medicinal Biotechnology,
CAMS&PUMC 1
TiantanXili, Beijing, China.*

Dr. Shobha D. Nadagir

*Dept of Microbiology. Karnatak Institute of
Medical Sciences,
Hubli. Karnatak State,
India.*

Isaiah Masinde Tabu

*Egerton University,
Egerton,
Kenya.*

Prof. Tzi Bun Ng

*School of Biomedical Sciences
The Chinese University of Hong Kong
Hong Kong,
China.*

Prof. Cristóbal Noé Aguilar González

*Group of Bioprocesses
Food Research Department
School of Chemistry
Universidad Autónoma de Coahuila
México.*

Prof. Caihong Dong

*State Key Laboratory of Mycology
Institute of Microbiology
Chinese Academy of Sciences
China.*

Dr. Tine Grebenc

*Department of Forest Physiology and Genetics
Slovenian Forestry Institute
Slovenia.*

Journal of Yeast and Fungal Research

Table of Content: Volume 9 Number 4 November, 2018

ARTICLE

Anticandidosic activity of selected medicinal plants from Côte d'Ivoire

Henriette Bosson-Vanga, Kpongbo Etienne Angora, Abibatou Konaté,
Akoua Valérie Bédia-Tanoh, Sébastien Miézan, Pulchérie Kiki-Barro,
Kondo Fulgence Kassi, Vincent Djohan, William Yavo and Hervé Menan

Full Length Research Paper

Anticandidosic activity of selected medicinal plants from Côte d'Ivoire

Henriette Bosson-Vanga^{1,2*}, Kpongbo Etienne Angora¹, Abibatou Konaté¹, Akoua Valérie Bédia-Tanoh¹, Sébastien Miézan¹, Pulchérie Kiki-Barro¹, Kondo Fulgence Kassi^{1,3}, Vincent Djohan¹, William Yavo¹ and Hervé Menan^{1,3}

¹Department of Parasitology, Mycology, Animal Biology and Zoology, Felix-Houphouët-Boigny University, BPV 34, Abidjan, Côte d'Ivoire.

²Department of Parasitology, Mycology, Institut Pasteur de Côte d'Ivoire, Abidjan, Côte d'Ivoire.

³Laboratory of Parasitology and Mycology of the Diagnosis and Research Centre on AIDS and the Others Infectious Diseases, 01 BPV 13, Abidjan, Côte d'Ivoire.

Received 11 October, 2018 ; Accepted 22 November, 2018

Superficial candidiasis is a real public health problem, especially in immunocompromised people and HIV-infected patients. Due to the toxic handicap and the high cost of antifungal drug treatment, people use medicinal plants, which are relatively more accessible. Medicinal plants are an important source of novel antimicrobial agents. This study aims to determine the *in vitro* antimicrobial activity of aqueous and methanolic extracts from different parts of plants species against *Candida albicans*. Six plants species from the flora of Côte d'Ivoire were screened for anticandidal activities by ethnobotanical study. These plants species were: *Bridelia ferruginea*, *Citrus aurantium*, *Pycnanthus angolensis*, *Desmodium adscendens*, *Zanthoxylum zanthoxyloides* Lam. and *Mareya micrantha* L. Methanolic and aqueous extracts from the powder of leaves and/or barks samples of each species were tested using the bioautographic method in F254 glass silica gel plate, with Miconazole as reference. Bioautography assay was used to account antifungal compounds in plant extracts. *Bridelia ferruginea*, *Pycnanthus angolensis*, and *Zanthoxylum zanthoxyloides* Lam showed good activity. The minimal inhibitory concentrations (MIC) values against *Candida albicans* ranged between 12.5 and 100 mg/ml. Bioautography results demonstrated that active chemical compounds of the alcoholic extracts of the plants (flavonoids, alkaloids, polyterpenes, polyphenols and sterols) were responsible for antimicrobial activity. The Ivorian pharmacopoeia is full of medicinal remedies. Three plants of our study have shown a good activity on strains of *Candida albicans*.

Key words: *Candida albicans*, antifungal activity, plants, bioautography.

INTRODUCTION

Candida yeast genus is the most common opportunistic pathogen among fungal infections causing high mortality and morbidity. It involves yeast pathogenic isolated from skin, mouth, intestinal tract, and vagina. Over the past few decades, the incidence of candidiasis has increased

especially with the growing number of immunocompromised patients (Essid et al., 2017). Resistance of *Candida* yeast genus to current medications, including the azoles, was reported (Arendrup, 2013). The search for new alternative

strategies is therefore important in order to fight *Candida* infections. Phytotherapy represents a more effective and less toxic alternative than standard drugs as anti-candida agents (Alves et al., 2014; Teodoro et al., 2015). Traditional medicine (TM) is often the primary mode of health care used by population in many countries due to poverty. It is an important and often underestimated part of health care. More than 80% of the population in Africa use TM to serve their health needs, according to the World Health Organisation (Anon n.d.). Plants are a primary source of new natural medicinal products. These traditional remedies use any part of plant, such as barks, leaves, flowers, roots and seeds. These remedies can be prepared from a single plant or a combination of different plants. Medicinal plants contain bioactive compounds with healing ability. These include saponins, tannins, essential oils, flavonoids, alkaloids and other chemical compounds found as secondary metabolites in plants (Mahlo et al., 2016). Africa medicinal flora is rich and medicinal plant knowledge nowadays has a great development especially in Côte d'Ivoire. The country benefits from an excellent floristic biodiversity and the indigenous people have old knowledge of the medicinal plant uses. Antimicrobial activities of extracts from *Bridelia ferruginea* (Afolayan et al., 2018; Alowanou et al., 2015), *Citrus aurantium* (Metoui et al., 2015; Ruiz-Pérez et al., 2016), *Pycnanthus angolensis* (Kuete et al., 2011), *Zanthoxylum zanthoxyloides* Lam. (Tine et al., 2017) and many other herbs have also been reported. Indeed, Kra Mathieu et al. showed that the hydroethanolic extracts from the four *Terminalia* species produced minimal fungicidal concentrations (MFC) values that are lower than the MFC value of ketoconazole (Mathieu et al., 2014). Other plant species like *Borreria latifolia*, *Borreria verticillata*, *Erigeron floribundus*, *Euphorbia hirta*, *Turraea heterophylla* and *Vernonia colorata* were tested (Bi et al., 2007).

In this study, we chose six Ivoirian plants species used by traditional healers for the treatment of superficial candidiasis: *Bridelia ferruginea*, *Citrus aurantium*, *Pycnanthus angolensis*, *Desmodium adscendens*, *Zanthoxylum zanthoxyloides* Lam. and *Mareya micrantha* L. These plants have never been evaluated in Côte d'Ivoire for their antifungal properties. We investigated their aqueous and methanolic extracts.

MATERIALS AND METHODS

Plants collection

For the identification of medicinal plants used to treat superficial

candidiasis, we bought them from sellers in different markets of Abidjan (Abobo, Adjamé and Treichville). These markets are located in the popular neighborhoods of the city of Abidjan. Different plants and recipes were identified from these sellers according to their skills. The formulas were combinations of plants used together. Then, the National Center of Floristry in Abidjan located at the University Félix Houphouët Boigny helped us to identify correctly all the plants bought in the markets. The selected plants for this study met the following criteria: (i) used by population in the treatment of superficial candidiasis; (ii) existed in the African pharmacopoeia; (iii) widespread in Côte d'Ivoire and easily accessible. Thus, six plants species from traditional medicine were selected namely: *B. ferruginea*, *C. aurantium*, *P. angolensis*, *D. adscendens*, *Z. zanthoxyloides* Lam. and *M. micrantha* L. The plant materials were the bark of *Pycnanthus angolensis*, the creeping stems of *Desmodium adscendens*, the roots of *Bridelia ferruginea* and the leaves of *C. aurantium*, *Z. zanthoxyloides* Lam. and *M. micrantha* L.

The plant materials were harvested in three parts of Côte d'Ivoire:

1. In the Banco forest, precisely on the slope overlooking the district "Anokoua" in the city of Abobo southern Abidjan (*D. adscendens*, *C. aurantium*, *P. angolensis* and *M. micrantha* L.),
2. in the city of Toumodi in central Côte d'Ivoire (*B. ferruginea*)
3. in the city of Korhogo in Northern Côte d'Ivoire (*Z. zanthoxyloides* Lam).

The plants materials were dried at room temperature (25°C) for about 8 weeks and then ground to fine powder. It was stored in airtight bottles in the dark until extraction to prevent oxidation.

Plant extraction

Preparation of total extracts of drugs

Each finely ground plant material (60 g) was extracted with 500 ml of solvents in erlenmeyer: water or methanol. Extracts (total aqueous, ethanolic 70% and residual extracts aqueous) were prepared according to the method described by Camara et al. (2016). Plant extracts were re-dissolved in acetone for microbiological assays and phytochemical analysis.

Phytochemical analysis

Colorimetric and precipitation reactions were used for phytochemical sorting; they highlight the following chemical groups: sterols and polyterpenes, polyphenols, flavonoids, tannins, quinonic substances and alkaloids. Chemical components of each extracts were analysed and used to determine their anticandidal activity.

C. albicans strains

Clinical strains were obtained from the parasitology and mycology laboratory of CeDRoS. The germ tube test and the chlamydo-spore formation assay were used for identification of *C. albicans* yeast by

*Corresponding author. E-mail: bossonhenriette@gmail.com.

Table 1. Parts of plants and their uses.

Plants	Family	Used parts	Form of use
<i>Bridelia ferruginea</i>	Euphorbiaceae	Roots	Maceration, decoction
<i>Citrus aurantium</i>	Rutaceae	Leafs, fruits	Cataplasm, maceration, decoction
<i>Desmodium adscendens</i>	Fabaceae	Leafs, Roots bark	Decoction, toothbrush
<i>Pycnanthus angolensis</i>	Myristicaceae	Stem bark	Decoction, cataplasm
<i>Mareya micrantha</i>	Euphorbiaceae	Leafs	Decoction, cataplasm
<i>Zanthoxylum zanthoxyloides</i>	Rutaceae	Leafs	Decoction, toothbrush

using young colonies (24 h).

Anticandidotic activity assay: Bioautography

A young *C. albicans* colony was collected and homogenized in 9 mL of sterile distilled water. The first inoculum was obtained by mixing 0.5 mL of the yeast suspension to 25 mL of Sabouraud medium liquid incubated at 30°C for 15 h. The second inoculum was obtained by mixing 0.5 mL of the first one to 25 mL of Sabouraud medium liquid incubated at 30°C for 8 h. Both incubations (15 and 8 h) allowed the yeasts to be in exponential growth phase.

In the next step, plant extracts (100 mg) were serially diluted at 50% with chloroform (1 mL). Thirty microliters of each dilution (S1, S2, S3, and S4) of the total plant extracts and 50 µL of the control (Miconazol: T) are put on a silica plate. The deposits are then quickly dried with an electric dryer. Sabouraud maltose agar medium was regenerated at 40°C and was mixed with the inoculum at the proportion of 0.5 mL of inoculum in 25 mL of malted Sabouraud medium. Then, 20 mL of this mixture was poured on the silica plate bearing the deposits of the serial extracts dilutions (alcoholic and aqueous). This step was performed on a heating plate to avoid solidification of the cast medium. The silica plate was incubated in the wet chamber at 30° for 24 h. After this time, the silica plate G was removed from the oven and sprayed with 4 mL of thiazolyl blue solution (2.5 mg/mL of thiazolyl blue in sterile distilled water). At the end, this silica gel plate was incubated again for 4 h in the wet chamber at 30°C. The reading was made with revelatory. The minimal inhibit concentration (MIC) was recorded as the lowest concentration of the extract that inhibited antifungal growth.

The antifungal test of each compound (alkaloids, polyterpenes, sterols and flavonoids) was carried out in the same way as the anticandidotic test on total plant extracts. These tests were performed three times to confirm the results.

RESULTS

The ethnobotanical study of plants with presumed activity on *C. albicans* for medicinal plant sellers identified 20 plants. Overall, medicinal plants are used in combination

with other plants for the same treatment. We therefore selected those that returned in the majority of the preparations. In the current study, six medicinal plants were investigated for their antifungal activities against *C. albicans*. Table 1 shows the parts of the plants used and their use. The commonly methods used remain decoction, which consists of introducing plant parts into boiling water.

Phytochemical sorting allowed extraction of chemical groups (sterols and polyterpenes, polyphenols, flavonoids, tannins, quinonic substances and alkaloids). Table 2 shows the chemical composition of the aqueous and alcoholic extracts. Sterols and polyterpenes, polyphenols, flavonoids and alkaloids were mainly found.

In the present study, the extracts of six medicinal plants were investigated for their antifungal activities against *C. albicans*. The details of the results are shown in Table 3. *C. albicans* strains are susceptible to alcoholic extract of three plants (*Bridelia ferruginea*, *Pycnanthus angolensis* and *Z. zanthoxyloides* Lam. The minimal inhibition concentration (MIC) values against *C. albicans* were 12.5 mg/ml for the three plants. This activity seems to be the result of chemical compounds extracted during phytochemical sorting. Determination of the active chemical compounds of the alcoholic extract showed that total flavonoid of *B. ferruginea*, total quinones of *P. angolensis* and total alkaloids of *Z. zanthoxyloides* Lam. gave an inhibition zone similar to the Miconazole one.

DISCUSSION

C. albicans infections are widespread. Under normal conditions, this fungus is harmless and found in the intestinal walls, mouth, vagina and skin. If the natural balance is upset, depending on the soil on which the *Candida* proliferates, it could cause superficial

Table 2. Phytochemical constituents.

Plants	Extract	Sterols / Polyterpenes	Polyphenols	Flavonoids	Tanins		Quinonics Substances	Alkaloids	
					Gal	Cat		D	B
<i>Bridelia ferruginea</i>	H ₂ O	+	+	-	-	-	-	+	+
<i>Citrus aurantium</i>	OH	+	+	+	-	-	-	+	+
<i>Mareya micrantha L.</i>	H ₂ O	-	+	-	-	+	-	-	-
<i>Desmodium adscendens</i>	OH	-	+	-	+	+	-	-	-
<i>Pycnanthus angolensis</i>	H ₂ O	+	+	+	-	-	-	+	+
<i>Zanthoxylum zanthoxyloides</i>	OH	+	+	+	-	-	+	+	+
	H ₂ O	-	-	+	-	-	-	+	+
	OH	+	+	+	-	-	-	+	+
	H ₂ O	-	+	+	-	-	-	-	-
	OH	-	+	+	-	-	-	+	+

+, positive reaction (presence of the searched chemical group); -, negative reaction (absence of the searched chemical group); gal, gallic acid; cat, catechin; D, Dragendorff; B, Bouchardat.

Table 3. Plants with anticandidal activity (*Bridelia ferruginea*, *Pycnanthus angolensis*, and *Zanthoxylum zanthoxyloides* Lam.).

Plants	Extract	Concentrations (mg/l)	Inhibition zone (mm)
<i>Bridelia ferruginea</i>	H ₂ O	100	0
		12.5	0
	OH	100	13 ± 2
		12.5	11 ± 2
<i>Pycnanthus angolensis</i>	H ₂ O	100	0
		12.5	0
	OH	100	13 ± 5
		12.5	11 ± 2
<i>Zanthoxylum zanthoxyloides</i>	H ₂ O	100	0
		12.5	0
	OH	100	13 ± 2
		12.5	11 ± 2
<i>Miconazole</i>		100	11 ± 4

candidiasis (cutaneous or mucous membrane) or deep candidiasis. Faced with this pathology, traditional medicine therapeutic is usually used in our regions. The current study aims to evaluate the anticandidal effectiveness of six plants and their main components. *Bridelia ferruginea*, *Citrus aurantium*, *Pycnanthus angolensis*, *Desmodium adscendens*, *Zanthoxylum zanthoxyloides* Lam. and *Mareya micrantha L.* are from African particularly Ivorian pharmacopoeia and their antifungal activities are known. *Bridelia ferruginea* for example is used for antiamebic, antianemic, antibacterial, anticonvulsant, anti-diabetic, antidiarrhoeal, antihelminthic,

anti-inflammatory, antimalarial, antinociceptive, antiviral, hypoglycemic and for abdominal pain (Awodele et al., 2015; Ngueyem et al., 2009). *Pycnanthus angolensis* showed antimalarial, antibacterial, hypoglycemic and anticancer activities (Fort et al., 2000; Gbolade and Adeyemi, 2008; Abrantes et al., 2008). *Z. zanthoxyloides* Lam. also has an action on hematological parameters and oxidation (Ogunbolude et al., 2014; Ikumawoyi et al., 2016). Decoction is the most commonly used method of preparation and is obtained by boiling plants in water. Otherwise, this method according to the stakeholders surveyed would be able to use all the plants' potential.

Most traditional healers use water for their decoctions because water is the only solvent available. A major challenge of using water for extraction is that non-polar bioactive compounds cannot be extracted. The type of solvent used in the extraction procedure determines the success of isolating compounds from plant material (Masoko and Nxumalo, 2013). We used two types of solvents in this study: water and methanol. The phytochemical composition revealed that both solvents were able to extract the same chemical compounds from the plants studied. However, in our study, the tests from extracts for antifungal activities against *C. albicans* showed that only alcoholic extracts were active. Other studies reported that alcohols, particularly methanol, allowed the best extraction of active substances from plants (Arama et al., 2016; Masoko and Makgapeetja, 2015).

The bioautography assay was used to determine antifungal activity. This study highlighted the anti-*Candida* potency of three from the six studied plants (*Bridelia ferruginea*, *Pycnanthus angolensis* and *Z. zanthoxyloides Lam.*). Many studies showed that *Bridelia ferruginea*, *Pycnanthus angolensis* and *Zanthoxylum zanthoxyloides Lam.* have antimicrobial activities in particular antifungal effect (Afolayan et al., 2018; Alowanou et al., 2015; Kuete et al., 2011; Tine et al., 2017). Besides, chemical groups from these three anti-*Candida* plants contained flavonoids quinones and alkaloids responsible for this activity. The antifungal activity of these compounds has been reported in the literature. Similarly, many authors tested the anticandidotic activity of flavonoids (Seleem et al., 2017; Ozçelik et al., 2006; Orhan et al., 2010). Kamdem Wabo et al. tested the anticandidotic activity of a new quinones as Pycnanthuquinone C isolated from *Pycnanthus angolensis* (Wabo et al., 2007). Alkaloids have also been reported to possess antifungal activity (Orhana et al., 2007). The remaining three plant species (*Citrus aurantium*, *Desmodium adscendens*, and *Mareya micrantha L.*) showed no activity in the bioautography screening against *C. albicans*. This could be explained by the fact that some of the active compounds were volatile and evaporated during drying period prior to bioautography. Biological activity synergism between different compounds in the extracts is also a possible reason (Mahlo et al., 2016).

The results presented in this study are encouraging. A wider study is needed to identify the effective components, mode of action and possible toxic effect *in vivo* of these ingredients. It is therefore important to investigate the potential of these plants as novel antifungal agents, targeting the multidrug resistant fungi of clinical importance.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

ACKNOWLEDGMENTS

Authors are grateful to the National Center of Floristry, the parasitology and mycology laboratory of CeDReS and the Department of Parasitology-Mycology for their support.

REFERENCES

- Abrantes M, Mil-Homens T, Duarte N, Lopes D, Cravo P, Madureira M, Ferreira MJU (2008). Antiplasmodial activity of lignans and extracts from *Pycnanthus angolensis*. *Planta Medica* 74:1408-1412.
- Afolayan M, Srivedavyasasri R, Asekun OT, Familoni OB, Ross SA (2018). Chemical and biological studies on *Bridelia ferruginea* grown in Nigeria. *Natural Product Research* 1-5.
- Alowanou GG, Olounlade AP, Azando EVB, Dedehou V, Daga FD, Hounzangbe-adote M (2015). A review of *Bridelia ferruginea*, *Combretum glutinosum* and *Mitragina inermis* plants used in zootherapeutic remedies in West Africa: historical origins, current uses and implications for conservation. *Journal of Applied Biosciences* 87:8003-8014-8014.
- Alves CT, Ferreira ICFR, Barros L, Silva S, Azeredo J, Henriques M (2014). Antifungal activity of phenolic compounds identified in flowers from North Eastern Portugal against *Candida* species. *Future Microbiology* 9:139-146.
- Anon (N.D). WHO | WHO traditional medicine strategy: 2014-2023. WHO. Available at: http://www.who.int/medicines/publications/traditional/trm_strategy14_23/en/ [Accessed November 11, 2018].
- Arama PF, Anyango B, Akenga T, Nyunja R, Khasabuli D (2016). In vitro antifungal activity of methanolic extracts of different senna *didymobotrya* (fresen.) h.s. irwin & barneby plant parts. *African journal of traditional, complementary, and alternative medicines: AJTCAM* 13:168-174.
- Arendrup MC (2013). *Candida* and candidaemia. Susceptibility and epidemiology. *Danish Medical Journal* 60:B4698.
- Awodele O, Amagon KI, Agbo J, Prasad MNV (2015). Toxicological evaluation of the aqueous stem bark extract of *Bridelia ferruginea* (Euphorbiaceae) in rodents. *Interdisciplinary Toxicology* 8:89-98.
- Bi FT, Kouame NF, Favel A, Fallague K (2007). Antifungal activity of some plants of the Ivorian flora. *Sciences & Nature* 4:117-122.
- Camara D, Bene K, Gnahoue G, Fofie NBY, Zirih GN (2016). Ethnobotanical Study, Evaluation of the Antifungal Activity on *Candida Albicans* and Toxicity on Hff Cells of *Bersama Abyssinica* (Fresen.), A Plant Of The Ivorian Pharmacopee. *European Scientific Journal*, ESJ 12:171.
- Essid R, Hammami M, Gharbi D, Karkouch I, Hamouda TB, Elkahoui S, Limam F, Tabbene O (2017). Antifungal mechanism of the combination of *Cinnamomum verum* and *Pelargonium graveolens* essential oils with fluconazole against pathogenic *Candida* strains. *Applied Microbiology and Biotechnology* 101:6993-7006.
- Fort DM, Ubillas RP, Mendez CD, Jolad SD, Inman WD, Carney JR, Chen JL, Ianiro TT, Hasbun C, Bruening RC, Luo J, Reed MJ, Iwu M, Carlson TJ, King SR, Bierer DE, Cooper R (2000). Novel antihyperglycemic terpenoid-quinones from *Pycnanthus angolensis*. *The Journal of Organic Chemistry* 65:6534-6539.
- Gbolade AA, Adeyemi AA (2008). Investigation of in vitro anthelmintic activities of *Pycnanthus angolensis* and *Sphenocentrum jollyanum*. *Fitoterapia* 79:220-222.
- Ikumawoyi V, Awodele O, Rotimi K, Fashina A (2016). Evaluation of the effects of the hydro-ethanolic root extract of *zanthoxylum zanthoxyloides* on hematological parameters and oxidative stress in cyclophosphamide treated rats. *African journal of traditional, complementary, and alternative medicines: AJTCAM* 13:153-159.
- Kuete V, Nono ECN, Mkounga P, Marat K, Hultin PG, Nkengfack AE (2011). Antimicrobial activities of the CH₂Cl₂-CH₃OH (1:1) extracts and compounds from the roots and fruits of *Pycnanthus angolensis* (Myristicaceae). *Natural Product Research* 25:432-443.

- Mahlo SM, Chauke HR, McGaw L, Eloff J (2016) Antioxidant and antifungal activity of selected medicinal plant extracts against phytopathogenic fungi. *African journal of traditional, complementary, and alternative medicines*: AJTCAM 13:216-222.
- Masoko P, Makgapeetja DM (2015) Antibacterial, antifungal and antioxidant activity of *Olea africana* against pathogenic yeast and nosocomial pathogens. *BMC complementary and alternative medicine* 15:409.
- Masoko P, Nxumalo KM (2013). Validation of antimycobacterial plants used by traditional healers in three districts of the limpopo province (South Africa). *Evidence-Based Complementary and Alternative Medicine: eCAM* 2013:586247.
- Mathieu KAK, Marcel AG, Djè DB, Sitapha O, Adama C, Joseph DA (2014). Anti-fungal activities of medicinal plants extracts of Ivorian pharmacopoeia. *Journal of Intercultural Ethnopharmacology* 3:159-166.
- Metoui N, Gargouri S, Amri I, Fezzani T, Jamoussi B, Hamrouni L (2015). Activity antifungal of the essential oils; aqueous and ethanol extracts from *Citrus aurantium* L. *Natural Product Research* 29:2238-2241.
- Ngueyem TA, Brusotti G, Caccialanza G, Finzi PV (2009). The genus *Bridelia*: a phytochemical and ethnopharmacological review. *Journal of Ethnopharmacology* 124:339-349.
- Ogunbolude Y, Ibrahim M, Elekofehinti OO, Adeniran A, Abolaji AO, Rocha JBT, Kamdem JP (2014). Effects of *Tapinanthus globiferus* and *Zanthoxylum zanthoxyloides* extracts on human leukocytes in vitro. *Journal of Intercultural Ethnopharmacology* 3:167-172.
- Orhan DD, Ozçelik B, Ozgen S, Ergun F (2010). Antibacterial, antifungal, and antiviral activities of some flavonoids. *Microbiological Research* 165:496-504.
- Orhana I, Ozçelik B, Karaoğlu T, Sener B (2007) Antiviral and antimicrobial profiles of selected isoquinoline alkaloids from *Fumaria* and *Corydalis* species. *Zeitschrift Fur Naturforschung. C, Journal of Biosciences* 62:19-26.
- Ozçelik B, Orhan I, Toker G (2006). Antiviral and antimicrobial assessment of some selected flavonoids. *Zeitschrift Fur Naturforschung. C, Journal of Biosciences* 61:632-638.
- Ruiz-Pérez NJ, González-Ávila M, Sánchez-Navarrete J, Toscano-Garibay JD, Moreno-Eutimio MA, Sandoval-Hernández T, Arriaga-Alba M (2016). Antimycotic Activity and Genotoxic Evaluation of *Citrus sinensis* and *Citrus latifolia* Essential Oils. *Scientific Reports* 6:25371.
- Seleem D, Pardi V, Murata RM (2017). Review of flavonoids: A diverse group of natural compounds with anti-*Candida albicans* activity in vitro. *Archives of Oral Biology* 76:76-83.
- Teodoro GR, Ellepola K, Seneviratne CJ, Koga-Ito CY (2015). Potential Use of Phenolic Acids as Anti-*Candida* Agents: A Review. *Frontiers in Microbiology* 6:1420.
- Tine Y, Diop A, Diatta W, Desjobert JM., Boye CSB, Costa J, Wélé A, Paolini J (2017). Chemical Diversity and Antimicrobial Activity of Volatile Compounds from *Zanthoxylum zanthoxyloides* Lam. according to Compound Classes, Plant Organs and Senegalese Sample Locations. *Chemistry & Biodiversity* 14. <http://dx.doi.org/10.1002/cbdv.201600125>
- Wabo HK, Tatsimo SN, Tane P, Connolly JD (2007). Pycnanthuquinone C: a new terpenoid-quinone from *Pycnanthus angolensis*. *Planta Medica* 73:187-189.

Related Journals:

