

OPEN ACCESS



# African Journal of **Biotechnology**

22 November 2018  
ISSN 1684-5315  
DOI: 10.5897/AJB  
[www.academicjournals.org](http://www.academicjournals.org)



**ACADEMIC  
JOURNALS**  
expand your knowledge

# About AJB

The African Journal of Biotechnology (AJB) is a peer reviewed journal which commenced publication in 2002. AJB publishes articles from all areas of biotechnology including medical and pharmaceutical biotechnology, molecular diagnostics, applied biochemistry, industrial microbiology, molecular biology, bioinformatics, genomics and proteomics, transcriptomics and genome editing, food and agricultural technologies, and metabolic engineering. Manuscripts on economic and ethical issues relating to biotechnology research are also considered.

## Indexing

[CAB Abstracts](#), [CABI's Global Health Database](#), [Chemical Abstracts \(CAS Source Index\)](#), [Dimensions Database](#), [Google Scholar](#), [Matrix of Information for The Analysis of Journals \(MIAR\)](#), [Microsoft Academic](#), [Research Gate](#)

## Open Access Policy

Open Access is a publication model that enables the dissemination of research articles to the global community without restriction through the internet. All articles published under open access can be accessed by anyone with internet connection.

The African Journals of Biotechnology is an Open Access journal. Abstracts and full texts of all articles published in this journal are freely accessible to everyone immediately after publication without any form of restriction.

## Article License

All articles published by African Journal of Biotechnology are licensed under the [Creative Commons Attribution 4.0 International License](#). This permits anyone to copy, redistribute, remix, transmit and adapt the work provided the original work and source is appropriately cited. Citation should include the article DOI. The article license is displayed on the abstract page the following statement:

This article is published under the terms of the [Creative Commons Attribution License 4.0](#)  
Please refer to <https://creativecommons.org/licenses/by/4.0/legalcode> for details  
about [Creative Commons Attribution License 4.0](#)

## **Article Copyright**

When an article is published by in the African Journal of Biotechnology, the author(s) of the article retain the copyright of article. Author(s) may republish the article as part of a book or other materials. When reusing a published article, author(s) should;

Cite the original source of the publication when reusing the article. i.e. cite that the article was originally published in the African Journal of Biotechnology. Include the article DOI

Accept that the article remains published by the African Journal of Biotechnology (except in occasion of a retraction of the article)

The article is licensed under the Creative Commons Attribution 4.0 International License.

A copyright statement is stated in the abstract page of each article. The following statement is an example of a copyright statement on an abstract page.

Copyright ©2016 Author(s) retains the copyright of this article.

## **Self-Archiving Policy**

The African Journal of Biotechnology is a RoMEO green journal. This permits authors to archive any version of their article they find most suitable, including the published version on their institutional repository and any other suitable website.

Please see <http://www.sherpa.ac.uk/romeo/search.php?issn=1684-5315>

## **Digital Archiving Policy**

The African Journal of Biotechnology is committed to the long-term preservation of its content. All articles published by the journal are preserved by [Portico](#). In addition, the journal encourages authors to archive the published version of their articles on their institutional repositories and as well as other appropriate websites.

<https://www.portico.org/publishers/ajournals/>

## **Metadata Harvesting**

The African Journal of Biotechnology encourages metadata harvesting of all its content. The journal fully supports and implement the OAI version 2.0, which comes in a standard XML format. [See Harvesting Parameter](#)

## Memberships and Standards



Academic Journals strongly supports the Open Access initiative. Abstracts and full texts of all articles published by Academic Journals are freely accessible to everyone immediately after publication.



All articles published by Academic Journals are licensed under the [Creative Commons Attribution 4.0 International License \(CC BY 4.0\)](#). This permits anyone to copy, redistribute, remix, transmit and adapt the work provided the original work and source is appropriately cited.



[Crossref](#) is an association of scholarly publishers that developed Digital Object Identification (DOI) system for the unique identification published materials. Academic Journals is a member of Crossref and uses the DOI system. All articles published by Academic Journals are issued DOI.

[Similarity Check](#) powered by iThenticate is an initiative started by CrossRef to help its members actively engage in efforts to prevent scholarly and professional plagiarism. Academic Journals is a member of Similarity Check.

[CrossRef Cited-by](#) Linking (formerly Forward Linking) is a service that allows you to discover how your publications are being cited and to incorporate that information into your online publication platform. Academic Journals is a member of [CrossRef Cited-by](#).



Academic Journals is a member of the [International Digital Publishing Forum \(IDPF\)](#). The IDPF is the global trade and standards organization dedicated to the development and

promotion of electronic publishing and content consumption.

## Contact

Editorial Office: [ajb@academicjournals.org](mailto:ajb@academicjournals.org)

Help Desk: [helpdesk@academicjournals.org](mailto:helpdesk@academicjournals.org)

Website: <http://www.academicjournals.org/journal/AJB>

Submit manuscript online <http://ms.academicjournals.org>

Academic Journals  
73023 Victoria Island, Lagos, Nigeria  
ICEA Building, 17th Floor,  
Kenyatta Avenue, Nairobi, Kenya.



## Editor-in-Chief

**Prof. N. John Tonukari**

Department of Biochemistry  
Delta State University  
Abraka,  
Nigeria.

**Ana I. L Ribeiro-Barros**

Department of Natural Resources,  
Environment and Territory  
School of Agriculture  
University of Lisbon  
Portugal.

**Estibaliz Sansinenea**

Chemical Science Faculty  
Universidad Autonoma De Puebla  
Mexico.

**Bogdan Sevastre**

Physiopathology Department  
University of Agricultural Science and  
Veterinary Medicine  
Cluj Napoca Romania.

**Parichat Phumkhachorn**

Department of Biological Science  
Ubon Ratchathani University  
Thailand.

**Mario A. Pagnotta**

Department of Agricultural and Forestry sciences  
Tuscia University  
Italy.

## Editorial Board Members

**Dr. Gunjan Mukherjee**

Agharkar Research Institute (ARI),  
Autonomous Institute of the Department of  
Science and Technology (DST) Government of  
India  
Pune, India.

**Prof. Dr. A.E. Aboulata**

Plant Pathology Research Institute (ARC)  
Giza, Egypt.

**Dr. S. K. Das**

Department of Applied Chemistry and  
Biotechnology  
University of Fukui  
Japan.

**Prof. A. I. Okoh**

Applied and Environmental Microbiology  
Research Group (AEMREG)  
Department of Biochemistry and Microbiology  
University of Fort Hare  
Alice, South Africa.

**Dr. Ismail Turkoglu**

Department of Biology Education  
Education Faculty  
Firat University  
Elazığ, Turkey.

**Dr. Huda El-Sheshtawy**

Biotechnological Application lab., Process,  
Design and Development  
Egyptian Petroleum Research Institute (EPRI)  
Cairo, Egypt.

**Prof. T. K. Raja**

Department of Biotechnology  
PSG College of Technology  
(Autonomous)  
Coimbatore India.

**Dr. Desobgo Zangue**

Steve Carly  
Food Processing and Quality Control  
University Institute of Technology  
(University of Ngaoundere) Cameroon.

**Dr. Girish Kamble**

Botany Department  
SRRL Science College Morshi India.

**Dr. Zhiguo Li**

School of Chemical Engineering  
University of Birmingham  
United Kingdom.

**Dr. Srecko Trifunovic**

Department of Chemistry  
Faculty of Science  
University of Kragujevac  
Serbia.

**Dr. Sekhar Kambakam**

Department of Agronomy  
Iowa State University USA.

**Dr. Carmelo Peter**

Bonsignore  
Department PAU – Laboratorio di  
Entomologia ed Ecologia Applicata  
Mediterranean University of Reggio  
Calabria  
Italy.



**Dr. Vincenzo Tufarelli**

Department of Emergency and Organ  
Transplant (DETO)  
Section of Veterinary Science and Animal  
Production  
University of Bari "Aldo Moro", Italy.

**Dr. Tamer El-Sayed Ali**

Oceanography Department  
Faculty of Science  
Alexandria University  
Alexandria, Egypt.

**Dr. Chong Wang**

College of Animal Science  
Zhejiang A&F University  
China.

**Dr. Christophe Brugidou**

Research Institute for Development (IRD)  
Center, France.

**Dr. Maria J. Poblaciones**

Department of Agronomy and Forest  
Environment Engineering  
Extremadura University,  
Spain.

**Dr. Anna Starzyńska-Janiszewska**

Department of Food Biotechnology  
Faculty of Food Technology  
University of Agriculture in Krakow  
Poland.

**Dr. Amlan Patra**

Department of Animal Nutrition  
West Bengal University of Animal and Fishery  
Sciences  
India.

**Dr. Navneet Rai**

Genome Center,  
University of California Davis, USA.

**Dr. Preejith Vachali**

School of Medicine  
University of Utah  
USA.

## Table of Content

**Characterization of fifteen microsatellite loci and genetic diversity analysis for the Ghanaian food security crop *Solenostemon rotundifolius* (Frafra potato)**

Lichun Hua, Denita Hadziabdic, Naalamle Amissah, Marcin Nowicki, Sarah L. Boggess, Margaret Staton, Nianjun Teng and Robert N. Trigiano

*Full Length Research Paper*

# Characterization of fifteen microsatellite loci and genetic diversity analysis for the Ghanaian food security crop *Solenostemon rotundifolius* (Frafra potato)

Lichun Hua<sup>1,4</sup>, Denita Hadziabdic<sup>1</sup>, Naalamle Amissah<sup>2</sup>, Marcin Nowicki<sup>1</sup>, Sarah L. Boggess<sup>1</sup>, Margaret Staton<sup>1</sup>, Nianjun Teng<sup>3</sup> and Robert N. Trigiano<sup>1\*</sup>

<sup>1</sup>University of Tennessee Institute of Agriculture, Department of Entomology and Plant Pathology, University of Tennessee, Knoxville, Tennessee, USA.

<sup>2</sup>School of Agriculture, Department of Crop Science, University of Ghana, Accra, Ghana.

<sup>3</sup>College of Horticulture, Nanjing Agricultural University, Nanjing, People's Republic of China.

<sup>4</sup>Department of Plant Development and (Epi) Genetics, Swammerdam Institute for Life Sciences, University of Amsterdam, The Netherlands.

Received 2 October, 2018; Accepted 20 November, 2018

**Frafra potato (*Solenostemon rotundifolius*) is an important food crop indigenous to Western Africa and a member of the Labiaceae (mint) family. It is nutrient-rich and drought tolerant; however, despite its potential as a food security crop, it has not received the research support needed to promote its widespread adoption and safeguard its diversity. As a result, Frafra potato is becoming extinct and restricted to indigenous communities where it holds traditional importance. It is imperative that conservation methods are explored, but before then, the genetic diversity of the species should be assessed. The objectives of this study were to develop simple sequence repeats (SSRs) for Frafra potato from *de novo* sequencing of the genome, and to provide a preliminary assessment of genetic diversity of Frafra potato in Ghana. To this end, 57 accessions of Frafra potato obtained from three geographic regions of Ghana were characterized using 15 discovered microsatellites. Forty-six multilocus genotypes (MLGs) were identified among the 57 accessions. The genetic diversity among the MLGs was moderate-to-high. Preliminary evaluation of accessions from the three collection zones indicated the presence of population structure among Frafra potato individuals collected in Ghana, Africa.**

**Key words:** Food security, Frafra potato, genetic diversity, microsatellites, simple sequence repeats, *Solenostemon rotundifolius*.

## INTRODUCTION

Frafra potato (*Solenostemon rotundifolius* (Poir.) J.K. Morton, known in northern Ghana as Frafra potato or Persa, in South Africa as Zulu potato, and in India as Chinese potato, is an herbaceous plant that originates

\*Corresponding author. E-mail: rtrigian@utk.edu. Tel: 865.386.1872. Fax: 865.974.4744.

Author(s) agree that this article remain permanently open access under the terms of the [Creative Commons Attribution License 4.0 International License](https://creativecommons.org/licenses/by/4.0/)

from tropical Africa (Aculey et al., 2011). Currently, it is extensively cultivated in Mali, Ghana, Nigeria, and South Africa as well as in tropical Asia. *S. rotundifolius* ( $2n = 64$ ) is a semi-succulent, aromatic herb between 40- 60 cm in height and is a member of the Lamiaceae or the mint family (Tindall, 1983). It produces edible, ovoid tubers up to 8 cm in length. The raw tubers are exceptionally nutritious, and 100 g of tissue contains 394 KJ (94 Kcal) of energy, 1.8 g protein, 0.2 g fat, 21.9 g carbohydrate, 17 mg calcium, 1.1 g fiber, and 6.0 mg iron (Anbuselvi, 2013). The tubers of Frafra potatoes are fermented to make alcoholic beverages (Opoku-Agyeman et al., 2007) or milled for flour, among other uses. However, Frafra potatoes' utility extends beyond that of food. Frafra potato tubers are popular for pharmaceutical and folk medicine applications. For example, the tubers are used to treat dysentery in rural Nigerian communities (Tindall, 1983). In addition to these uses, Frafra potato is also drought-tolerant (Scholten, 2007), probably because it efficiently processes and uses water (Nanema et al., 2009). Based on its superior advantages in nutritional composition, medicinal properties, and the ability to grow in tropical areas, Frafra potato is used as food for bridging the hunger gap (Sugri et al., 2013).

Despite the beneficial characteristics of Frafra potato, research into conservation and breeding of the species has been limited (Opoku-Agyeman et al., 2007; Kwarteng et al., 2018). Hence, it is imperative that the available FRgermplasm is collected and thereafter morphologically and genetically characterized to identify the best accessions for genetic crop improvement. To achieve this goal, specific microsatellite loci for Frafra potato were developed for assessing the genetic diversity among accessions from different regions in Ghana.

Microsatellite loci or simple sequences repeats (SSR) are repeated motifs of 2- to 6 nucleotides that are found throughout the genome and occur mainly in non-coding regions, but are also present in coding segments (Ellegren, 2004). These markers have become relatively routine to discover at a very reasonable cost with the advent of high-throughput and advanced sequencing techniques for genomes. They are very efficient tools for assessing genetic diversity and population structure among the different cultivars and accessions grown by farmers as well as potentially useful accessions growing in the uncultivated environs. The aim of our study was to develop microsatellite loci specific to Frafra potato and preliminarily assess the population structure of the accessions via the genetic data gleaned from SSRs.

## MATERIALS AND METHODS

### Plant materials and DNA extraction

Fifty-seven *S. rotundifolius* accessions were collected from farms in the upper east (24), upper west (18), and the northern regions (15) of Ghana, and grown at the Department of Crop Science, University of Ghana, Accra, Ghana. Genomic DNA was extracted from frozen

leaves of the accessions using a CTAB extraction protocol (Cota-Sánchez et al., 2006) and quantified using a NanoDrop® ND-1000 Spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA).

### Microsatellite loci development

Genomic DNA from a single plant of *S. rotundifolius* '4E016', an upper east accession, was sequenced at the University of Texas, Austin Genomic Sequencing and Analysis facility using an Illumina MiSeq sequencer (Illumina Inc., CA, USA). Paired-end reads were trimmed with the Skewer program version 0.2.2 (Jiang et al., 2014). Sequencing adapters and low quality (mean quality <30), and short reads (<30 bases) were discarded. Read quality control was performed using FastQC (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>). *De novo* assembly was performed using ABySS v1.9.0 (Simpson et al., 2009) with a k-mer size of 64. Sequence filtering for low complexity repeats was completed using the utility DustMasker (Morgulis et al., 2006) on the resulting unitigs. Perfect microsatellite loci were identified using an in-house developed perl script. The minimum and maximum motif frequency defined were 6- to 20 bp di- or tri-nucleotide repeats and 4 to -20 bp tetra-nucleotide repeats. Also, forward and reverse primers flanking each marker were designed using Primer3 (Koressaar and Remm, 2007). Optimum primer size was 21 bp (range 18-27), annealing temperature was 60°C, and a GC content of 40- to -60%.

### Microsatellite loci amplification

The PCR conditions described by Wadl et al. (2011) with some modifications were used to screen and characterize 50 microsatellite loci for *S. rotundifolius*. PCR was performed using 10 µl reaction containing the following in final concentrations: 4 ng genomic DNA, 1×PCR Gold Buffer, 2.5 mM MgCl<sub>2</sub> (Applied Biosystems, Carlsbad, California, USA), 0.25 mM dNTPs, 5% dimethyl sulfide (DMSO; Fisher Scientific, Pittsburgh, Pennsylvania, USA), 0.4 U AmpliTaq Gold Polymerase (Applied Biosystems), 0.25 µM of forward and reverse primer pairs (Integrated DNA Technologies, Coralville, Iowa, USA) and sterile, distilled water. The contents of the reaction mixtures were subjected to the same thermal cycling program found in Wadl et al. (2011). PCR products were separated using the QIAxcel Capillary Electrophoresis System (QIAGEN, Valencia, California, USA) and sized with a 25 to 500 bp size marker and an internal 15/600 alignment marker (Wang et al., 2009).

### Data analyses

QIAxcel ScreenGel (Version 1.5.0) was used to determine the sizes of the PCR products. Flexibin was employed to convert raw allele length data into discrete allelic classes (Amos et al., 2006). The R packages Poppr (Kamvar et al., 2014, 2015) and hierfstat (Goudet, 2005) were utilized for primary data analyses including the number of alleles per locus, observed and expected heterozygosity, evenness, and allelic richness. The population structure across clone corrected dataset ( $n = 46$ ) was evaluated using STRUCTURE v2.3.4 (Pritchard et al., 2000), and Principal Coordinate Analysis (PCoA) with GenAlEx v6.5 (Peakall and Smouse, 2012). Analysis of molecular variance (AMOVA) was conducted with Arlequin v3.5.1.2 (Excoffier and Lischer 2010). Although STRUCTURE and AMOVA analyses were in agreement with PCoA, they were completed with limited accessions of Frafra potato. Large-scale population analyses require larger sample sizes, hence we present preliminary STRUCTURE and AMOVA results.

**Table 1.** Characteristics of 15 microsatellite loci developed for 57 accessions of *S. rotundifolius* (Frafra potato) collected in Ghana, Africa.

GenBank accession number	Locus	Primer sequences (5'-3')	Repeat motif	T <sub>m</sub> (°C)	Number of alleles	Size range (bp)	H <sub>o</sub>	H <sub>e</sub>	Evenness	Allelic richness
MH784481	Sr_001	F: CAGAAGGTGAGGGAGAAGTGG R: GAAAGAGGTAAGCAGTGCAGC	(AT) <sub>8</sub>	60	5	86-117	0.04	0.65	0.85	3.74
MH784482	Sr_009	F: GTGAGTGAGAGACTTATTTGCC R: AGCTTGACTGTACTCGACTCG	(AT) <sub>8</sub>	60	3	227-233	0.11	0.42	0.65	2.85
MH784483	Sr_011	F: TCAACCCACATACTTGGCG R: ACGGATCCTTGCTTGGATTGG	(AG) <sub>15</sub>	61	5	110-141	0.21	0.54	0.66	3.59
MH784484	Sr_017	F: TAAACAGCTAGGCTCCAACCC R: CGTCCCTCCATTTCTGTATTGC	(AG) <sub>8</sub>	60	5	186-208	1.00	0.53	0.85	2.69
MH784485	Sr_018	F: GTTGTGATGGCCGGATTGC R: ACATGAGAGATAGGCAACGAGC	(GA) <sub>7</sub>	60	4	180-196	0.57	0.45	0.74	2.49
MH784486	Sr_020	F: TCTGAACCATTGATTAAGCGG R: TCTTCTCACCTGATTTCTCTCC	(AT) <sub>6</sub>	57	9	132-161	0.64	0.77	0.76	4.43
MH784487	Sr_021	F: CCAACTCTATTCGTCGGCAGC R: AATCCATACCATACGCCAGGG	(AAT) <sub>11</sub>	61	5	194-219	0.40	0.34	0.50	3.13
MH784488	Sr_023	F: GACATTTGGCGCTTTCTCACC R: TCCCAACAGTTTGATCCTCCC	(CTG) <sub>6</sub>	60	4	350-368	1.00	0.68	0.86	3.60
MH784489	Sr_024	F: CTGCTGTAGCTTTCCCTTTGC R: AACTGGGCATGCTACTACTCG	(TGT) <sub>6</sub>	60	5	162-177	1.00	0.65	0.78	3.82
MH784490	Sr_034	F: ATTGGCGGTTATTGGTTGACC R: CCCACCATACACATATATAATCGGC	(TTA) <sub>6</sub>	59	4	250-261	0.20	0.42	0.59	3.20
MH784491	Sr_037	F: AATCCCATCAACAGCCTGAGG R: ATTGCTGGAAGAAATGCTGGC	(TGC) <sub>7</sub>	60	6	157-171	1.00	0.74	0.84	4.66
MH784492	Sr_039	F: ATGAGGCAGAGAGAGGACTCC R: TGCACCACTGTCTTCTTCTCC	(GAT) <sub>6</sub>	60	4	144-154	1.00	0.67	0.86	3.83

Table 1. Contd.

MH784493	Sr_044	F: TCGTG TAGCAATTTCTCACTGC R: GCCAGTGATTAGTTGCACTCG	(TCT) <sub>7</sub>	60	5	241-253	0.02	0.72	0.88	3.78
MH784494	Sr_045	F: CATCGAATCTCAGATCGTGCC R: GTCGCCATCAAGTCTACTGCC	(CGG) <sub>6</sub>	61	5	108-126	1.00	0.70	0.83	4.44
MH784495	Sr_046	F: TTCTCTTCCAATTCCTCCGCC R: TGCAGGGAAGGTTTAGGTTGG	(CTT) <sub>7</sub>	60	6	115-129	0.57	0.67	0.77	4.20

Ho, observed heterozygosity; He, expected heterozygosity.

## RESULTS

One gDNA sample ('4E016') was sequenced. Based on the 3,791,769 sequences (reads), 53,124 microsatellites were discovered. Among them, di-nucleotide repeats accounted for the highest (66.9%), whereas tri- or tetra-nucleotide repeats accounted for 11.5 and 21.5%, respectively. Primer pairs for 15,502 loci were designed based on these microsatellites and the percentages for di-, tri-, and tetra-nucleotide repeats were 70.9, 13.3, and 15.8%, respectively. Fifty primer pairs were screened with six randomly chosen genomic DNA samples, and the fifteen primer pairs that amplified definitive polymorphic alleles without extraneous peaks were employed to analyze the 57 accessions. Clone correction with Poppr identified 11 clones of which five were from the upper east region, one from the upper west region, and five from the northern region. Clone-corrected data ( $n = 46$ ) indicated the presence of seventy-five alleles among 15 polymorphic loci. The number of alleles per locus ranged between 3 (Sr\_009) and 9 (Sr\_020), with an average of 5 alleles per locus (Table 1). Observed heterozygosity of the 46 accessions was between 0.02 and 1.00, whereas the

expected heterozygosity was 0.34 to 0.77. Compliance with the Hardy-Weinberg equilibrium was not evaluated because *S. rotundifolius* is reproduced clonally. Evenness and allelic richness were 0.76 and 3.36 across all loci, respectively, ranging from 0.50 (Sr\_021) to 0.88 (Sr\_044) for evenness and 2.49 (Sr\_018) and 4.66 (Sr\_037) (Table 1). Analysis of microsatellite data from the three collection zones in Ghana with STRUCTURE and principal coordinate analysis (PCoA) indicated the possibility of three different clusters. However, a more extensive collection of accessions is required to confirm these findings (Figure 1).


## DISCUSSION

In this study, the first set of microsatellites from *S. rotundifolius* has been developed, and these loci provide a set of markers to evaluate the genetic diversity of cultivated Frafra potato. The genetic diversity among the sampled Frafra potato accessions was moderate - to- high, and in our sample of 57 accessions, there were only 80% MLGs indicating clonal reproduction among these accessions. Balloux et al. (2003) indicated that asexual or clonally reproducing populations of

plants sustain higher genetic diversity at loci compared to sexually reproducing plants. However, the report also stated that clonal plants typically have fewer genotypes which appear to be true for Frafra potato. Preliminary results of this research suggest the presence of population structure among the accessions from the three regions. However, more accessions must be included in the data set before this finding can be confirmed. It can also be inferred that trade of plants among local farmers contributes to levels of genetic diversity found here. Furthermore, this research data is restricted to the limited number of genotypes collected in three regions of Ghana and does not provide a complete picture of the current status of this indigenous crop that is vulnerable to potential abiotic and biotic stresses.

## Conclusion

The microsatellite markers developed for this study were successfully deployed to assess the genetic diversity of Frafra potato in Ghana. The present study will facilitate further genetic and more complete population studies of this critical African food security crop across a more



**Figure 1.** Principal coordinate analysis (PCoA) of *S. rotundifolius* (Frafra potato) accessions from Ghana, Africa. Analysis of clone corrected data revealed three groups among the collection zones. UW = Upper West; UE = Upper East and NR = Northern Region of Ghana.

extensive geographical area.

## CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

## ACKNOWLEDGEMENTS

This research was sponsored by The University of Tennessee, Nanjing Agricultural University, and the University of Ghana.

## REFERENCES

- Aculey K, Quainoo A, Mahanu G (2011). Feasibility studies on the potential of grafting and budding of frafra potato (*Solenostemon rotundifolius*). *Journal of Horticulture and Forestry* 3:327-332.
- Amos W, Hoffman J, Frodsham A, Zhang A, Best S, Hill AVS (2006). Automated binning of microsatellite alleles: problems and solutions. *Molecular Ecology Notes* 7:10-14.
- Anbuselvi S, Priya MH (2013). Nutritional and anti-nutritional constituent of *Plectranthus rotundifolius*. *International Journal Pharmaceutical Science Review and Research* 22:213-215.
- Balloux F, Lehmann L, de Meeus T (2003). The population genetics of clonal and partially clonal diploids. *Genetics* 164(4):1635-1644.
- Cota- Sánchez JH, Remarchuk K, Ubayasena K (2006). Ready-to-use DNA extracted with a CTAB method adapted for herbarium specimens and mucilaginous plant tissue. *Plant Molecular Biology Reporter* 24(2):161-167.
- Ellegren H (2004). Microsatellites: simple sequences with complex evolution. *Nature Review Genetics* 5:435-445. DOI:10.1038/nrg1348
- Excoffier L, Lischer HEL (2010). Arlequin suite ver 3.5: A new series of programs to perform population genetics analyses under Linux and Windows. *Molecular Ecology Resources* 10:564-567.
- Goudet J (2005). Hierfstat, a package for R to compute and test variance components and F-statistics. *Molecular Ecology Notes* 5:184-186.
- Jiang H, Lei R, Ding SW, Zhu S (2014). Skewer: a fast and accurate adapter trimmer for next-generation sequencing paired-end reads. *BMC Bioinformatics* 15:182.
- Kamvar ZN, Brooks JC, Grunwald NJ (2015). Novel R tools for analysis of genome-wide population genetic data with emphasis on clonality. *Frontiers of Genetics* 6:208. DOI: doi: 10.3389/fgene.2015.00208
- Kamvar ZN, Tabima JF, Grunwald NJ (2014). Poppr: an R package for genetic analysis of populations with clonal, partially clonal, and/or sexual reproduction. *Peer J*. 2:e281.
- Koressaar T, Remm M (2007). Enhancements and modifications of primer design program Primer3. *Bioinformatics* 23:1289-1291.
- Kwarteng AO, Ghunney T, Adu Amoah R, Nyadanu D, Abogoom J, Nyam KC, Ziyaaba JZ, Danso EO, Whyte T, Asiedu DD (2018). Current knowledge and breeding avenues to improve upon Frafra potato (*Solenostemon rotundifolius* (Poir.) JK Morton). *Genetic Resources and Crop Evolution* 65:659-669.
- Morgulis A, Gertz EM, Schäffer AA, Agarwala R (2006). A fast and symmetric DUST implementation to mask low-complexity DNA sequences. *Journal of Computational Biology* 13:1028-1040.
- Nanema RK, Traore ER, Bationo-Kando P, Zongo JD (2009). Morphoagronomical characterization of *Solenostemon rotundifolius* (Poir. J.K. Morton) (Lamiaceae) germplasm from Burkina Faso. *International Journal of Biological and Chemical Sciences* 3:1100-1113.
- Opoku-Agyeman MO, Bennett-Lartey SO, Vodouhe RS, Osei C, Quarcoo E, Boateng SK, Osekere EA (2007). Morphological characterization of frafra potato (*Solenostemon rotundifolius*) germplasm from the savannah regions of Ghana. In: Vodouhe R, Atta-Krah K, Achigan-Dako GE, Eyog-Matig O, Avohou H (eds) *Plant genetic resources and food security in West and Central Africa. Regional Conference, Ibadan, Nigeria* pp. 26-30.

- Peakall R, Smouse PE (2012). GenAEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research -- an update. *Bioinformatics* 28:2537-2539.
- Pritchard JK, Stephens M, Donnelly P (2000). Inference of population structure using multilocus genotype data. *Genetics* 155:945-959
- Scholten W (2007). Agricultural development and water use in the Central Rift Valley of Ethiopia: A rapid appraisal. Essay (Bachelor) of University of Twente. <https://essay.utwente.nl/68647/1/Scholten-Wolter.pdf>
- Simpson JT, Wong K, Jackman SD, Schein JE, Jones SJ, Birol I (2009). ABySS: a parallel assembler for short read sequence data. *Genome Research* 19:1117-1123.
- Sugri I, Kusi F, Kanton RAL, Nutsugah SK, Zakaria M (2013). Sustaining Frafra potato (*Solenostemon rotundifolius* Poir.) in the food chain; current opportunities in Ghana. *Journal of Plant Science* 1:68-75.
- Tindall HD (1983). *Vegetables in the tropics*. ELBS. Macmillan Education Ltd., Houndmills, Basingstoke, Hampshire RG 21(2):5.
- Wadi PA, Dattilo AJ, Scheffler BE, Trigiano RN (2011). Development of microsatellite loci for the endangered species *Pityopsis ruthii* (Asteraceae). *American Journal of Botany* 98:e342-5.
- Wang, X, Rinehart TA, Wadi PA, Windham MT, Spiers JM, Johnson DH, Trigiano RN (2009). A new electrophoresis technique to separate microsatellite alleles. *African Journal of Biotechnology* 8(11):2432-2436.



**Related Journals:**

