

Full Length Research Paper

Species-specific detection of *Dickeya* sp. (*Pectobacterium chrysanthemi*) in infected banana tissues, soil and water

Peiqian Li, Birun Lin* Huifang Shen and Xiaoming Pu

Plant Protection Research Institute, Guangdong Academy of Agricultural Sciences, Guangzhou 510640, PR China.

Accepted 21 October, 2011

***Dickeya* sp. (*Pectobacterium chrysanthemi*) is a bacterial pathogen that can cause soft rot disease on banana pseudostem tissue. A species-specific polymerase chain reaction (PCR) assay was developed for rapid and sensitive detection of the pathogenic bacteria in diseased plant tissues, soil and artificially infested irrigation water. Based on differences in internal transcribed spacer (ITS) sequences of *Dickeya* sp. (*P. chrysanthemi*) and other bacteria, a pair of species-specific primers was synthesized. Specificity was tested against eight bacterial organisms associated with banana. The LF/LR primers amplified only a single PCR band of 171 bp from *Dickeya* sp. (*P. chrysanthemi*). The detection sensitivity was determined to be 0.44 fg for pure genomic DNA per 25 µl reaction volume. The results suggest that the assay detected the pathogen more rapidly and accurately than standard isolation methods. The PCR-based methods developed here could simplify both plant disease diagnosis and pathogen monitoring, as well as guide plant disease management.**

Key words: Soft rot, banana, detection, polymerase chain reaction (PCR).

INTRODUCTION

Dickeya sp. (*Pectobacterium chrysanthemi*) is a plant-pathogenic enterobacterium responsible for soft rot disease in a wide range of species (Lee and Yu, 2006). The gram-negative bacterium, important pathogen, first identified on banana in 2009 at Panyu, Guangzhou, China (Lin et al., 2010), was accountable for significantly economic losses to banana production. This pathogen is survives in stagnant irrigation water and soil, often in association with decaying crop residues. Usually contamination or infection is a subclinical level; such latent infections cannot be detected by visual inspection.

Polymerase chain reaction (PCR) techniques offer advantages over traditional methods of detection and diagnosis. The practice of diagnosing plant pathogens using

PCR has previously been described (Henson and French, 1993; Lévesque, 2001; McCartney et al., 2003; Atkins and Clark, 2004). Plant diseases can be controlled most effectively if diagnostics is introduced at an early stage of disease development. Information resulting from molecular diagnostics could be used to make more rational decisions about the choice and use of proper agrochemicals at optimal application times. Accurate and early diagnosis of a specific pathogen is greatly important in a crop protection system or could be useful in selection within the breeding programs. Standard diagnostic methods, such as visual assessment of symptoms, as well as microscopic or cultural methods, often have limited potential. External symptoms can be masked, especially if pathogens occur simultaneously. The microscopic method evaluation mostly requires specialized training, experience and the taxonomical knowledge of an investigator and cultural techniques can also be time-consuming (Matusinsky et al., 2010). Following the appearance of the PCR technology, because of sensitivity, accuracy and relatively rapid results, molecular detection method has been widely used for pathogen diagnoses.

*Corresponding author. E-mail: linbrgdpp@163.com. Fax: +86 020 85161048.

Abbreviations: PCR, Polymerase chain reaction; ITS, internal transcribed spacer.



Figure 1. Symptoms appearing on healthy banana leaves after inoculation treatments. **A**, Leaves injected with *Dickeya* sp. (*Pectobacterium chrysanthemi*) bacterial suspension; **B**, leaves injected with distilled water, designed as control.

In this study, PCR primers derived from 16S-23S rDNA internal transcribed spacer (ITS) sequences were designed for the specific detection of *Dickeya* sp. (*P. chrysanthemi*) (Gurtler and Stanisich, 1996; Kwon et al., 2000). The specificity and sensitivity of the reaction were tested and the PCR protocols were used to detect diseased plant tissues, irrigation water, and infested soil samples collected in the field.

MATERIALS AND METHODS

Isolates

In 2009, *Dickeya* sp. (*P. chrysanthemi*) strains were isolated from naturally diseased banana pseudostem tissue at the growth stage showing soft rot symptoms in various cultivation areas of Guangdong province, China. Individual colonies isolated were further assayed by injecting 20 μ l 10⁹CFU/ml bacterial suspension into the healthy banana leaves to test pathogenicity (Lee and Yu, 2006) (Figure 1).

Bacterial strains and DNA extraction

The bacterial strains used in this study are listed in Table 1. The reference strains were grown in 30 ml of nutrient broth for 24 to 48 h on an orbital shaker at 35°C. Cultures were harvested in late log phase and genomic DNA was extracted using a commercial DNA purification kit (QIAGEN, France). DNA of infected and non-infected banana pseudostem tissue was extracted according to Tooley et al. (1997) using 1 g of tissue samples that were previously ground in liquid nitrogen. Soil samples DNA was extracted directly according to the method of Li (Li and Hartman, 2003). A TENP extraction buffer (containing 50 mM Tris, pH 8.0; 20 mM ethylene diamine tetraacetic (EDTA) disodium salt pH 8.0, 100 mM NaCl and 1% polyvinylpyrrolidone) was added to the lyophilized soil suspended in 0.5 ml 0.4% dry milk powder solution. After vortexing, samples were incubated at 65°C for 30 min, and then 10 ml of 20% (w/v) sodium dodecyl sulfate (SDS) was added. The samples were incubated at 65°C for another 30 min, then samples were centrifuged at 18 000 *g* for 10 min. The supernatant was transferred

to a new centrifuge tube, and half volume of 7.5 M ammonium acetate and two volumes of 100% ethanol at -20°C for 2 h were added. DNA was pelleted, rinsed with 70% ethanol, dried, dissolved in sterile water and stored at -20°C. DNA yield and purity were determined spectrophotometrically.

PCR amplification conditions

PCRs were carried out in a GeneAmp PCR systems 9700 thermocycler (Applied Biosystems, Courtaboeuf, France) in 25 μ l containing 1 \times Mg²⁺-free PCR buffer (Gibco-BRL, Burlington, Ontario, Canada), 2.0 mM MgCl₂, 0.16 mM dNTPs each, 0.4 μ M primer each, 2.5 units of Taq DNA polymerase (Gibco/BRL), and 1 μ l of genomic DNA (10-15 ng/ μ l) with the following thermal profile: 94°C for 5 min followed by 35 cycles of 94°C for 30 s, 53°C for 20 s, and 72°C for 30 s, and a final extension at 72°C for 10 min.

16S-23S rDNA gene ITS PCR amplification

The *Dickeya* sp. (*P. chrysanthemi*) 16S-23S rDNA ITS region was amplified by PCR. The PCR product with the expected size of approximately 440 bp was obtained with universal primers (primer1: 5'-GAAGTCGTAACAAGG-3'; primer2: 5'-CAAGGCATCCACCGT-3') for bacterial ITS amplification described earlier by Jensen et al. (1993). The purified PCR products were cloned into the plasmid vector pMD20-T (TaKaRa), and positive clones were sequenced in Invitrogen Company. The sequence was submitted to GenBank.

Specific primers design

Specific primers for *Dickeya* sp. (*P. chrysanthemi*) were designed by the comparison of the ITS of 18 different *Erwinia* sequences from GenBank (Table 2). The pair of primers were designed using Primer3Plus software (Untergasser et al., 2007), and they were analyzed using the Net-Primer (Biosoft International) software (NetPrimer) which identifies regions of likely secondary structure (hairpins) and potential primer dimer formation and assigns values for melting temperature (T_m). The specificities of the primers were checked using FASTA and basic local alignment search tool (BLAST) searches of the GenBank databases.

Table 1. Bacterial isolates used to screen the specificity of the primers.

Species	Host	Origin	Amplification with primers LF/LR
<i>Dickeya</i> sp. (<i>Pectobacterium chrysanthemi</i>)	Banana	Guangdong	+
<i>Pectobacterium chrysanthemi</i>	Unknown	ATCC11663	+
<i>Erwinia amylovora</i>	Unknown	Zhejiang	-
<i>Erwinia carotovora</i> subsp. <i>carotovora</i>	Carrot	Yunnan	-
<i>Xanthomonas campestris</i> pv. <i>musacearum</i>	Banana	CGMCC	-
<i>Ralstonia solanacearum</i> race 2	Banana	CGMCC	-
<i>Pantoea stewartii</i> subsp. <i>Stewartii</i>	Corn	Shandong	-
<i>Xanthomonas oryzae</i> pv. <i>oryzae</i>	Rice	Guangdong	-
<i>Curtobacterium flaccumfaciens</i> pv. <i>flaccumfaciens</i>	Unknown	CGMCC	-
<i>Pseudomonas syringae</i> pv. <i>syringae</i>	Unknown	CGMCC	-

+, 171bp product amplified by primers LF/LR; -, no amplified products; CGMCC, China general microbiological culture collection; ATCC, American type culture collection.

Table 2. GeneBank accession numbers and sequences compared to develop species-specific primers LF/LR for *Dickeya* sp. (*Pectobacterium chrysanthemi*).

Accession number	Species	Accession number	Species
GQ181062	<i>Erwinia amylovora</i>	EF427910.1	<i>Erwinia pyrifoliae</i>
HM130661.1	<i>Pectobacterium carotovorum</i>	EF427911.1	<i>Erwinia pyrifoliae</i>
HM130662.1	<i>Pectobacterium carotovorum</i>	EF427912.1	<i>Erwinia pyrifoliae</i>
HM130663.1	<i>Pectobacterium carotovorum</i>	EF427913.1	<i>Erwinia pyrifoliae</i>
HM130664.1	<i>Pectobacterium carotovorum</i>	EF427914.1	<i>Erwinia pyrifoliae</i>
EF427906.1	<i>Erwinia pyrifoliae</i>	EF422407.1	<i>Erwinia pyrifoliae</i>
EF427907.1	<i>Erwinia pyrifoliae</i>	AF232681.1	<i>Erwinia chrysanthemi</i>
EF427908.1	<i>Erwinia pyrifoliae</i>	AF232682.1	<i>Erwinia chrysanthemi</i>
EF427909.1	<i>Erwinia pyrifoliae</i>	AF232683.1	<i>Erwinia chrysanthemi</i>

Test of primers specificity and sensitivity

The designed primer pairs were tested for their specificity against the DNA of *Dickeya* sp. (*P. chrysanthemi*) isolated on diseased banana pseudostem tissue and other bacterial pathogens associated with disease of crops. Furthermore, the potential for detection was tested in banana pseudostem tissue infected with *Dickeya* sp. (*P. chrysanthemi*) compared with non-infected ones. The sensitivity of diagnostic PCR was determined for *Dickeya* sp. (*P. chrysanthemi*) by testing a 10-fold dilution series of known DNA concentration. DNA extracts were initially tested at concentrations on the order of 44 ng/μl; the concentration was decreased until no products could be observed by agarose gels. All reactions were repeated at least three times.

RESULTS

16S-23S rDNA gene ITS sequence analysis

16S–23S rDNA coding spacer regions of the *Dickeya* sp.

(*P. chrysanthemi*) were amplified using universal primers (primer1/2) for bacterial ITS amplification (Jensen et al., 1993) and products were cloned and sequenced (Genbank no:HQ882841). The 440 bp length sequence was identical with sequence from *Dickeya* sp. (*P. chrysanthemi*) strain ATCC 11663 (Type strain, GenBank Accession no. AF232681.1) submitted previously, which further enhance authenticity of our study.

Specificity and sensitivity of PCR amplification

Specific primers for *Dickeya* sp. (*P. chrysanthemi*) (LF 5'-TTCGTCTAGAGGCCAGGAC-3' and LR 5'-TCAGCTTGTTCCGGATTGTT-3') were designed. In the essential specificity test, the primer set LF/LR was able to amplify a unique DNA fragment of approximately 171 bp (Figure 2) from all *Dickeya* sp. (*P. chrysanthemi*) isolates tested from naturally diseased banana pseudostem tissue.




Figure 2. Agarose gel electrophoresis of PCR-amplified products using the specific primers LF/LR. Lane 1, 2000bp DNA marker; lane 2, *P. chrysanthemi* (ATCC11663); lanes 3 to 4, *Dickeya* sp. (*Pectobacterium chrysanthemi*) isolates; lane 5, infected banana tissues; lane 6, *Erwinia amylovora*; lane 7, *Erwinia carotovora* subsp. *carotovora*; lane 8, *Xanthomonas campestris* pv. *musacearum*; lane 9, *Ralstonia solanacearum* race 2; lane 10, *Pantoea stewartii* subsp. *Stewartii*; lane 11, *Xanthomonas oryzae* pv. *oryzae*; lane 12, *Curtobacterium flaccumfaciens* pv. *flaccumfaciens*; lane 13; *Pseudomonas syringae* pv. *syringae*; lanes 14, non-infected banana tissues; lane 15, ddH₂O.

However, the other reference strains tested showed no amplification product. The primer set was therefore used for further analysis. The primers LF/LR were used to examine infected and non-infected plant materials and one sharp band (171 bp) was amplified in infected banana tissues and non-infected not.

The sensitivity of diagnostic PCR was assessed by determining the minimum amount of *Dickeya* sp. (*P. chrysanthemi*) DNA. Amplification reactions were conducted with 10-fold decreasing amounts of genomic DNA of *Dickeya* sp. (*P. chrysanthemi*). In a 25 μ l reaction volume assay, primers LF/LR were able to detect 0.44 fg of pure genomic DNA (Figure 3A).

Detection in plant tissues

The PCR assay with the pair of species-specific primers LF/LR detected *Dickeya* sp. (*P. chrysanthemi*) in naturally infected plant tissues from Hainan, Guangdong, Guangxi, and Fujian. This PCR method succeeded in amplifying the expected 171 bp product using DNA samples extracted from all soft-rot infected banana samples (Figure 3B). In contrast, no PCR product was amplified from healthy banana tissues. This indicates that the DNA amplified in PCR assays by primer sets LF/LR was derived from *Dickeya* sp. (*P. chrysanthemi*).

Detection in infested field soil and artificially infested irrigation water

DNA extracted from field soil samples collected from 12 infested banana fields in Guangdong and twice-autoclaved field soil sample (healthy banana fields as negative control) was subjected to PCR using primers LF/LR. The DNA samples extracted from 1 g soil were suspended in 20 μ l ultrapure water used for PCR amplification. 10 of the 12 samples provided 171 bp PCR products amplified by LF/LR (Figure 3C), while PCR with the pair of primers yielded the 171 bp band in irrigation water artificially infested with *Dickeya* sp. (*P. chrysanthemi*) 10⁹ CFU/ml bacterial suspension. The detection limit of PCR template was 10³ CFU/ml in 25 μ l volumes reaction (Figure 3D), indicating that the method used in this study can sensitively detect the pathogen in water.

DISCUSSION

Molecular technologies have become some of the most important tools for identifying and detecting plant pathogens (Matusinsky et al., 2010). The rapid and reliable detection of pathogens is important for disease management, plant breeding, and epidemiological studies. In this study, our objective was to develop a rapid, sensitive, and effective




Figure 3. A, Sensitivity of PCR for detection of *Dickeya* sp. (*Pectobacterium chrysanthemi*). Lane 1, 2000bp DNA marker; lanes 2 to 9, amplified products using DNA at concentrations of 440 pg, 44 pg, 4.4 pg, 440 fg, 44 fg, 4.4 fg, 440 ag and 44 ag in a 25 μ l PCR reaction; lane 10, ddH₂O. The same results were obtained in 3 replicates; B, PCR amplification of DNA extracted from diseased plants; lane 1, 2000bp DNA marker; lane 2, DNA from *Dickeya* sp. (*Pectobacterium chrysanthemi*); lanes 3 to12, amplified products using DNA from diseased banana tissues; lane 13, DNA from healthy banana tissues; C, products amplified from DNA extracted from soil samples; lane 1, 2000bp DNA ladder marker; lane 2, DNA from *Dickeya* sp. (*Pectobacterium chrysanthemi*); lanes 3 to 14, amplified products using DNA from infested soil samples; lane 15, negative control; D, products amplified from different concentrations of irrigation water infested with *Dickeya* sp. (*P. chrysanthemi*) bacterial suspension; lane 1, 2000bp marker; lane 2, DNA from *Dickeya* sp. (*Pectobacterium chrysanthemi*); lanes 3-11, Amplified products using infested irrigation water at concentrations of 10⁹ CFU/ml, 10⁸ CFU/ml, 10⁷ CFU/ml, 10⁶ CFU/ml, 10⁵ CFU/ml, 10⁴ CFU/ml, 10³ CFU/ml, 10² CFU/ml, 10 CFU/ml; lane 12, ddH₂O.

molecular method using species-specific PCR to identify and detect *Dickeya* sp. (*P. chrysanthemi*) which cause an odorous soft rot of the center of banana rhizome (Lin et al., 2010).

PCR-based assays applied to detecting plant pathogens have been described (Henson and French, 1993; Tooley et al., 1997; Cullen et al., 2002; Mercado-Blanco et al., 2003) as well as to

monitor microorganisms in soils (Cullen et al., 2001; Lees et al., 2002; Carnegie et al., 2003). Most of the primers were designed for the rDNA region, and particularly the ITS region (Nazar et al., 1991; Ma and Michailides, 2002; Zhao et al., 2007) or other conserved genes, for example, β -tubulin (Hirsch et al., 2000), elongation factors (Filion et al., 2003; Li and Hartman, 2003), mito-

chondrial DNA (mtDNA) (Frederick et al., 2000; Tooley et al., 2006) and so on. The same to the previous study (Nazar et al., 1991; Ma and Michailides, 2002; Zhao et al., 2007), we designed specific primers from *Dickeya* sp. (*P. chrysanthemi*) 16S-23S rDNA ITS sequence.

The rDNA genes are essential for the survival of all organisms and are highly conserved in the

bacterial and other kingdoms. In addition, sequence information of 16S-23S spacer region suggests that considerable variation can occur between species in both the length and the sequence of this region. Therefore spacer variations between strains, species and genera may be used for identification and typing purposes (Gurtler and Stanisich, 1996). In our research, 16S–23S rDNA coding spacer regions of *Dickeya* sp. (*P. chrysanthemi*) were cloned and sequenced, and specific primers (LF/LR) were designed to amplify PCR products of 171 bp.

Some diseases have similar symptoms in early stages and infection is a subclinical level. Besides, agricultural field soil is a complex ecosystem with a diverse microbial community (Torsvik and Øvreås, 2002). There are many hundreds of different species of microorganism including fungi, bacteria and nematodes that are found in field soil. Thus, it is difficult and time-consuming to detect the presence and distinguish *Dickeya* sp. (*P. chrysanthemi*) from the other microbes in plant tissues and soil with standard diagnostic methods. However, the PCR detection method reported here can provide a definitive diagnosis of *Dickeya* sp. (*P. chrysanthemi*) in infected banana tissues, soils and water within hours. Moreover, this method is very easy to use and requires minimal training. The specific primer set LF/LR could not only distinguish *Dickeya* sp. (*P. chrysanthemi*) from other pathogens but could also accurately diagnose and survey the pathogen from diverse areas of China. The primers were sensitive enough to accurately amplify as little as 0.44 fg of diluted pure genomic DNA and 10³ CFU/ml irrigation water artificially inoculated with *Dickeya* sp. (*P. chrysanthemi*) suspension in 25 µl reaction volume assay (Figures 3A and 3D). This level of detection limit may be of particular importance where it is desirable to detect the pathogen before disease symptoms have developed and most of plants are infected.

This PCR-based detection can be used together with standard diagnostic methods focused on plant pathology. If needed, results of analyses will be available in a few hours. Thus, farmers can get early information about the disease pathogen, which can contribute to good decision-making about a strategy of disease management. To summarize, results from this study demonstrate the usefulness of the PCR-based molecular detection of *Dickeya* sp. (*P. chrysanthemi*) in banana production.

ACKNOWLEDGEMENTS

This research was supported by National natural science foundation of China (NO.30971958) and Special agro-scientific research in the public interest program (201003004). We thank Dr B.Z. Feng for the assistance with the manuscript revision and English improvement.

REFERENCES

Atkins SD, Clark IM (2004). Fungal molecular diagnostics: a mini review. *J. Appl. Genet.* 45: 3-15.

- Carnegie SF, Choiseul JW, Roberts AMI (2003). Detection of *Colletotrichum coccodes* and *Helminthosporium solani* in soils by bioassay. *Plant Pathol.* 52: 13–21.
- Cullen DW, Lees AK, Toth IK, Duncan JM (2001). Conventional PCR and real-time quantitative PCR detection of *Helminthosporium solani* in soil and on potato tubers. *Eur. J. Plant Pathol.* 107: 387–398.
- Cullen DW, Lees AK, Toth IK, Duncan JM (2002). Detection of *Colletotrichum coccodes* from soil and potato tubers by conventional and quantitative real-time PCR. *Plant Pathol.* 51: 281-292.
- Filion M, St-Arnaud M, Jabaji-Hare SH (2003). Direct quantification of fungal DNA from soil substrate using real-time PCR. *J. Microbiol. Methods* 53: 67-76.
- Frederick RD, Snyder KE, Tooley PW, Berthier-Shaad Y, Peterson GL, Bonde MR, Shaad NW, Knorr DA (2000). Identification and differentiation of *Tilletia indica* and *T. walkeri* using the polymerase chain reaction. *Phytopathology*, 90: 951-960.
- Gurtler V, Stanisich VA (1996). New approaches to typing and identification of bacteria using the 16s-23s rDNA spacer region. *Microbiology*, 142: 3-16.
- Henson JM, French R (1993). The polymerase chain reaction and plant disease diagnostics. *Annu. Rev. Phytopathol.* 31: 81-109.
- Hirsch PR, Mauchline TH, Mendum TA, Kerry BR (2000). Detection of the nematophagous fungus *Verticillium chlamydosporium* in nematode-infested plant roots using PCR. *Mycol. Res.* 104: 435-439.
- Jensen MA, Webster JA, Straus N (1993). Rapid identification of bacteria on the basis of polymerase chain reaction-amplified ribosomal DNA spacer polymorphisms. *Appl. Environ. Microbiol.* 59: 945-952.
- Kwon SW, Myung IS, Go SJ (2000). Detection of *Pectobacterium chrysanthemi* using specific PCR primers designed from the 16S-23S rDNA intergenic spacer region. *Plant Pathol. J.* 16(5): 252-256.
- Lee YA, Yu CP (2006). A differential medium for the isolation and rapid identification of a plant soft rot pathogen, *Erwinia chrysanthemi*. *J. Microbiol. Methods.* 64: 200-206.
- Lees AK, Cullen DW, Sullivan L, Nicolson MJ (2002). Development of conventional and quantitative real-time PCR assays for the detection and identification of *Rhizoctonia solani* AG-3 in potato and soil. *Plant Pathol.* 51: 293-302.
- Lévesque AC (2001). Molecular methods for detection of plant pathogens e what is the future? *Can. J. Plant Pathol.* 24: 333-336.
- Li S, Hartman GL (2003). Molecular detection of *Fusariumsolani* f.sp. *glycines* in soybean roots and soil. *Plant Pathol.* 52: 74-83.
- Lin BR, Shen HF, Pu XM, Tian XS (2010). First Report of a soft rot of banana in mainland China caused by a *Dickeya* sp. (*Pectobacterium chrysanthemi*) *Plant Dis.* 94: 640.
- Ma Z, Michailides TJ (2002). A PCR-based technique for identification of *Fusicoccum* sp. from pistachio and various other hosts in California. *Plant Dis.* 86: 515-520.
- Matusinsky P, Frei P, Mikolasova R, Svacinova I, Tvaruzek L, Spitzer T (2010). Species-specific detection of *Bipolaris sorokiniana* from wheat and barley tissues. *Crop Prot.* 29: 1325-1330.
- McCartney HA, Foster SJ, Fraaije BA, Ward E (2003). Molecular diagnostics for fungal plant pathogens. *Pest Manage. Sci.* 59: 129-142.
- Mercado-Blanco J, Collado-Romero M, Parrilla-Araujo S, Rodriguez-Jurado D, Jimnez-Daz RM (2003). Quantitative monitoring of colonization of olive genotypes by *Verticillium dahliae* pathotypes with real-time polymerase chain reaction. *Physiol. Mol. Plant Pathol.* 63: 91-105.
- Nazar RN, Hu X, Schmidt J, Culham D, Robb J (1991). Potential use of PCR amplified ribosomal intergenic sequences in the detection and differentiation of *Verticillium* wilt pathogens. *Physiol. Mol. Plant Pathol.* 39: 1-11.
- Tooley PW, Bunyard BA, Carras MM, Hatziloukas E (1997). Development of PCR primers from internal transcribed spacer region 2 for detection of *Phytophthora* species infecting potatoes. *Appl. Environ. Microbiol.* 63: 1467-1475.
- Tooley PW, Martin FN, Carras MM, Frederick RD (2006). Real-time fluorescent polymerase chain reaction detection of *Phytophthora ramorum* and *Phytophthora pseudosyringae* using mitochondrial gene regions. *Phytopathology*, 96: 336-345.

Torsvik V, Øvreås L (2002). Microbial diversity and function in soil: from genes to ecosystems. *Curr. Opin. Microbiol.* 5: 240–245.

Untergasser A, Nijveen H, Rao X, Bisseling T, Geurts R, Leunissen JAM (2007). Primer3Plus, an enhanced web interface to Primer3. *Nucleic Acids Res.* 35: 71-74.

Zhao J, Wang XJ, Chen CQ, Huang LL, Kang ZS (2007). A PCR-based assay for detection of *Puccinia striiformis* f. sp. *tritici* in wheat. *Plant Dis.* 91: 1669-1674.