

Full Length Research Paper

# Co-transfer of *gfp*, CHS and *hptII* genes into *Oncidium* Sharry Baby PLB using the biolistic gun

Ng Chea Yee<sup>1</sup>, Janna Ong Abdullah<sup>1\*</sup>, Maziah Mahmood<sup>2</sup>, Nazir Basiron<sup>3</sup>

<sup>1</sup>Department of Microbiology, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, 43400 UPM, Serdang, Selangor Darul Ehsan, Malaysia.

<sup>2</sup>Department of Biochemistry, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, 43400 UPM, Serdang, Selangor Darul Ehsan, Malaysia.

<sup>3</sup>Malaysian Institute for Nuclear Technology Research (MINT), Bangi, 43000, Kajang, Selangor Darul Ehsan, Malaysia.

Accepted 25 June, 2008

This study was aimed at assessing the biolistic method of co-transforming non-linked genes on separate gene cassettes into *Oncidium* Sharry Baby's protocorm-like-body (PLB). An expression vector containing a synthetic green fluorescence protein (*sgfp*) gene driven by the *Cauliflower Mosaic Virus* (*CaMV*) 35S promoter and another vector containing the antisense chalcone synthase (CHS) and hygromycin resistant (*hpt II*) genes, were successfully co-bombarded into *Oncidium* Sharry Baby PLB. Six critical parameters (PLB size, time course of *gfp* transient expression in PLB, DNA concentration, PLB age, presence of spermidine and CaCl<sub>2</sub> in the DNA-microcarrier precipitation, and duration of PLB on fresh medium prior bombardment) were optimized based on transient *gfp* expression. One month after bombardment, the PLB were subjected to 5 µg/ml hygromycin selection for 8 months. A total of 137 regenerated putative transformants were randomly selected and verified by polymerase chain reaction (PCR) analysis. The results indicated the presence of the transgenes *gfp*, *hptII* and antisense CHS in 28, 61, and 11% of the selected putative transformants, respectively.

**Key words:** Biolistic gun, co-transformation, orchidaceae, protocorm-like-body.

## INTRODUCTION

The family *Orchidaceae* includes a number of commercially important hybrids that are grown for cut flowers and potted plants. Cut-flower production and potted plant cultivation have been mainstays for agro-industries worldwide. *Oncidium* sp. is a popular ornamental plant for worldwide market, and is produced as a high value cash crop plant for cut-flower in Malaysia. The fragrant types provide additional commercial values. However, some of these fragrant orchids possess dull flower colours and are not suitable for use as cut flowers. On top of that, the

orchid industries in Malaysia are facing great competition from neighbouring exporting countries. Therefore, there is substantial interest in the improvement and production of these commercially valuable plants to give Malaysia a better competitive edge in worldwide market. Unfortunately orchids usually have long juvenile periods and reproductive cycle (several years), which restrict genetic improvement using the traditional sexual hybridization method. Therefore, the application of modern genetic engineering techniques provides an alternative for orchid improvement of flower colour and morphology in order to increase the commercial value of the orchid.

Until recently, transgenic orchid plants have been reported for only a few orchid genera. The first transgenic orchid plant was reported for *Dendrobium* which was generated using biolistic bombardment. The transformation was verified using kanamycin selection and polymerase chain reaction analysis (Kuehnle and Sugii, 1992). Chia et al. (1994) also transformed *Dendrobium*

\*Corresponding author: E-mail: [janna@biotech.upm.edu.my](mailto:janna@biotech.upm.edu.my).  
Tel: 603-89466697. Fax: 603-89430913.

**Abbreviations:** *CaMV*, *Cauliflower Mosaic Virus*; **CHS**, chalcone synthase; *hpt II*, hygromycin phosphotransferase type II; *sgfp*, synthetic green fluorescence protein; **PLB**, protocorm-like-body; **PCR**, polymerase chain reaction.

through biolistic bombardment and used a non-invasive selection system while Tee and Marziah (2005) used the GFP and GUS reporter systems for their *Dendrobium* callus transformation. Yu et al. (2001) transformed *Dendrobium* by inoculating thin sections of the PLB with *Agrobacterium tumefaciens*. Transformation of other orchid genera includes *Cymbidium* (Yang et al., 1999), *Phalaenopsis* (Anzai et al., 1996; Belarmino and Mii, 2000) and *Vanilla* (Malabadi and Nataraja, 2007). The various orchid genera each have its own optimal transformation conditions. Therefore, an effective transformation system for gene transfer into *Oncidium* PLB needs to be established for producing new varieties of *Oncidium* orchid.

Most available useful genes are usually not cloned conveniently in the same vector with the preferred reporter or selectable marker gene. Hence, an alternative to reconstructing a new transforming vector harbouring all the desired genes would be to co-transform available constructs into the target tissues (Ramesh and Gupta, 2005; Yao et al., 2007). Besides that, to avoid unwanted DNA in the transgenic plant, it is also preferred to have the genes on separate gene cassettes (Zhao et al., 2007). In either case, co-transformation theoretically provides equal chances for both genes on different vectors to enter the cell. The present work was carried out with the aim of assessing the biolistic system for the introduction of common marker genes (*gfp* and *hpt*) and a flower colour gene (CHS) on different vectors into *Oncidium* Sharry Baby PLB. PLB are the choice target tissues because they are the most easily obtained materials for most orchids and they have high capability of regenerating into new plants. Five critical parameters (size of PLB, time course of *gfp* transient expression in PLB, concentration of DNA, age of PLB, spermidine and  $\text{CaCl}_2$  in DNA-microcarrier precipitation and duration of PLB in fresh medium prior bombardment) were evaluated to provide useful information for future transformation work on biolistic introduction of specific beneficial genes into PLB of *Oncidium* Sharry Baby.

## MATERIALS AND METHODS

### Plasmids

The plasmid pSM-CHS (a generous gift from Dr. Abdullah Sipat, Department of Biochemistry, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia) contains an antisense chalcone synthase (CHS) gene and a hygromycin phosphotransferase (*hpt* II) gene driven by the *Cauliflower Mosaic Virus (CaMV)* 35S promoter that confers hygromycin resistance. p35S is approximately 4 kb, was obtained from Sheen et al. (1995). It carries a synthetic (s) *gfp* gene driven by the *CaMV* 35S promoter in pUC18, and an *amp* gene that confers ampicillin resistance. All bombardments were carried by co-transforming the two plasmids, p35S and pSMCHS, at a 1:1 ratio.

### Plant materials

The protocorm-like-body (PLB) of *Oncidium* Sharry Baby were used as target tissues. These PLB were multiplied and maintained with

four-week sub-culturing interval in half-strength MS basal salt liquid medium (Murashige and Skoog, 1962) supplemented with 2% (w/v) sucrose, at pH 5.7 on a rotary shaker at 120 rpm with a 16 h photoperiod at  $25 \pm 1^\circ\text{C}$ . Light was provided by fluorescent tubes with a photon flux density of  $150 \mu\text{mol}/\text{m}^2/\text{s}^1$ . After four weeks of sub-culturing, secondary PLB were scalpel separated into individual PLB and sub-cultured onto  $\frac{1}{2}$  MS basal salt solid medium. A total of 20 individual PLB were plated at the centre of a 90-mm diameter Petri dish prior bombardment. Each Petri dish was bombarded once. Each parameter was performed in triplicates.

### Biolistic bombardment

Bombardment was carried out using the PDS-1000/He System (BioRad). All the gold microcarriers, macrocarriers, stopping discs, and rupture discs were purchased from Bio-Rad (Hercules), California, USA. Precipitation of the DNA onto gold microcarriers was done according to the manufacturer's instructions and all the bombardments were performed according to the standard procedure. Bombardments were carried out at the following conditions: 1100 psi rupture disc pressure, rupture disc to macrocarrier distance (11 mm) and stopping plate to target tissue distance (8 cm), 11 mm macrocarrier to stopping plate distance, and 27 mmHg vacuum pressure. Two controls were included, which consisted of unbombarded PLB and PLB bombarded with microcarriers only (not coated with DNA). The bombardment tissues were then incubated at  $25 \pm 0.5^\circ\text{C}$  under 16 hours illumination provided by fluorescent tubes with a photon flux density of  $150 \mu\text{mol}/\text{m}^2/\text{s}^1$  for 24 h prior to transient GFP expression observation. Five critical transformation parameters (size of PLB, time course of GFP transient expression of PLB, concentration of DNA, age of PLB, presence of spermidine and  $\text{CaCl}_2$  in DNA-microcarrier precipitation and duration of single PLB in fresh medium prior bombardment) were assessed in this study.

### GFP monitoring and transient assessment

Assessment of the bombardment parameters was based on transient *gfp* expression and subsequent hygromycin selection. The *gfp* expressing cells were detected using a fluorescence stereomicroscope (Leica MZFL 111) equipped with GFP2 (excitation filter: 480/40 nm; barrier filter: 510 nm) and GFP3 (excitation filter: 470/40 nm and barrier filter: 520/50 nm) filters. An imaging system (Leica DC 200) was attached to the fluorescence microscope to capture the image in real time using the Leica DC Viewer software.

### Selection of putative transformants

Hygromycin was used as a selective agent in the transformation. Four-week-old individual PLB, measuring 4 - 6 mm at the longest diameter, were subjected to a range of hygromycin treatment: 0, 1, 2, 3, 4, 5, 10, 15, 20, 25 and 30  $\mu\text{g}/\text{ml}$ , on solid  $\frac{1}{2}$  MS medium to determine the minimal killing concentration. A total of 60 single PLB for each concentration was tested. The cultures were incubated at  $25 \pm 0.5^\circ\text{C}$  under 16 h illumination provided by fluorescent tubes with a photon flux density of  $150 \mu\text{mol}/\text{m}^2/\text{s}^1$  for 1 month. Data was recorded based on the percentage of surviving PLB. The minimal lethal concentration obtained was used to select the putative transformants in subsequent experiments.

The transformed PLB were maintained on  $\frac{1}{2}$  MS solid medium for one month with subsequent transfer to fresh  $\frac{1}{2}$  MS solid medium supplemented with the optimal hygromycin concentration for putative transformants selection. Surviving and newly formed secondary PLB were further sub-cultured at four-week intervals on fresh  $\frac{1}{2}$  MS solid medium containing the same concentration of hygromycin as

previously for at least 8 months. After 8 months, the PLB were verified for the presence of the transgenes using PCR. Total genomic DNA were extracted from the leaves and analyzed for the presence of the antisense CHS, *hpt II* and *gfp* genes.

#### Genomic DNA isolation from leaves of regenerated bombarded plantlets

Genomic DNA was isolated from the leaves of 5 - 6 cm *in vitro* plantlets using the FTA Classic Card (Whatman®) according to manufacturer's instructions. Briefly, the leaf (0.5 - 1 cm) was placed directly onto the FTA card and covered with a parafilm. Pressure was applied to the leaf with a blunt object such as a tack hammer. The leaf extract was drawn through to the back of the FTA card and the card was then allowed to dry for at least an hour at room temperature. A sample disc measuring 2.0 mm was punched out from the dried spot and placed in a PCR amplification tube. Two hundred microliters (μl) of the FTA Purification Reagent (Whatman®) was added to the PCR tube and incubated for 5 min at room temperature with moderate manual mixing. All spent FTA Purification Reagent was removed and discarded using a pipette. This adding and discarding of the FTA Purification reagent was repeated twice. Two hundred microliters (μl) of TE-1 Buffer (10 mM Tris-HCl, 0.1 mM EDTA, pH 8.0) was then added to the tube and incubated for 5 min at room temperature. All spent TE-1 Buffer was removed. This TE-1 Buffer wash was repeated once again before the disc was allowed to dry at room temperature for about an hour prior to the PCR analysis.

#### PCR analysis

The PCR reactions were carried out using the DNA Thermal Cycle 480 machine (Perkin-Elmer). Amplifications of the *hptII*, CHS antisense and *gfp* genes were each carried out in a 25 μl reaction volume containing 10 x PCR buffer (10 mM Tris-Cl, pH 8.8, 50 mM KCl and 0.8% Nonidet P40), 1.5 mM MgCl<sub>2</sub>, 5 mM dNTP mix (2.5 mM of each nucleotide dATP, dCTP, dGTP, and dTTP), 5 U *Taq* polymerase, 15 pmole primer each and template DNA (disc). Amplification of the antisense CHS gene and *CaMV35S promoter* was done with the following primers: 5'ACGACACTCTCGTCTACTCCA3' and 5' ICCRAANCCRAANAGIACNCCCA 3' resulting in a 1.3 kb fragment. The PCR run condition was set to an initial denaturation step of 1 min at 94°C and subsequent 30 cycles of denaturation (94°C, 1 min), annealing (52°C, 1 min) and extending (72°C, 1 min) followed by a final extending step at 72°C for 5 min. A 750 bp fragment representing the coding region of *sgfpS65T* gene was amplified using the primers 5'ATGGTGAGCAAGGGCGAGGAG3' and 5' TTAGTTGTACAGCTCGTCCAT3' (Elliott et al., 1999) with the PCR run condition of an initial denaturation step of 1 min at 94°C and subsequent 30 cycles of denaturation (94°C, 1 min), annealing (60°C, 1 min) and extending (72°C, 1 min) followed by a final extension step at 72°C for 5 min. The *hptII* gene was amplified using the primers 5' TCGTCCATCAGTTTGCC-3' and 5'AAAAGCCTGAACTCACCGC-3' to give a 500bp fragment and the PCR conditions as described by Upadhyaya et al. (2000).

#### Statistical analysis

All data were analysed using one-way ANOVA and the differences contrasted using DUNCAN's multiple range test performed at the level 5% using SPSS 10.0 (SPSS Inc. USA).


## RESULTS AND DISCUSSION

### Minimal killing concentration of hygromycin on non-transformed *Oncidium Sharry Baby* PLB


Determination of the minimal fatal concentration of selective agent on the target tissue is an essential step in a transformation protocol (Chauvin et al., 1999), which entails killing almost all non-transformed cells and allowing the transformed cells to survive and regenerate. Evaluation of the survival of non-transformed PLB on various concentrations of hygromycin (0, 1, 2, 3, 4, 5, 10, 15, 20, 25 and 30 μg/ml) showed that 100% of the PLB were killed at 4 μg/ml (Figure 1). All the PLB turned dark brown, an indication of cells death. At 3 μg/ml, only 3% of the PLB remained green and growing. This indicates that 4 μg/ml of hygromycin is sufficient to select putative transformants. Contrary to Liao et al. (2003) who reported that 5 μg/ml of hygromycin was sufficient for selecting their *Agrobacterium*-transformed *Oncidium Sharry Baby* PLB. However, as a precaution to avoid potential escapees in the selection process, 5 μg/ml of hygromycin was used as the lethal dosage throughout the selection of putative transformants in all subsequent transformation experiments.

### Assessment of transformation parameters

Six critical transformation parameters were assessed based on transient *gfp* expression and/or hygromycin selection (Table 1, Figure 2) to increase the transformation efficiency. Selection of the right size of target tissue is important because it is an indicator of the developmental stage of the tissue. According to Schenk et al (1998), *gfp* expression varied greatly with respect to shape and size of explant. In this study, 4-week-old single PLB, measuring 1 - 2 mm, 3 - 4 mm and 5 - 6 mm (diameter width) size ranges, were subjected to single bombardment. The results as summarized in Figure 3 show that PLB of 5 - 6 mm size range gave the highest transient *gfp* expression while the 1 - 2 mm size range PLB gave the lowest expression. Differences in *gfp* expression between the 5 - 6 mm size range and both 1 - 2 mm and 3 - 4 mm size ranges were significant ( $p = 0.5$ ) but they were not significant between the 1 - 2 mm and 3 - 4 mm size ranges. However, all the bombarded PLB of size ranges 1 - 2 mm, 3 - 4 mm and 5 - 6 mm turned brown and died after 2 months of selection on 5 μg/ml hygromycin. This may be because this parameter was the first to be optimized, so the transformation condition was not optimal yet. Consequently optimization of other parameters was assessed to improve the transformation efficiency. To avoid low survival rate of smaller size target under the bombarding conditions, bigger PLB of 5 - 6 mm size range was chosen as the target size for subsequent experiments. Another reason is this size range is the



(B)



**Figure 1.** Minimum inhibitory level of hygromycin as selective agent. **(A)** Non-transformed PLB were subjected to various concentrations of hygromycin (0, 1, 2, 3, 4, 5, 10, 15, 20, 25 and 30  $\mu\text{g/ml}$ ) for one month. For each selection, a total of 60 PLB were tested. Data plotted are the means  $\pm$  SD of three replicates. **(B)** Lethal effect exhibited by PLB on media containing different concentrations of hygromycin. A: control, 0  $\mu\text{g/ml}$ ; B: 1  $\mu\text{g/ml}$ ; C: 2  $\mu\text{g/ml}$ ; D: 3  $\mu\text{g/ml}$ ; E: 4  $\mu\text{g/ml}$ ; F: 5  $\mu\text{g/ml}$ . Green signifies surviving PLB (indicated with arrow).

**Table 1.** Bombardment parameters assessed in the co-bombardment of pSM-CHS and p35S into *Oncidium Sharry Baby* PLB.


Parameter	Conditions assessed
Size of PLB	1-2 mm, 3-4 mm, 5-6 mm
Time course of <i>gfp</i> transient expression	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 days
Total amount of plasmid DNA (ratio of plasmids 1:1)	0, 0.5, 1.0, 1.5, 2.0 $\mu$ g/bombardment
Age of PLB	1-, 2-, 3-, 4-week old
Pre-culture period on fresh medium prior bombardment	0, 1, 2, 3, 4, 5, 6, 7 days
*Preparation of DNA coating with gold microcarriers	CS, C, S, O

\*CS means plasmid DNA was coated with microcarriers with the presence of spermidine and calcium chloride.

C means plasmid DNA was coated with microcarriers with presence of calcium chloride only.

S means plasmid DNA was coated with microcarriers with presence of spermidine only.

O means plasmid DNA was coated without the presence of spermidine or calcium chloride.




**Figure 2.** *gfp* expression in bombarded *Oncidium Sharry Baby* PLB. Arrows indicate *gfp* expressing cells (green) visualized under Leica MZFLIII fluorescence microscope equipped with GFP2 filters (Excitation filter: 480/40 nm).

maximum size range of the *Oncidium Sharry Baby* PLB in culture.


Determining the most suitable observation time for *gfp* expression after bombardment is also essential to optimize the transformation system. Tee et al. (2003) reported *gfp* expressing cells were detected 3 hours post-bombardment in their *Dendrobium* PLB. From the result obtained in this study, the earliest *gfp* expressing cell was detected on the same day of bombardment, and it showed that the number of *gfp* expressing cells increased during the first two days (Figure 4). The highest number

of *gfp* expressing cells and the most intense *gfp* expression were observed 2 days post-bombardment. This shows that the best time to quantify cells transiently expressing *gfp* is 2 days post-bombardment.

The use of appropriate concentrations of DNA is important in order to produce efficient DNA-microcarrier binding. In assessing the amount of DNA used per bombardment, it was found that 1.0  $\mu$ g of DNA per bombardment gave the highest transient *gfp* expression (Figure 5A). Reducing or increasing the DNA concentration beyond 1.0  $\mu$ g significantly reduced the *gfp* expression.




**Figure 3.** Effect of different PLB size ranges on transformation efficiency. Data was collected 2 days post-bombardment. Error bars correspond to standard deviation ( $n = 3$ ). The average number of GFP spots is per Petri dish (each dish contained 20 pieces of 4-week-old, 5-6 mm size range PLB). Different letters indicate values are significantly different ( $p = 0.05$ ).



**Figure 4.** Time course of *gfp* expression in PLB. Data was collected 2 days post-bombardment. Error bars correspond to standard deviation ( $n = 3$ ). The average number of GFP spots is per Petri dish (each dish contained 20 pieces of 4-week-old, 5 - 6 mm size range PLB). Different letters indicate values are significantly different ( $p = 0.05$ ): 1<sup>ab</sup>, 2<sup>a</sup>, 3<sup>a</sup>, 4<sup>a</sup>, 5<sup>ab</sup>, 6<sup>abc</sup>, 7<sup>abc</sup>, 8<sup>abc</sup>, 9<sup>cd</sup>, 10<sup>de</sup>, 11<sup>de</sup>, 12<sup>e</sup>, 13<sup>e</sup>, 14<sup>e</sup>.

These might probably due to insufficient amount of microcarriers to coat the available DNA molecules in the solution or maybe the large aggregates were less efficient in penetrating the target tissues thereby increasing cell injury and eventually cells death (Klein et al., 1988; Li et al., 1994). Parveez et al. (1998) also reported that


higher concentrations were not only ineffective but expensive. Only one – fifth of those PLB bombarded with 0.5  $\mu\text{g}$  DNA survived on the hygromycin selection medium (Figure 5B). Even though this DNA concentration did not give the highest *gfp* expression level, it gave the highest number of surviving PLB. In contrast, 1.0  $\mu\text{g}$  DNA



**Figure 5.** Effects of different DNA concentrations on *gfp* expression and survival on hygromycin selection. Each point correspond to standard deviation ( $n = 3$ ), with 20 pieces of 4-week-old, 5 - 6 mm size range PLB per replicate. Different letters indicate values are significantly different ( $p = 0.05$ ). **(A)** *gfp* expression in PLB bombarded with different concentrations of total plasmid DNA (0, 0.5, 1.0, 1.5, 2.0 µg, with plasmids ratio of 1:1). Data was collected 2 days post-bombardment. **(B)** PLB surviving on hygromycin selection. The bombarded PLB were subjected to 5 µg/ml hygromycin selection for 8 months.

per bombardment gave the highest transient *gfp* expression but not the highest percentage of survivals. Overall, the results showed that high concentration of DNA (beyond 1.0 µg) resulted in low survival rates.

Subsequent assessment of the suitable target tissue age for bombardment was carried out. According to Sanford et al. (1993), actively dividing cells were more receptive to receive DNA and gave higher gene expression.




**Figure 6.** Effect of PLB age on *gfp* expression and survival on hygromycin selection. Data was collected 2 days post-bombardment. Error bars correspond to standard deviation ( $n = 3$ ). Different letters indicate values are significantly different ( $p = 0.05$ ). (A) The average number of GFP spots is per Petri dish (20 pieces of 4-week-old, 5-6 mm size range PLB). Each bombardment contains three replicates. (B) Bombarded PLB were subjected to 5  $\mu$ g/ml hygromycin selection for 8 months.

In this work, PLB of 5-6 mm size range obtained from 1, 2, 3 and 4-week-old cultures, were subjected to bombardment, then transient *gfp* expression analysis and hygromycin selection. Neither 1 nor 2-week-old cultures exhibited good *gfp* expression (Figure 6A) compared to the 3 and 4-week-old PLB. After 8 months of hygromycin

selection, only 2% of 1-week-old bombarded PLB survived (Figure 6B). The numbers of surviving and proliferating PLB increased to 10 to 25% when 2, 3 and 4-week-old PLB were used as target tissues. Considering that majority of the PLB at these culture ages ranged from 5 – 6 mm in size, we therefore chose 4-week-old






**Figure 7.** Effect of duration of PLB on fresh medium prior bombardment on *gfp* expression and survival on hygromycin selection. Data was collected 2 days post-bombardment. Error bars correspond to standard deviation ( $n = 3$ ). Different letters indicate values are significantly different ( $p = 0.05$ ). (A) The average number of GFP spots is per Petri dish (20 pieces of 4-week-old, 5 - 6 mm size range PLB). Each bombardment contains three replicates. (B) Percentage of PLB survived after 8 months on 5  $\mu\text{g/ml}$  hygromycin selection.

PLB to be the most suitable culture age for bombardment.

The effect of the culture period of the PLB on fresh medium prior bombardment was further examined. The PLB were sub-cultured onto  $\frac{1}{2}$  MS solid medium for various durations (0, 1, 2, 3, 4, 5, 6 and 7 days) before subjecting to bombardment. The result shows that transient *gfp* expression is proportional to the pre-culturing period from 0 to 7 days (Figure 7A). Pre-culture period

beyond 7 days was not carried out in this study because it was previously observed that beyond 7 days, the PLB started to shoot. After 8 months on hygromycin selection, only 40% of the PLB that were not subjected to a pre-culturing period prior bombardment, survived (Figure 7B). The number of survivals increased to 67% with PLB that were pre-cultured for 1-day prior to bombardment. These bombarded PLB also gave the highest transient *gfp* expression. Hence, this time period was chosen for pre-




**Figure 8.** Effect of spermidine and CaCl<sub>2</sub> in coating of DNA with gold microcarriers on *gfp* expression and survival on hygromycin selection. Calcium chloride and spermidine were added together or alone in the DNA-microcarrier preparation tube. "0" means absence of calcium chloride and spermidine in the preparation. Data was collected 2 days post-bombardment. Error bars correspond to standard deviation (n = 3). The average number of GFP spots is per Petri dish (20 pieces of 4-week-old, 5 - 6 mm size range PLB). Each bombardment contains three replicates. Different letters indicate values are significantly different ( $p = 0.05$ ). (A) *gfp* expression; (B) Selection on 5  $\mu$ g/ml hygromycin for 8 months.

culturing the PLB on fresh medium prior bombardment.

Calcium and spermidine are two important components added in the solution mixture during precipitation of the DNA onto gold microcarriers preparation in order to aid

the binding of DNA molecules to the gold microcarriers. It was found in this study that high transient *gfp* expression was observed when spermidine was used alone in the preparation (Figure 8A). This result parallels that of



**Figure 9. PCR amplification of *chs*, *hptII* and *gfp* genes.** Genomic DNA harvested from randomly picked independent putative transformants that survived on 5  $\mu$ g/ml hygromycin selection were subjected to PCR amplification of the specific transgene. M is 1Kb DNA ladder (A) specific 750 bp *gfp* gene fragment from 3 independent putative transformants (lanes 2, 3, 4), lane 1 is negative control (untransformed plantlet). No amplification of *gfp* gene from putative transformant shown in lane 2; (B) 500bp *hpt II* gene fragment from 9 independent putative transformants (lanes 1 - 9); (C) specific 1.3kb *chs* gene fragment from 7 independent putative transformants (lanes 1 - 7), lane 8 is positive control (plasmid), lane 9 is negative control (untransformed plantlet). No amplification of the *chs* gene was obtained from putative transformant shown in lane 4; (D) shooted PLB that survived on hygromycin selection.

Janna et al. (2006) observation for *Dendrobium Sonia* transformation. However, hygromycin selection recorded higher survivals (67%) for the PLB bombarded with DNA prepared with CaCl<sub>2</sub> alone (Figure 8B), and spermidine is detrimental when included in the preparation process. Hence, only calcium is preferred for the DNA preparation in future experiments.

### PCR analysis

Untransformed PLB turned brown and necrotic on the selection medium within 2 months, while the hygromycin-resistant PLB remained green and regenerated into plantlets during the 8 months of hygromycin selection (Figure 9D). A total of 137 regenerated putative transformants were randomly picked to verify the presence of the transgenes (antisense CHS, *gfp*, and *hpt II*) by PCR analysis. Eleven of the putative transformants were positive for the antisense CHS gene, 61 positives for the *hpt II* gene and 28 for the *gfp* gene (Figure 9A, B, C).

### Conclusion

Theoretically, gene transfer using biolistic gun (PDS 1000/He system) is applicable to all kinds of cells and tissues. However, the DNA delivery conditions and transient *gfp* expression in each species and tissues involved are different and hence, optimization of the delivery conditions is important in order to improve the transformation process (Nan and Kuehnle, 1995). Using the biolistic system, we have successfully introduced unlinked genes on different gene cassettes, p35S which carries a synthetic green fluorescence protein (*sgfp*) gene and pSM-CHS which carries the antisense chalcone synthase (CHS) and hygromycin resistance (*hptII*) genes, into *Oncidium Sharry Baby* PLB. We have shown that hygromycin at 5.0 µg/ml is a good selective agent for *Oncidium Sharry Baby* PLB transformation. Four-week-old PLB of 5 – 6 mm size range and pre-incubated on fresh medium for 1 day prior to bombardment, are optimal for transformation using 1.0 µg DNA prepared in the presence of calcium only. Transformation, selection, and regeneration of the transformants were also carried out successfully where 137 randomly selected regenerated putative transformants were verified to be PCR positives for the antisense CHS, *gfp*, and *hpt II* genes at 11, 28 and 61%, respectively.

In conclusion, this study demonstrates that co-transformation of unlinked genes on different cassettes for *Oncidium Sharry Baby* PLB is possible using the established optimized conditions. This will facilitate the creation of new varieties through the introduction of commercially beneficial genes on different gene cassettes into *Oncidium Sharry Baby* PLB. Because of the time and economic savings without having to re-construct

new gene cassette, the established protocol will also facilitate functional studies of orchid genes at the molecular and cellular levels.

### ACKNOWLEDGEMENTS

The authors are grateful to the Ministry of Science, Technology and Innovation of Malaysia for supporting this work under IRPA grant. Special thanks to Dr. A. Elliot (CSIRO, Brisbane, AUS) and Dr. J. Sheen (Boston, USA) for kindly providing us with the GFP constructs.

### REFERENCES

- Anzai H, Ishii Y, Shichinohe M, Katsumata K, Nojiri C, Morikawa H, Tanaka M (1996). Transformation of *Phalaenopsis* by particle bombardment. *Plant Tissue Cult. Lett.* 13: 265-272.
- Belarmino MM, Mii M (2000). *Agrobacterium* –mediated genetic transformation of a *Phalaenopsis* Orchid. *Plant Cell Rep.* 19: 435-442.
- Chauvin J-E, Marhadour S, Cohat J, Le Nard M (1999). Effets of gelling agents on *in vitro* regeneration and kanamycin efficiency as a selective agent in plant transformation procedures. *Plant Cell Tissue Organ Cult.* 58: 213-217.
- Chia TF, Chan YS, Chua NH (1994). The firefly luciferase gene as a non-invasive reporter for *Dendrobium* transformation. *Plant J.* 6: 441-446.
- Elliott AR, Campbell JA, Dugdale B, Brettell RIS, Grof CPL (1999). Green-fluorescent protein facilitates rapid *in vivo* detection of genetically transformed plant cells. *Plant Cell Rep.* 18: 707-714.
- Janna OA, Maziah M, Parveez GKA, Saleh K (2006). Factors affecting delivery and transient expression of β-glucuronidase gene in *Dendrobium Sonia* protocorm-like-body. *Afr. J. Biotechnol.* 5: 88-94.
- Klein TM, Gradziel T, Fromm ME, Sanford JC (1988). Factors influencing gene delivery into *Zea mays* cells by high velocity microprojectiles. *Bioresour. Technol.* 6: 559-563.
- Kuehnle AR, Sugii N (1992). Transformation of *dendrobium* orchid using particle bombardment of protocorms. *Plant Cell Rep.*, 11: 484-488.
- Li YH, Trembley FM, Seguin A (1994). Transient transformation of pollen and embryogenic tissues of white spruce (*Picea glauca Moench*) resulting from microprojectile bombardment. *Plant Cell Rep.* 13: 661-665.
- Liau CH, You SJ, Prasad V, Hsiao, HH, Lu JC, Yang NS, Chan MT (2003). *Agrobacterium tumefaciens*-mediated transformation of an *Oncidium* orchid. *Plant Cell Rep.* 21: 993-998.
- Malabadi RB, Nataraja K (2007). Genetic transformation of *Vanilla planifolia* by *Agrobacterium-tumefaciens* using shoot tip sections. *Res. J. Bot.* 2(2): 86-94.
- Murashige T, Skoog T (1962). A revised method for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant* 15: 473-497.
- Nan GL, Kuehnle AR (1995). Factors affecting gene delivery by particle bombardment of *Dendrobium* orchids. *In Vitro Cell Dev. Biol. Plant.* 31: 131-136.
- Parveez GKA, Chowdhury MKU, Saleh NM (1998). Physical parameters affecting transient GUS gene expression in oil palm (*Elaeis guineensis* Jacq.) embryogenic calli via microprojectile bombardment. *Indian Crop. Prod.* 8: 17-27.
- Ramesh M, Gupta AK (2005). Transient expression of beta-glucuronidase gene in indica and japonica rice (*Oryza sativa* L.) callus after different stages of co-bombardment. *Afr. J. Biotechnol.* 4(7): 596-600.
- Sanford J, Smith FD, Russel JA (1993). Optimizing the biolistic process for different biological applications. *Method Enzymol.* 217: 483-509.
- Schenk PM, Elliot AR, Manner JM (1998). Assessment of transient gene expression in plant tissue using the green fluorescent protein as a reference. *Plant Mol. Biol. Rep.* 16: 313-322.

- Sheen J, Hwang S, Niwa Y, Kobayashi H, Galbraith DW (1995). Green fluorescent protein as a new vital marker in plant cells. *Plant J.* 8: 777-784.
- Tee CS, Marziah M (2005). Optimization of biolistic bombardment parameters for *Dendrobium* Sonia 17 calluses using GFP and GUS as the reporter system. *Plant Cell Tissue Organ Cult.* 80: 77-89.
- Tee CS, Marziah M, Tan CS, Abdullsh MP (2003). Evaluation of different promoters driving the GFP reporter gene and selected target tissues for particle bombardment of *Dendrobium* Sonia 17. *Plant Cell Rep.* 21: 452-458.
- Upadhyaya NM, Surin B, Ramm K, Gaudron J, Schunmann PHD, Taylor W, Waterhouse PM, Wang MB (2000). *Agrobacterium*-mediated transformation of Australian rice cultivars Jarrah and Amaroo using modified promoters and selectable markers. *Austr. J. Plant Physiol.* 27: 201-210.
- Yang J, Lee HJ, Shin DH, Oh SK, Seon JH, Paek KY, Han KH (1999). Genetic transformation of *Cymbidium* orchid by particle bombardment. *Plant Cell Rep.* 18: 978-984.
- Yao Q, Cong L, He GY, Chang JL, Li KX, Yang GX (2007). Optimization of wheat co-transformation procedure with gene cassettes resulted in an improvement in transformation efficiency. *Mol. Biol. Rep.* 34(1): 61-67.
- Yu H, Yang SH, Goh C J (2001). *Agrobacterium*-mediated transformation of a *Dendrobium* orchid with the 1 knob gene DOH1. *Plant Cell Rep.* 20: 301-305.
- Zhao Y, Qian Q, Wang HZ, Huang DN (2007). Co-transformation of gene expression cassettes via particle bombardment to generate safe transgenic plant without any unwanted DNA. *In Vitro Cell. Dev. Biol.* 43(4): 325-334.