

## Review

# Inhibitory effect of essential oil on aflatoxin activities

Lokman Alpsoy

Department of Biology, Faculty of Art and Science, Fatih University, Istanbul, Turkey.  
E-mail: [lalpsoy@fatih.edu.tr](mailto:lalpsoy@fatih.edu.tr). Tel: +90(212)8663300-2029. Fax: +90(212)8663402.

Accepted 29 March, 2010

**Aflatoxins, which are well-known to be mutagenic, carcinogenic, teratogenic, hepatotoxic and immunosuppressive, also inhibit several metabolic systems. Aflatoxins are biologically active secondary metabolites produced by certain strains of *Aspergillus parasiticus*, *Aspergillus nominus* and *Aspergillus flavus*. Many different substances, such as essential oils, flavanoids, could inhibit the aflatoxin production and growth of *Aspergillus*. In this study, aflatoxins biosynthesis, aflatoxins damaged and aflatoxins with essential oils interaction are evaluated.**

**Key words:** Aflatoxins, essential oils, antioxidant, oxidative stress.

## INTRODUCTION

Aflatoxins (*Aspergillus flavus* toxins) are biologically active secondary metabolites produced by certain strains of *Aspergillus parasiticus*, *Aspergillus nominus* and *Aspergillus flavus* (Cotty et al., 1994). The aflatoxin producing fungi are widely distributed in nature and can grow over a wide range of environmental conditions (Holmquist et al., 1983). Aflatoxins have been detected in cereal grains, oil seeds, fermented beverages made from grains, milk, cheese, meat, nut products, fruit juice and numerous other agricultural commodities (Bullerman, 1986).

Aflatoxins have been shown to be hepatotoxic, carcinogenic, mutagenic and teratogenic to different species of animals (Wogan et al., 1974; Eaton and Gallagher, 1994; Abdel-Wahhab et al., 1999). Aflatoxin B<sub>1</sub> (AFB<sub>1</sub>) is the most prevalent and carcinogenic of the aflatoxins and the International Agency for Research on Cancer (IARC) classify AFB<sub>1</sub> as a group I carcinogen (that is, an agent that is carcinogenic to humans). Epidemiological studies also indicate that areas in the world with high levels of aflatoxin are correlated with high incidence of liver cancer (IARC, 1985).

AFB<sub>1</sub> caused damage by two different way in the cells. Firstly, AFB<sub>1</sub> (C<sub>17</sub>H<sub>12</sub>O<sub>6</sub>) is activated to AFB<sub>1</sub>-8,9-oxide

and forms adduct primarily at N7 position of guanine and is responsible for its mutagenic and carcinogenic effects (Wang and Groopman, 1999; Denissenko et al., 1999). Secondly, aflatoxins especially AFB<sub>1</sub>, produce reactive oxygen species (ROS) such as superoxide radical anion, hydrogen peroxide and lipid hydroperoxides; though these do not appear to interact with DNA, but they are precursors to the hydroxyl radical. The hydroxyl radicals interact with DNA and produces mutations (Halliwell and Gutteridge, 1999).

Numerous diverse compounds and extracts containing activity inhibitory to aflatoxin biosynthesis have been reported. The most of these inhibitors are plant-derived such as phenylpropanoids, terpenoids and alkaloids (Holmes et al., 2008). A group of plant-derived inhibitors is essential oils (EO) that possess antimicrobial activities against *A. parasiticus* and/or *A. flavus* (Rasoli and Owlia, 2005; Kumar et al., 2007; Rasoli et al., 2008; Bluma and Etcheverry, 2008).

Up to date, no review directly has been carried out to evaluate the protective effects of essential oils against the aflatoxins. The objective of this study is to explain the protective effects of essential oil against the growth of *A. parasiticus* and *A. flavus*, synthesis of aflatoxins as well as the damage of aflatoxins.

**Abbreviations:** AFB<sub>1</sub>, Aflatoxin B<sub>1</sub>, ROS, reactive oxygen species; EO, essential oils; 8-OHdG, 8-hydroxy-2'-deoxyguanosine; G.C, guanine-cytosine; T.A, thymine-adenine; SOD, superoxide dismutase; GPx, glutathione peroxidase; CAT, catalase.

## BIOSYNTHESIS OF AFLATOXINS

Aflatoxins belong to the polyketide class of secondary metabolites and are synthesized by enzymes encoded




Figure 1. Inhibition of aflatoxin synthesis.

within a large gene cluster (Yabe and Nakajima 2004; Yu et al., 2004). The initial step in the generation of the polyketide backbone of aflatoxins is proposed to involve polymerization of acetate and nine malonate units (with a loss of  $\text{CO}_2$ ) by a polyketide synthetase in a manner analogous to fatty acid biosynthesis (Dutton, 1988; Bhatnagar et al., 1992).

Synthesis of aflatoxins is control by specific enzymes which are expressed by DNA through many steps. Each step in gene expression, transcription, RNA transport and processing, translation, protein processing and localization can in theory be inhibited by natural plant products or other agents (Trail et al., 1995).

Many inhibitors of aflatoxin biosynthesis may act at three levels: (1) Modulate environmental and physiological factors affecting aflatoxin biosynthesis, (2) inhibit signaling circuits upstream of the biosynthetic pathway, or (3) directly inhibit gene expression or enzyme activity in the pathway (Figure 1). While the mode of action of most inhibitory compounds is unknown, there is a little evidence for the described compounds having an effect on gene transcription or enzyme activity of individual steps in the biosynthetic pathway. More likely, the known inhibitory compounds either alter known environmental and physiological modulators of aflatoxin biosynthesis or they alter signaling transduction pathways in the upstream regulatory network (Figure 1) (Holmes et al., 2008).

## DAMAGES OF AFLATOXINS IN CELLS

Aflatoxins, which are well-known to be potentially mutagenic, carcinogenic, teratogenic, hepatotoxic and immunosuppressive, also inhibit several metabolic systems (Minto and Townsend, 1997; Wogan, 1999). The carcinogenic mechanism of aflatoxin B<sub>1</sub> (AFB<sub>1</sub>) has been extensively studied. It has been shown that AFB<sub>1</sub> is activated by hepatic cytochrome P450 enzyme system to produce a highly reactive intermediate, AFB<sub>1</sub>-8,9-epoxide, which subsequently binds to nucleophilic sites in DNA and the major adduct 8,9-dihydro-8-(N7 guanyl)-9-hydroxy AFB<sub>1</sub> (AFB<sub>1</sub> N7-Gua) is formed (Sharma and Farmer, 2004; Klein, et al., 2002; Bedard and Massey, 2006). The formation of AFB<sub>1</sub>-DNA adducts is regarded as a critical step in the initiation of AFB<sub>1</sub>-induced hepatocarcinogenesis (Preston and Williams, 2005).

AFB<sub>1</sub> also has been shown to induce 8-hydroxy-2'-deoxyguanosine (8-OHdG) formation in rat (Shen et al., 1995; Yarborough et al., 1996) and duck (Barraud et al., 2001) liver in *in vivo* treatment. The mutagenic specificity of 8OHGua provides a potential tag for assessing the role of ROS mutagenesis in human cancer. It is of interest that the spectrum of p53 tumor suppressor gene mutations includes G.C+T.A transversions in about half of non-small cell carcinomas of the lung and nearly three-quarters of primary liver carcinomas (Hollstein et al.,

1991).

Although the mechanism underlying the hepatotoxicity of aflatoxins is not fully understood, several reports suggest that toxicity may ensue through the generation of intracellular ROS during the metabolic processing of AFB<sub>1</sub> by cytochrome P450 in the liver (Towner et al., 2003; Sohn et al., 2003). Free radicals provoked by various environmental chemicals as well as endogenous metabolism are involved in a number of diseases like tumors, inflammation, shock, atherosclerosis, diabetes, infertility, gastric mucosal injury and ischemia due to the oxidative damage to DNA, lipids and proteins and which can result in failure of cellular functions (Kasai, 2002). Free radicals also contribute to G. C + T. A transversions by the production of 8OHGua in liver DNA, therapies designed to reduce damage by oxygen free radicals (Ames, 1983) during chronic hepatitis would be predicted to delay the onset of primary liver cancer (Cheng et al., 1992).

To control the level of ROS and to protect cells under stress conditions, living tissues contain several enzymes (SOD, GPx, and CAT) and many antioxidant substances. The effect of ROS is balanced by the antioxidant action of non-enzymatic antioxidants, as well as by antioxidant enzymes. Such antioxidant defenses are extremely important as they represent the direct removal of free radicals (prooxidants), thus, providing maximal protection for biological sites (Valko et al., 2006).

A lot of antioxidant compounds such as EO, phenolics compounds and secondary metabolites, which are synthesized by plants, serve in defense against ROS. The antioxidant properties of essential oil of plant origin have been studied in recent years. A strong correlation has been found between the essential oil level and the antioxidant activity potential.

Some EOs and other extracts (vitamins, riboflavin, carotenoids, beta-carotene, alfa-carotene, lycopene, ascorbic acid, curcumin, several flavonoids, phenolic compounds and synthetic phenolic compounds) of plants could potentially provide protection against aflatoxins especially AFB<sub>1</sub> (Nyandieka et al., 1990; Webster et al., 1996; Rasoli and Owlia, 2005; Agar et al., 2005; Kumar et al., 2007; Rasoli et al., 2008; Bluma and Etcheverry, 2008; Alpsoy et al., 2009). In addition, many essential oils could reduce toxic and mutagenic effect of aflatoxins. Most studies indicated that antiaflatoxigenic properties may be due to inhibition of penetration of *A. flavus* and *A. parasiticus* (Rasoli et al., 2008; Bluma and Etcheverry, 2008).

## IMPACT OF ESSENTIAL OIL ON GROWTH OF ASPERGILLUS

Essential oils are volatile, natural and complex compounds characterized by a strong odour and are produced from aromatic plants as secondary metabolites. They are

usually obtained by steaming or hydro-distillation which was first developed in the Middle Ages by the Arabs. Having been known for their antiseptic and medical properties such as bactericidal, virucidal and fungicidal and fragrance, they were used in embalment, preservation of foods and for the treatment of antimicrobial, analgesic, sedative, anti-inflammatory, spasmolytic and locally anesthetic diseases. Essential oils can contain about 20-60 components in quite different concentrations. They are characterized by two or three major components at fairly high concentrations (20-70%) compared to others present in trace amounts (Bakkali et al., 2008).

Until now, many studies have revealed that *Aspergillus* growth was completely inhibited by many plants EOs. The effects of essential oils of 58 plant species (18 Family) were examined on the development of *A. flavus* and/or *A. parasiticus* (Table 1). EOs were extracted from leaf, stem and flower and they were also purchased from local market. Used different concentration of EOs was found to inhibit the development of *Aspergillus* species.

There are complex interactions of environmental factors, like water availability, which influence the efficacy of essential oils. It is possible to use a combination of them to reduce growth and aflatoxin production of *A. flavus* and *A. parasiticus*. The antifungal efficacy of plant essential oils may be attributable to the oil compositions. This result may be explained by the high content of some of these substances in the plants essential oils. Many researches indicate that the components of essential oil reduce the growth rate of *A. flavus* and *A. parasiticus*. Major components can constitute up to 85% of the EO, whereas other components are present only as a trace.

Some researchers reported that there is a relationship between the chemical structures of the most abundant compounds in the essential oils and the antimicrobial activity. Compositional analysis of EOs showed that compounds such as carvacrol, r-cimene, a-terpinolene, anethol and eugenol were the main components present in the different EO studied. According to Baydar et al. (2004) phenolic components, such as eugenol, anethole, carvacrol, its precursors r-cimene and g-terpinolene and its isomers thymol, are chiefly responsible for the antimicrobial properties of essential oils. Also, the physical nature of essential oils, that is, low molecular weight combined with pronounced lipophilic tendencies allow them to penetrate cell membrane more quickly than other substances (Pawar and Thaker, 2007).

However, there is evidence that minor components have a critical part to play in antimicrobial activity, possibly by producing a synergic effect between other components (Burt, 2004). The antimicrobial activity of essential oils or their constituents such as thymol, carvacrol and vanillin could be in different ways; (1) The result could be in the form of damage to the enzymatic cell system, including those associated with energy production and synthesis of structural compounds (Conner and Beuchat, 1984 a,b), (2) denaturation of the enzymes responsible for spore germination or interference

**Table 1.** Examples of plants that has antifungal activity.

| Species                           | Family     | Part of plant       | Aspergillus                             | References   |
|-----------------------------------|------------|---------------------|---|--|
| <i>Rosmarinus officinalis</i>     | Lamiaceae  | Leaf                | <i>A. parasiticus</i>                   | Rasoli et al., 2008  |
| <i>Ocimum basilicum</i>           | Lamiaceae  | Leaf                | <i>A. flavus</i> , <i>A. paraiticus</i> | Montes-Belmont and Carvajal, 1998;<br>Soliman and Badeaa, 2002; Atanda et al., 2007                        |
| <i>Thymus eriocalyx</i>           | Lamiaceae  | Leaf                | <i>A. parasiticus</i>                   | Rasoli and Abyaneh, 2004   |
| <i>Thymus x-porlock</i>           | Lamiaceae  | Leaf                | <i>A. parasiticus</i>                   | Rasoli and Abyaneh, 2004   |
| <i>Satureja hortensis</i>         | Lamiaceae  | Leaf                | <i>A. parasiticus</i>                   | Razzaghi-Abyaneh et al., 2008  |
| <i>Ocimum gratissimum</i>         | Lamiaceae  | Leaf                | <i>A. flavus</i>                        | Neguefact et al., 2004   |
| <i>Thymus vulgaris</i>            | Lamiaceae  | All of plant, Leaf  | <i>A. flavus</i> , <i>A. paraiticus</i> | Montes-Belmont and Carvajal, 1998;<br>Soliman and Badeaa, 2002; Neguefact et al., 2004; Kumar et al., 2007 |
| <i>Mentha viridis</i>             | Lamiaceae  | Leaf                | <i>A. flavus</i> , <i>A. paraiticus</i> | Soliman and Badeaa, 2002   |
| <i>Mentha piperita</i>            | Lamiaceae  | Leaf, Leaf and stem | <i>A. flavus</i> , <i>A. paraiticus</i> | Montes-Belmont and Carvajal, 1998; Bluma et al., 2007  |
| <i>Origanum vulgare</i>           | Lamiaceae  | Leaf, Leaf and stem | <i>A. flavus</i> , <i>A. paraiticus</i> | Montes-Belmont and Carvajal, 1998; Bluma et al., 2007  |
| <i>Minthostachys verticillata</i> | Lamiaceae  | Leaf and Stem       | <i>A. flavus</i> , <i>A. paraiticus</i> | Bluma et al., 2007   |
| <i>Lavandula officinalis</i>      | Lamiaceae  | Leaf                | <i>A. flavus</i>                        | Kumar et al., 2007   |
| <i>Mentha arvensis</i>            | Lamiaceae  | Leaf                | <i>A. flavus</i>                        | Kumar et al., 2007   |
| <i>Ocimum canum</i>               | Lamiaceae  | Leaf                | <i>A. flavus</i>                        | Kumar et al., 2007   |
| <i>Pogostemon cablin</i>          | Lamiaceae  | Leaf                | <i>A. flavus</i>                        | Kumar et al., 2007   |
| <i>Matricaria chamomilla</i>      | Asteraceae | Leaf                | <i>A. flavus</i> , <i>A. paraiticus</i> | Soliman and Badeaa, 2002   |
| <i>Calendula officinalis</i>      | Asteraceae | Leaf                | <i>A. flavus</i> , <i>A. paraiticus</i> | Soliman and Badeaa, 2002   |
| <i>Achillea millefolium</i>       | Asteraceae | Leaf                | <i>A. flavus</i> , <i>A. paraiticus</i> | Soliman and Badeaa, 2002   |
| <i>Achillea fragrantissima</i>    | Asteraceae | Leaf                | <i>A. flavus</i> , <i>A. paraiticus</i> | Soliman and Badeaa, 2002   |
| <i>Artemisia nilagirica</i>       | Asteraceae | Leaf                | <i>A. flavus</i>                        | Kumar et al., 2007   |
| <i>Eupatorium cannabinum</i>      | Asteraceae | Leaf                | <i>A. flavus</i>                        | Kumar et al., 2007   |
| <i>Chrysactinia mexicana</i>      | Asteraceae | Flower              | <i>A. flavus</i>                        | Alvarez-Castellanos et al., 2001   |
| <i>Trachyspermum copticum</i>     | Apiaceae   | Seed                | <i>A. parasiticus</i>                   | Rasoli et al., 2008  |
| <i>Coriandrum sativum</i>         | Apiaceae   | Leaf                | <i>A. parasiticus</i>                   | Atanda et al., 2007  |
| <i>Pimpinella anisum</i>          | Apiaceae   | Leaf, Seed          | <i>A. flavus</i> , <i>A. paraiticus</i> | Soliman and Badeaa, 2002; Bluma et al., 2007;<br>Bluma et al., 2008  |
| <i>Carum carvi</i>                | Apiaceae   | Leaf                | <i>A. flavus</i> , <i>A. paraiticus</i> | Soliman and Badeaa, 2002; Kumar et al., 2007   |
| <i>Foeniculum vulgare</i>         | Apiaceae   | Leaf                | <i>A. flavus</i> , <i>A. paraiticus</i> | Soliman and Badeaa, 2002   |
| <i>Entandrophragma utile</i>      | Meliaceae  | Leaf                | <i>A. parasiticus</i>                   | Anthony, 2006  |
| <i>Khaya grandifoliola</i>        | Meliaceae  | Leaf                | <i>A. parasiticus</i>                   | Anthony, 2006  |
| <i>Lovoa trichilioides</i>        | Meliaceae  | Leaf                | <i>A. parasiticus</i>                   | Anthony, 2006  |
| <i>Pseudoedrela kotschyi</i>      | Meliaceae  | Leaf                | <i>A. parasiticus</i>                   | Anthony, 2006  |
| <i>Trichilia heudelotii</i>       | Meliaceae  | Leaf                | <i>A. parasiticus</i>                   | Anthony, 2006  |

Table 1. Contd

|                                 |               |                              |   |   |
|---------------------------------|---------------|------------------------------|---|---|
| <i>Cymbopogon citratus</i>      | Poaceae       | Leaf, Commerical             | <i>A. flavus</i>                        | Paranagama et al., 2003; Neguefact et al., 2004; Souza et al., 2005; Helal et al., 2007 |
| <i>Syzygium aromaticum</i>      | Myrtaceae     | Leaf, Airal part, Flower bud | <i>A. flavus</i>                        | Montes-Belmont and Carvajal, 1998; Bluma et al., 2007; Bluma et al., 2008               |
| <i>Cymbopogon flexuosus</i>     | Poaceae       | Leaf                         | <i>A. flavus</i>                        | Kumar et al., 2007  |
| <i>Cymbopogon martinii</i>      | Poaceae       | Leaf                         | <i>A. flavus</i>                        | Kumar et al., 2007  |
| <i>Melaleuca leucadendron</i>   | Myrtaceae     | Leaf                         | <i>A. flavus</i>                        | Kumar et al., 2007  |
| <i>Vetiveria zizanioides</i>    | Poaceae       | Leaf                         | <i>A. flavus</i>                        | Kumar et al., 2007  |
| <i>Eucalyptus globulus</i>      | Myrtaceae     | Commerical, Airal part       | <i>A. flavus</i>                        | Souza et al., 2005; Bluma et al., 2007  |
| <i>Eugenia uniflora</i>         | Myrtaceae     | Commerical                   | <i>A. flavus</i>                        | Souza et al., 2005  |
| <i>Cinnamomum cassia</i>        | Lauraceae     | Leaf                         | <i>A. flavus</i>                        | Atanda et al., 2007   |
| <i>Laurus nobilis</i>           | Lauraceae     | Leaf                         | <i>A. flavus</i>                        | Atanda et al., 2007   |
| <i>Cinnamomum zeylanicum</i>    | Lauraceae     | Commerical; Leaf             | <i>A. flavus</i> , <i>A. paraiticus</i> | Montes-Belmont and Carvajal, 1998; Soliman and Badeaa, 2002; Carmo et al., 2008         |
| <i>Lippia alba</i>              | Verbenaceae   | Commerical, Leaf             | <i>A. flavus</i>                        | Souza et al., 2005; Kumar et al., 2007  |
| <i>Lippia turbinata</i>         | Verbenaceae   | Airal part                   | <i>A. flavus</i>                        | Bluma et al., 2007; Bluma et al., 2008  |
| <i>Lippia microphylla</i>       | Verbenaceae   | Commerical                   | <i>A. flavus</i>                        | Souza et al., 2005  |
| <i>Teloxys ambrosioides</i>     | Amaranthaceae | Leaf                         | <i>A. flavus</i>                        | Montes-Belmont and Carvajal, 1998   |
| <i>Aegle marmelos</i>           | Rutaceae      | Leaf                         | <i>A. flavus</i>                        | Kumar et al., 2007  |
| <i>Chenopodium ambrosioides</i> | Amaranthaceae | Leaf                         | <i>A. flavus</i>                        | Kumar et al., 2007  |
| <i>Citrus limon</i>             | Rutaceae      | Commerical                   | <i>A. flavus</i>                        | Souza et al., 2005  |
| <i>Monodora myristica</i>       | Annonaceae    | Seed                         | <i>A. flavus</i>                        | Neguefact et al., 2004  |
| <i>Zingiber officinale</i>      | Zingiberaceae | All of plant, Leaf           | <i>A. flavus</i>                        | Neguefact et al., 2004; Kumar et al., 2007  |
| <i>Agrimonia eupatoria</i>      | Rosaceae      | Leaf                         | <i>A. flavus</i> , <i>A. paraiticus</i> | Soliman and Badeaa, 2002  |
| <i>Peumus boldus</i>            | Monimiaceae   | Leaf, Commerical             | <i>A. flavus</i> , <i>A. paraiticus</i> | Souza et al., 2005; Bluma et al., 2007; Bluma et al., 2008                              |
| <i>Pelargonium graveolens</i>   | Geraniaceae   | Leaf                         | <i>A. flavus</i>                        | Kumar et al., 2007  |
| <i>Santalum album</i>           | Santalaceae   | Wood                         | <i>A. flavus</i>                        | Kumar et al., 2007  |
| <i>Hedeoma multiflora</i>       | Malpighiaceae | Airal part, Leaf             | <i>A. flavus</i>                        | Bluma et al., 2007, Bluma et al., 2008  |
| <i>Sardinia juniperus</i>       | Cupressaceae  | Leaf                         | <i>A. flavus</i>                        | Cosentino et al, 2003   |

with the aminoacid involved in germination (Nychas, 1995) and (3) irreversible damage in cell

wall, cell membrane and cellular organelles when *A. parasiticus* and *A. flavus* were exposed to

different essential oils (Rasoli and Owlia, 2005; Helal et al., 2007).

**Table 2.** Examples of plants that has antiaflatoxigenic activity.

| Species                  | Family        | Part of plant                | Aflatoxins               | References  |
|--------------------------|---------------|------------------------------|--------------------------|---|
| <i>R. officinalis</i>    | Lamiaceae     | Leaf                         | Aflatoxin                | Rasoli et al., 2008   |
| <i>O.basilicum</i>       | Lamiaceae     | Leaf                         | Aflatoxin, AFB1 and AFG1 | Montes-Belmont and Carvajal, 1998; Soliman and Badeaa, 2002; Atanda et al., 2007                        |
| <i>T. eriocalyx</i>      | Lamiaceae     | Leaf                         | Aflatoxin                | Rasoli and Abyaneh, 2004  |
| <i>T.x-porlock</i>       | Lamiaceae     | Leaf                         | Aflatoxin                | Rasoli and Abyaneh, 2004  |
| <i>S. hortensis</i>      | Lamiaceae     | Leaf                         | AFB1 and AFG1            | Razzaghi-Abyaneh et al., 2008   |
| <i>T. vulgaris</i>       | Lamiaceae     | All of plant, Leaf           | Aflatoxin                | Montes-Belmont and Carvajal, 1998; Soliman and Badeaa, 2002; Neguefact et al., 2004; Kumar et al., 2007 |
| <i>M. viridis</i>        | Lamiaceae     | Leaf                         | Aflatoxin                | Soliman and Badeaa, 2002  |
| <i>M. chamomilla</i>     | Asteraceae    | Leaf                         | Aflatoxin                | Soliman and Badeaa, 2002  |
| <i>C. officinalis</i>    | Asteraceae    | Leaf                         | Aflatoxin                | Soliman and Badeaa, 2002  |
| <i>A. millefolium</i>    | Asteraceae    | Leaf                         | Aflatoxin                | Soliman and Badeaa, 2002  |
| <i>A. fragrantissima</i> | Asteraceae    | Leaf                         | Aflatoxin                | Soliman and Badeaa, 2002  |
| <i>T. copticum</i>       | Apiaceae      | Seed                         | Aflatoxin                | Rasoli et al., 2008   |
| <i>C. sativum</i>        | Apiaceae      | Leaf                         | AFB1 and AFG1            | Atanda et al., 2007   |
| <i>P. anisum</i>         | Apiaceae      | Leaf, Seed                   | Aflatoxin                | Soliman and Badeaa, 2002; Bluma et al., 2007; Bluma et al., 2008  |
| <i>C.carvi</i>           | Apiaceae      | Leaf                         | Aflatoxin                | Soliman and Badeaa, 2002; Kumar et al., 2007  |
| <i>F. vulgare</i>        | Apiaceae      | Leaf                         | Aflatoxin                | Soliman and Badeaa, 2002  |
| <i>E. utile</i>          | Meliaceae     | Leaf                         | Aflatoxin                | Anthony, 2006   |
| <i>K. grandifoliola</i>  | Meliaceae     | Leaf                         | Aflatoxin                | Anthony, 2006   |
| <i>L. trichilioides</i>  | Meliaceae     | Leaf                         | Aflatoxin                | Anthony, 2006   |
| <i>P.kotschyi</i>        | Meliaceae     | Leaf                         | Aflatoxin                | Anthony, 2006   |
| <i>T. heudilotii</i>     | Meliaceae     | Leaf                         | Aflatoxin                | Anthony, 2006   |
| <i>C. citratus</i>       | Poaceae       | Leaf, Commerical             | AFB1                     | Paranagama et al., 2003; Neguefact et al., 2004; Souza et al., 2005; Helal et al., 2007                 |
| <i>S. aromaticum</i>     | Myrtaceae     | Leaf, Airal part, Flower bud | Aflatoxin, AFB1          | Montes-Belmont and Carvajal, 1998; Bluma et al., 2007; Bluma et al., 2008                               |
| <i>E. globulus</i>       | Myrtaceae     | Commerical, Airal part       | AFB1                     | Souza et al., 2005; Bluma et al., 2007  |
| <i>C. cassia</i>         | Lauraceae     | Leaf                         | AFB1 and AFG1            | Atanda et al., 2007   |
| <i>L. nobilis</i>        | Lauraceae     | Leaf                         | AFB1 and AFG1            | Atanda et al., 2007   |
| <i>C. zeylanicum</i>     | Lauraceae     | Commerical; Leaf             | Aflatoxin                | Montes-Belmont and Carvajal, 1998; Soliman and Badeaa, 2002; Carmo et al., 2008                         |
| <i>L. turbinata</i>      | Verbenaceae   | Airal part                   | Aflatoxin, AFB1          | Bluma et al., 2007; Bluma et al., 2008  |
| <i>C. ambrosioides</i>   | Amaranthaceae | Leaf                         | AFB1                     | Kumar et al., 2007  |
| <i>A. eupatoria</i>      | Rosaceae      | Leaf                         | Aflatoxin                | Soliman and Badeaa, 2002  |
| <i>P. boldus</i>         | Monimiaceae   | Leaf, Commerical             | Aflatoxin                | Souza et al., 2005; Bluma et al., 2007; Bluma et al., 2008  |
| <i>H. multiflora</i>     | Malpighiaceae | Airal part, Leaf             | Aflatoxin, AFB1          | Bluma et al., 2007, Bluma et al., 2008  |

## INHIBITION OF AFLATOXIN PRODUCTION BY ESSENTIAL OILS

The effects of EO and its principal components showed inhibitory activity on aflatoxin biosynthesis by *A. flavus*. Many researcher reported that EOs of plants was able to inhibit both growth and/or mycotoxin production. The antiaflatoxigenic activity of EOs of different 32 plants are indicated by many researcher (Table 2). For instance, the

EOs of *Pimpinella anisum*, *Peumus boldus*, *Hoya multiflora*, *Syzygium aromaticum* and *Lippia turbinata* inhibited aflatoxin production.

Essential oils, such as anisum and boldus, could be safely used as a preservative material on some foods because they stopped fungal growth and AFB<sub>1</sub> accumulation. They could also be added to grain in storage to protect it from fungal infection. These oils could be used as a substitute for chemical fungicides.

They may also prove valuable as 'lead structure' for the development of synthetic compounds as they are natural and nontoxic to humans and animals alike (Soliman and Badeaa, 2002; Bluma et al., 2007). The antiaflatoxigenic actions of EO may be related to inhibition of the ternary steps of aflatoxin biosynthesis involving lipid peroxidation and oxygenation (Bluma et al., 2007)

## INHIBITION OF AFLATOXIN DAMAGE BY ESSENTIAL OIL

The essential oils can decrease the damaged effect of aflatoxins by two different ways. Firstly, DNA binding formation of aflatoxins is reduced by essential oils. Secondly, aflatoxins cause increase of reactive oxygen species and essential oils react with ROS. Therefore, essential oils protect the cells from harmful impact of aflatoxins.

Recently, the natural products such as plant extracts have been identified as potential candidates against AFB<sub>1</sub>. A few study show that essential oils reduce DNA binding of aflatoxin. Essential oils from common spices such as nutmeg, ginger, cardamom, celery, xanthoxylum, black pepper, cumin and coriander were tested for their ability to suppress the formation of DNA adducts by AFB<sub>1</sub> *in vitro* in a microsomal enzyme-mediated reaction. All oils were found to inhibit adduct formation very significantly and in a dose-dependent manner. The adduct formation appeared to be modulated through the action on microsomal enzymes, because an effective inhibition on the formation of activated metabolite was observed with each oil. The enzymatic modulation is perhaps due to the chemical constituents of the oils and this could form a basis for their potential anticarcinogenic roles (Hashim et al., 1994). In another research, the effects of garlic oil, such as diallyl disulfide (DADS) and diallyl sulfide (DAS) on AFB<sub>1</sub>-induced DNA damage in cultured primary rat hepatocytes are shown. About 0.5 and 2 mM DAS or 0.5 and 1 mM DADS significantly decreased the DNA damage induced by AFB<sub>1</sub> as compared with the AFB<sub>1</sub> control, according to the unscheduled DNA synthesis test (Shenn et al., 2001).

## CONCLUSION

These results indicate the potential of essential oils as natural preservatives in food against *A. parasiticus* and *A. flavus*, the well known casual agents of foodborne diseases and food poisonings. There are a few research on aflatoxin-induced oxidative damage and essential oil interaction. Therefore, this area needs more study.

## REFERENCES

Abdel-Wahhab MA, Nada SA, Amra HA (1999). Effect of aluminosilicate and bentonite on aflatoxin-induced developmental toxicity in rats. J.

- Appl. Toxicol. 19: 199-204.
- Agar G, Alpsoy L, Yildirim N (2005). The protective role of selenium against the genotoxicity induced by aflatoxin B<sub>1</sub> in root cells of crop plants. *Fresenius Environ. Bull.* 14(10): 849-853.
- Alpsoy L, Yildirim A, Agar G (2009). The Antioxidant Effects of Vitamin A, C and E on Aflatoxin B<sub>1</sub>-Induced Oxidative Stress in Human Lymphocytes. *Toxicol. Ind. Health* 25(2): 121-127.
- Alvarez-Castellanos PP, Bishop CD, Pascual-Villalobos MJ (2001). Antifungal activity of the essential oil of flowerheads of garland chrysanthemum (*Chrysanthemum coronarium*) against agricultural pathogens. *Phytochemistry*, 57: 99-102.
- Ames BN (1983). Dietary Carcinogens and Anticarcinogens - Oxygen Radicals and Degenerative Diseases. *Science*, 221:1256-1264.
- Anthony KO (2006). Effects of Essential oils of some Meliaceae Plants on Aflatoxin Production and Growth of *Aspergillus parasiticus*. *J. Food Technol.* 4: 322-324.
- Atanda OO, Akpan I, Oluwafemi F (2007). The potential of some spice essential oils in the control of *A. parasiticus* CFR 223 and aflatoxin production. *Food Control*, 18: 601-607
- Bakkali F, Averbeck S, Averbeck D, Idaomar M (2008). Biological effects of essential oils- A review. *Food Chem. Toxicol.* 46: 446-475
- Barrauc L, Douki T, Guerret S, Chevallier M, Jamard C, Trepo C (2001). The role of duck hepatitis B virus and aflatoxin B<sub>1</sub> in the induction of oxidative stress in the liver. *Cancer Detect. Prev.* 25: 192-201.
- Baydar H, Sagdic O, Ozcan G, Karadogan G (2004) Antibacterial activity and composition of essential oils from Origanum, Thymbra and Satureja species with commercial importance in Turkey. *Food Control*, 15: 169-172.
- Bedard LL, Massey TM (2006). Aflatoxin B<sub>1</sub>-induced DNA damage and its repair. *Cancer Lett.* 241: 174-183
- Beric T, Nikolic B, Stanojevic J, Vukovic- Gacic B, Knez evic-Vukcevic J (2008). Protective effect of basil (*Ocimum basilicum* L.) against oxidative DNA damage and mutagenesis. *Food Chem. Toxicol.* 46: 724-732.
- Bhatnagar D, Ehrlich KC, Cleveland TE (1992). Oxidation-reduction reactions in biosynthesis of secondary metabolites. In *Biosynthesis of Secondary Metabolites Chapter 10*: pp. 255-286. Edited by Town Publishers.
- Bluma R, Amaiden MR, Daghero J, Etcheverry M (2007). Control of *Aspergillus* section Flavi growth and aflatoxin accumulation by plant essential oils. *J. Appl. Microbiol.* 1723: 1364-5072.
- Bluma R, Amaiden MR, Etcheverry M (2008). Screening of Argentine plant extracts: Impact on growth parameters and aflatoxin B<sub>1</sub> accumulation by *Aspergillus* section Flavi. *Int. J. Food Microbiol.* 122: 114-125.
- Bluma RV, Etcheverry MG (2008). Application of essential oils in maize grain: Impact on *Aspergillus* section Flavi growth parameters and aflatoxin accumulation. *Food Microbiol.* 25: 324-334
- Bullerman LB (1986). Mycotoxins and food safety. *Food Technol.* 40: 59-66.
- Burits M, Bucar F (2000). Antioxidant Activity of Nigella sativa Essential Oil. *Phytother. Res.* 14: 323-328
- Burt S (2004). Essential oils: their antimicrobial properties and potential applications in foods – a review. *Int. J. Food Microbiol.* 94: 223-253.
- Carmo ES, Lima EO, Souza EL, Sousa FB (2008). Effect of Cinnamomum zeylaicum blume essential oil on the growth and Morphogenesis of some potentially pathogenic *Aspergillus* species. *Braz. J. Microbiol.* 39: 91-97.
- Cheel J, Theoduloz C, Rodriguez J, Schmeda-Hirschmann G (2005). Free Radical Scavengers and Antioxidants from Lemongrass [*Cymbopogon citratus* (DC.) Stapf.]. *J. Agric. Food Chem.* 53: 2511-2517.
- Cheng KC, Cahill DS, Kasai H, Nishimura S, Loeb LA (1992). 8-Hydroxyguanine, an abundant form of oxidative DNA damage, causes G/T and A/C substitutions. *J. Biol. Chem.* 267: 166-172.
- Conner DE, Beuchat LR (1984a). Effects of essential oils from plants On growth of food spoilage yeast. *J. Food Sci.* 49: 429-434.
- Conner DE, Beuchat LR (1984b) Sensitivity of heat-stressed yeast to essential oils of plant. *Appl. Environ. Microbiol.* 47: 229-233.
- Cosentino S, Barra A, Pisano B, Cabizza M, Pirisi FM, Palmas F (2003). Composition and antimicrobial properties of Sardinian juniperus essential oils against foodborne pathogens and spoilage micro-

- organisms. *J. Food Protection*, 66: 1288-1291.
- Cotty PJ, Bayman P, Egel DS, Elias D (1994). Agriculture, aflatoxins, and *Aspergillus*. In *The Genus Aspergillus*, pp. 1-27. Edited by Powell KA, Fenwick A, Peberdy JF. New York: Plenum Press.
- Denissenko MF, Cahill J, Kondriakova TB, Gerber N, Pfeifer GP (1999). Quantitation and mapping of aflatoxin B<sub>1</sub>-induced DNA damage in genomic DNA using aflatoxin B<sub>1</sub>-8, 9-epoxide and microsomal activation systems. *Mutat. Res.* 425: 205-211.
- Dutton MF (1988). Enzymes and aflatoxin biosynthesis. *Microbiol. Rev.* 2: 274-295.
- Eaton DL, Gallagher EP (1994). Mechanisms of aflatoxin carcinogenesis. *Annu. Rev. Pharmacol. Toxicol.* 34: 135-172.
- Farag RS, Shalaby AS, El-Baroty GA, Ibrahim NA, Ali MA, Hassan EM (2004). Chemical and Biological Evaluation of the Essential Oils of Different *Melaleuca* Species. *Phytother. Res.* 18: 30-35.
- Halliwell B, Gutteridge JMC (1999). Oxidative stress: adaptation damage, repair and death, in: Halliwell B, Gutteridge JMC (Eds.), *Free Radicals in Biology and Medicine*, Oxford University Press, New York, pp. 246-350.
- Hashim S, Aboobaker VS, Madhubala R, Bhattacharya RK Rao AR (1994). Modulatory effects of essential oils from spices on the formation of DNA adducts by aflatoxin B<sub>1</sub> in vitro. *Nutr. Cancer*, 21: 169-175.
- Helal GA, Sarhan MM, Abu Shahla ANK, Abou El-Khair EK (2007). Effects of *Cymbopogon citratus* L. essential oil on the growth, morphogenesis and aflatoxin production of *Aspergillus flavus* ML2-strain. *J. Basic Microbiol.* 47: 5-15
- Hollstein M, Sidransky D, Vogelstein B, Harris CC (1991). *Sci.* 253: 49-53.
- Holmes RA, Boston RS, Payne GA (2008). Diverse inhibitors of aflatoxin biosynthesis. *Appl. Microbiol. Biotechnol.* 78: 559-572.
- Holmquist GU, Walker HH, Stahr HM (1983). Influence of temperature, pH, water activity and antifungal agents on growth of *Aspergillus flavus* and *A. parasiticus*. *J. Food Sci.* 48: 778-782.
- IARC (1985). Aflatoxin, IARC Monographs (Suppl. 7): 83-87.
- Kasai H (2002). Chemistry-based studies on oxidative DNA damage: formation, repair, and mutagenesis *Free Radic. Biol. Med.* 33: 450-456.
- Klein PJ, Van Vleet TR, Hall JO, Coulombe Jr, RA (2002). Biochemical factors underlying the age-related sensitivity of turkeys to aflatoxin B<sub>1</sub>. *Comp. Biochem. Physiol.* 132: 193-201.
- Kumar R, Dubey N, Tiwari OP, Tripathi YB, Sinha KK (2007). Evaluation of some essential oils as botanical fungitoxicants for the protection of stored food commodities from fungal infestation. *J. Sci. Food Agric.* 87: 1737-1742.
- Kumar R, Mishra AK, Dubey NK, Tripathi YB (2007). Evaluation of *Chenopodium ambrosioides* oil as a potential source of antifungal, anti-aflatoxigenic and antioxidant activity. *Int. J. Food Microbiol.* 115: 159-164.
- Minto RE, Townsend CA (1997). Enzymology and molecular biology of aflatoxin biosynthesis. *Chem. Rev.* 97: 2537-2555.
- Montes-Belmont R, Carvajal M (1998). Control of *Aspergillus flavus* in maize with plant essential oils and their components. *J. Food Prot.* 61(5): 616-619.
- Neguefact J, Leth V, Amvam Zollo PH, Mathur SB (2004). Evaluation of five essential oils from aromatic plants of Cameroon for controlling food spoilage and mycotoxin producing fungi. *Int. J. Food Microbiol.* 94: 329-334.
- Nyandieka HS, Wakhis J, Kilonzo MM (1990). Association of reduction of AFB<sub>1</sub>-induced liver tumours by antioxidants with increased activity of microsomal enzymes. *Indian J. Med. Res.* 92: 332-336.
- Nychas GJE (1995). Natural antimicrobial from plants. In *New Methods of Food Preservations* ed. Gould GW. pp. 58-89. Glasgow, UK: Blackie Academic and Professional.
- Oktaç M, Gulcin I, Kufrevioglu OI (2003). Determination of in vitro antioxidant activity of fennel (*Foeniculum vulgare*) seed extracts. *Lebensm.-Wiss.U.-Technol.* 36: 263-271.
- Paranagama PA, Abeysekera KHT, Abeywickrama K, Nugaliyadde L (2003). Fungicidal and anti-aflatoxigenic effects of the essential oil of *Cymbopogon citratus* (DC.) Stapf. (lemongrass) against *Aspergillus flavus* Link. isolated from stored rice. *Lett. Appl. Microbiol.* 37: 86-90.
- Pawar VC, Thaker VS (2007) In vitro efficacy of 75 essential oils against *Aspergillus niger*. *Mycoses*, 49: 316-323.
- Preston RJ, Williams GM (2005). DNA-reactive carcinogens: mode of action and human cancer hazard. *Crit. Rev. Toxicol.* 35: 673-683.
- Rasoli I, Abyaneh MR (2004). Inhibitory effects of Thyme oils on growth and aflatoxin production by *Aspergillus parasiticus*. *Food Control*, 15: 479-483.
- Rasoli I, Fakoor MH, Yadegarinia D, Gachkar L, Allameh A, Rezaei MB (2008). Antimycotoxigenic characteristics of *Rosmarinus officinalis* and *Trachyspermum copticum* L. essential oils. *Int. J. Food Microbiol.* 122: 135-139.
- Rasoli I, Owlia P (2005). Chemoprevention by thyme oils of *Aspergillus parasiticus* growth and aflatoxin production. *Phytochemistry*, 66: 285102856.
- Razzaghi-Abyaneh M, Shams-Ghahfarokhi M, Yoshinari T, Rezaee MB, Jaimand K, Nagasawa H, Sakuda S (2008). Inhibitory effects of *Satureja hortensis* L. essential oil on growth and aflatoxin production by *Aspergillus parasiticus*. *Int. J. Food Microbiol.* 123: 228-233.
- Sharma RA, Farmer PB (2004). Biological relevance of adduct detection to the chemoprevention of cancer. *Clin. Cancer Res.* 10: 4901-4912.
- Shenn LY, Wu CC, Lii CK, Tsai SJ (2001). Effect of diallyl sulfide and diallyl disulfide, the active principles of garlic, on the aflatoxin B<sub>1</sub>-induced DNA damage in primary rat hepatocytes. *Toxicol. Lett.* 122: 45-52.
- Shen H, Ong C, Lee B, Shi C (1995). Aflatoxin B<sub>1</sub>-induced 8-hydroxydeoxyguanosine formation in rat hepatic DNA. *Carcinog* 16: 419-422.
- Sohn DH, Kim YC, Oh SH, Park EJ, Li X, Lee BH (2003). Hepatoprotective and free radical scavenging effects of *Nelumbo nucifera*. *Phytomedicine*, 10: 165-169.
- Soliman KM, Badeaa RI (2002). Effect of oil extracted from some medicinal plants on different mycotoxigenic fungi. *Food Chem. Toxicol.* 40: 1669-1675.
- Souza EL, Lima EO, Freire KRL, Sousa CP (2005). Inhibitory Action of Some Essential Oils and Phytochemicals On the Growth of Various Moulds Isolated From Foods. *Braz. Arch. Biol. Technol.* 48(2): 245-250.
- Towner RA, Qian SY, Kadiiska MB, Mason RP (2003). In vivo identification of aflatoxin-induced free radicals in rat bile. *Free Radic. Biol. Med.* 35: 1330-1340.
- Trail F, Mahanti N, Linz J (1995). Molecular biology of aflatoxin biosynthesis. *Microbiology*, 141: 755-765.
- Valko M, Rhodes CJ, Moncola J, Izakovic M, Mazura M (2006). Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chemico Biol. Interact.* 160: 1-40.
- Wang JS, Groopman JD (1999). DNA damage by mycotoxins. *Mutat. Res.* 424: 167-181.
- Webster RP, Gawde MD, Bhattacharya RK (1996). Effect of different vitamin A status on carcinogen-induced DNA damage and repair enzymes in rats. *In Vivo* 10: 113-118.
- Wogan GN, Paglialonga S, Newberne PN (1974). Carcinogenic effects of low dietary levels of aflatoxin B<sub>1</sub> in rats. *Food Cosmet. Toxicol.* 12: 681-685.
- Wogan GN (1999). Aflatoxin as a human carcinogen. *Hepatology*, 30: 573-575.
- Yabe K, Nakajima H (2004). Enzyme reactions and genes in aflatoxin biosynthesis. *Appl. Microbiol. Biotechnol.* 64: 745-755.
- Yarborough A, Zhang YJ, Hsu TM, Santella RM (1996). Immunoperoxidase detection of 8-hydroxydeoxyguanosine in aflatoxin B<sub>1</sub>-treated rat liver and human oral mucosal cells. *Cancer Res.* 56: 683-688.
- Yu JJ, Chang PK, Ehrlich KC, Cary JW, Bhatnagar D, Cleveland TE, Payne GA, Linz JE, Woloshuk CP, Bennett JW (2004). Clustered pathway genes in aflatoxin biosynthesis. *Appl. Environ. Microbiol.* 70: 1253-1262.