

Review

***Agrobacterium*-induced hypersensitive necrotic reaction in plant cells: a resistance response against *Agrobacterium*-mediated DNA transfer**

Danladi Dada Kuta^{1*} and Leena Tripathi²

¹Biotechnology Advanced Laboratory, Sheda Science and Technology Complex (SHESTCO),
P.M.B. 186 Garki-Abuja, Nigeria

²Biotechnology Research Unit, International Institute of Tropical Agriculture (IITA), Nigeria
C/o L W Lambourn; Carolyn House; 26 Dingwall Rd; Croydon CR9 3EE; UK

Accepted 6 July, 2005

High necrosis and poor survival rate of target plant tissues are some of the major factors that affect the efficiency of *Agrobacterium*-mediated T-DNA transfer into plant cells. These factors may be the result of, or linked to, hypersensitive defense reaction in plants to *Agrobacterium* infection, which may involve the recognition of specific signals from the *Agrobacterium* that triggers the burst of reactive oxygen species at the infection site. Evidences of *Agrobacterium*-induced necrosis in target plant tissues and its link to reactive oxygen species are presented. Application of antioxidants, addition of acetosyringone and optimization of pre-culture conditions suppress the *Agrobacterium*-induced hypersensitive necrotic response in target plant tissues, thereby enhancing stable transformation.

Key words: *Agrobacterium*; hypersensitive reaction; necrosis; signal transduction; oxidative burst; transformation.

INTRODUCTION

Genetic transformation has become an important tool for crop improvement. At present time gene transfer by *Agrobacterium* is the established method of choice for the genetic transformation of most plant species. Compared to direct gene transfer methodologies (particle bombardment, electroporation, etc), *Agrobacterium*-mediated transformation offers several advantages such as the possibility to transfer only one or few copies of DNA fragments carrying the genes of interest at higher efficiencies with lower cost and the transfer of very large DNA fragments with minimal rearrangement (Hiei et al., 1997; Gheysen et al., 1998; Hansen and Wright, 1999;

Shibata and Liu, 2000). The most important advantage, however, is the possibility of producing transgenic plants, which are free of marker genes (Komari et al., 1996; Mathews et al., 2001). This has and will continue to have enormous implications with regards to approval by regulatory agencies, public acceptance and marketability of transgenic crops.

Recent advances in molecular biology of *Agrobacterium*-mediated transformation have improved our understanding of the mechanisms of recognition, induction of vir genes and transfer of T-DNA into plant cells by the *Agrobacterium* (Gustavo et al., 1998). However, the efficiency of *Agrobacterium*-mediated T-DNA transfer to plant cells depends not only on the successful recognition and colonization of plant cells by the *Agrobacterium*, but also on the responses of the plant cells to the *Agrobacterium* infection process

*Corresponding Author E-Mail: drkuta2002@yahoo.com.

(Zambryski, 1988; Binns, 1990). *Agrobacterium*-mediated transformation involves interaction between two biological systems and is affected by various physiological conditions (Bhalla and Smith, 1998).

Plant cells are known to possess the ability of recognizing invading pathogens and activating defense signal transduction leading to hypersensitive necrotic responses (Lamb et al., 1989; Mehdy, 1994; Dangl et al., 1996; Hammond-Kosack and Jones, 1996; Blumwald et al., 1998; Somssich and Hahlbrock, 1998; Richter and Ronald, 2000). The relationship of *Agrobacterium* to host plants is unique among plant pathogens. Many aspects of the plant-*Agrobacterium* interaction are not yet fully understood. It was earlier reported that *Agrobacterium* does not induce the hypersensitive response in target plants, even though the bacterium introduces several proteins into the host cell (Robinette and Matthyse, 1990). However, there are now several reports of high necrosis and poor survival rate of target plant tissues during the process of *Agrobacterium*-mediated T-DNA transfer (Pu and Goodman, 1992; Deng et al., 1995; Perl et al., 1996; Mercuri et al., 2000; Chakrabarty et al., 2002; Das et al., 2002). This could be the consequence of plant's hypersensitive reaction to *Agrobacterium* infection. Recently, it was demonstrated that plants can modulate their gene expression in response to *Agrobacterium* infection and that *Agrobacterium* can actually trigger the plant defense machinery (Ditt et al., 2001).

Hypersensitive reaction (HR) is known to be one of the plant defense responses and it is generally characterized by a rapid, localized cell death around the infection site and the accumulation of antimicrobial agents (Hammond-Kosack and Jones, 1996; Richter and Ronald, 2000). It is the sequence of events during HR that subsequently lead to necrosis of the collapsed cells (Goodman and Novacky, 1994). In this review article, the mechanisms by which the plant cells perceive and transduce signals from *Agrobacterium tumefaciens* to activate hypersensitive defense responses are suggested. The implications of *Agrobacterium*-induced plant defense responses for stable transformation are discussed and methods to suppress the defense responses proposed.

AGROBACTERIUM-INDUCES NECROSIS AND CELL DEATH IN INVADDED PLANT TISSUES

Plant tissue necrosis and cell death is reported to be one of the major factors that reduce the efficiency of *Agrobacterium*-mediated transformation (Gustavo et al., 1998), and is often observed in many crops. Pu and Goodman (1992) and Sangwan et al. (1992) were among the first investigators to report on *Agrobacterium*-induced necrosis in plant tissues. While Pu and Goodman (1992) observed *Agrobacterium*-induced necrosis in tissues of grape explants, Sangwan et al. (1992) reported that

Agrobacterium infection led to necrosis in target cells of *Arabidopsis thaliana*. The role of T-DNA genes in the induction of necrosis in host tissues was later demonstrated (Deng et al., 1995) in an experiment with grape plants. Perl et al. (1996), also experimenting with grape tissues, observed that co-cultivation with *Agrobacterium* resulted in host tissue necrosis and mortality. Interestingly, the necrotic response of the grape calli was observed not during co-cultivation, but 48 h after transfer of calli to *Agrobacterium*-free medium. These observations of *Agrobacterium*-induced necrosis in target plant tissues are gradually generating lots of interest among researchers. Recently, Hansen (2000) worked with maize tissues and observed that co-cultivation with *Agrobacterium* leads to rapid tissue necrosis and cell death. High tissue necrosis was also reported on leaf-discs of grape after co-cultivation with *Agrobacterium* (Das et al., 2002). In this case, the degree of necrotic reaction appears to depend on several transformation parameters, including explant age, pre-culture period, bacterial inoculum density, and infection duration. This confirms the earlier observations of Kumria et al. (2001) that high bacterial density ($A_{600} = 0.7 - 1.0$ with 10 min infection) or prolonged infection time (15-30 min with the optimal $A_{600} = 0.3 - 0.6$) adversely affect the growth and regeneration of callus of *Indica* rice during *Agrobacterium*-mediated transformation. Similarly, Chakrabarty et al. (2002) reported that exposure of cauliflower hypocotyl explants to undiluted culture of *Agrobacterium* ($OD_{600} = 0.5$) resulted in severe necrosis of the explants whereas diluted culture (1:10 and 1:20 dilution) reduced necrosis to greater extent. Also, the hypocotyls were hypersensitive to *Agrobacterium* infection when no pre-culture was allowed and necrotic reaction was enhanced on explants from 4-day-old seedlings in comparison to 7-day-old seedlings.

It appears, therefore, that exposure of plant tissues to *Agrobacterium* leads to tissue necrosis and cell death, which may invariably affect transformation efficiency. First, necrosis and cell death may occur in the cell layer where T-DNA is transferred. Transgenic cells that are imbedded in such necrotic tissues may be inhibited with regards to regeneration, thus reducing the recovery of transgenic cell clones (Potrykus, 1990). Necrotic tissues are also known to accumulate antimicrobial substances (Goodman and Novacky, 1994) that may inhibit the potential of *Agrobacterium* to colonize plant cells and transfer T-DNA. The active release of chemical signal, which induces the *vir* genes in *Agrobacterium*, occurs only in living, but not in dead necrotic cells (Shaw et al., 1991). Moreover, dead necrotic cells may also attract opportunistic microorganisms under *in vitro* conditions, leading to serious contamination that subsequently inhibits plant regeneration. Therefore, necrosis in host plant tissue during *Agrobacterium*-mediated T-DNA transfer drastically reduces transformation efficiency.

The optimization of *Agrobacterium*-mediated transfor-

systems may therefore require proper understanding of regulatory mechanisms of the *Agrobacterium*-induced necrotic reaction in plants.

PLANT TISSUE NECROSIS AND CELL DEATH AS DEFENSE MECHANISM AGAINST *AGROBACTERIUM*-MEDIATED GENE TRANSFER

Plant perception of signals from *Agrobacterium*

An efficient plant defense response usually requires the recognition of specific signal molecules from the invading pathogen (Blumwald et al., 1998; Richter and Ronald, 2000). The presence of chemical signaling between *Agrobacterium* and plant cell has been suggested earlier (Chilton, 1993). Though there are several reports on plant-excreted signals that induce *Agrobacterium* infection (Citovsky et al., 1982; Bolten et al., 1986; Cangelosi et al., 1990; Shaw et al., 1991), we have not come across any report of a concerted study on the signal molecules from the *Agrobacterium* that may elicit defense response in target plant tissues. However, there are reports on genotype-strain specificity during *Agrobacterium*-mediated transformation of plants (Owens and Cress, 1985; Byrne et al., 1987; Hobbs et al., 1989; Fillipone and Penza, 1992), which may indicate the presence of specific signals from specific *Agrobacterium* strain that could be recognized by specific plant genotype. Each *Agrobacterium*-susceptible plant cell (competent cell) has been shown to contain polysaccharide-polysaccharide binding sites recognizable by *Agrobacterium* (Sangwan et al., 1992). It has earlier been shown that the first step in the transfer of T-DNA molecule from *Agrobacterium* to plant is the recognition of a susceptible plant cell (Zambryski, 1988). Therefore, plant cell can be highly susceptible or non-susceptible to *Agrobacterium* infection, depending on the genotype of the host plant and the strain of the *Agrobacterium* (Jordan and Hobbs, 1994).

Non-susceptibility of plant cells to colonization by pathogen is known to be due to successful recognition of the invading pathogen by the plant cell, which generates an internal signal that triggers early defense responses in the plant cells (Somssich and Hahlbrock, 1998). This may also explain the non-susceptibility of plant cells of particular genotypes to infection by particular strain of *Agrobacterium* (Hobbs et al., 1989). The earliest defense reaction observed in non-susceptible plant cells following pathogen attack is oxidative burst (Mehdy, 1994).

***Agrobacterium*-induced oxidative burst in target plant cells**

Oxidative burst is the large and rapid generation of reactive oxygen species (superoxide, hydrogen peroxide,

hydroxyl, peroxy and alkoxy radicals), which can cause cell damage. It is now widely accepted that the key component of the oxidative burst is hydrogen peroxide. The well-known reactivity of hydrogen peroxide is not due to its reactivity per se, but requires the presence of a metal reductant to form the highly reactive hydroxyl radical, which is the strongest oxidizing agent known and reacts with organic molecules at diffusion-limited rates (Mehdy, 1994; McKersie, 1996). Numerous enzymes use hydrogen peroxide as a substrate in oxidation reactions, but the most prominent among the enzymes is peroxidase. Perl et al. (1996) observed that elevated levels of peroxidase activity in grape tissues correlated with *Agrobacterium*-induced necrosis in the host tissues during *Agrobacterium*-mediated transformation. Interestingly, the increase in peroxidase activity in the target grape tissue was always observed, not before or during co-cultivation with *Agrobacterium*, but 24-36 h after co-cultivation.

Peroxidase is known to mediate oxidative cross-linking of structural proteins in the cell wall (Somssich and Hahlbrock, 1998), and the *Agrobacterium*-induced increase in peroxidase activity in grape tissues could therefore confirm the role of oxidative burst in hypersensitive necrotic responses in plant to *Agrobacterium* infection. The reactive oxygen species (ROS) produced during pathogen-induced oxidative burst could be toxic enough to directly kill the attacking *Agrobacterium* (Wojtaszek, 1997). ROS can also lead to the induction of pathogenesis-related (PR) proteins (Mehdy, 1994), that may inhibit the potential of *Agrobacterium* to colonize and transfer T-DNA to plant cells. Transgene inactivation has been reported as a defense response of plants for expression of foreign DNA (Finnegan and McElroy, 1994), and ROS may play a role in such defense response. This is because the sugar and the base moieties of DNA are susceptible to oxidation by the hydroxyl radical, causing base degradation, single strand breakage, and cross-linking of DNA to protein (McKersie, 1996).

Therefore, it could be speculated that during incompatible plant-*Agrobacterium* interaction, the following sequence of events may occur in the target plant tissues: the first step is perception of specific signal(s) from the invading *Agrobacterium*, followed by the over-production of ROS (oxidative burst) at the site of *Agrobacterium* infection. Then the generated oxygen radicals may lead to plant cell death and necrosis, bacterial cell death, induction of pathogenesis-related genes, followed by the production of antimicrobial substances (phytoalexins, etc) and oxidation of sugar and base moieties of DNA. Therefore, proper understanding of these plant defense signal transduction events could assist in the development of strategies to suppress the *Agrobacterium*-induced defense responses and enhance the efficiency of *Agrobacterium*-mediated transformation, especially in 'recalcitrant' species.

CONTROLLING PLANT DEFENSE RESPONSES AGAINST *AGROBACTERIUM*-MEDIATED T-DNA TRANSFER

There are reports on successful experiments for optimization of transformation protocols. The strategies employed in such experiments can be grouped into two: 1. Quenching of the *Agrobacterium*-induced oxidative burst; and 2. Reprogramming of an *Agrobacterium*-incompetent plant cell into a competent one.

Quenching of *Agrobacterium*-induced oxidative burst

The activity of oxidative burst in plant defense responses could be suppressed by the addition of antioxidants such as ascorbic acid, cysteine, citric acid, PVPP, PVP, DTT, and cyclitols (myo-inositol). Some of these compounds are known to scavenge reactive oxygen species, thereby quenching oxidative burst. The application of a mixture of antioxidants has been shown to improve the efficiency of *Agrobacterium*-mediated transformation in some crops (Perl et al., 1996; Das et al., 2002). The combination of PVPP and DTT was found by Perl et al. (1996) to improve the viability of embryogenic grape calli. They observed that tissue necrosis was completely inhibited by these antioxidants while *Agrobacterium* virulence was not affected. In their review of *Agrobacterium*-mediated transformation of plants, Gustavo et al. (1998) reported that efficient transformation of monocotyledonous crops like sugarcane was possible only when a mixture of antinecrotic compounds with remarkable antioxidative activity was added. Recently, Das et al. (2002) applied the double-layer antioxidant method as described by Perl et al. (1996) to control the problem of *Agrobacterium*-induced necrosis during transformation of grape leaf-discs. Therefore, compounds with potential to quench oxidative burst could be used to arrest *Agrobacterium*-induced necrosis in host tissues, thereby improving transformation efficiency.

Reprogramming a resistant plant cell into a susceptible one

To make plant tissues susceptible to *Agrobacterium* infection, they have to be induced to undergo cellular dedifferentiation (Sangwan et al., 1992). Pathogen recognition may be weakened in dedifferentiating cells probably due to perturbation of membrane structure. Sangwan et al. (1992) explained that pre-culture of explants prior to *Agrobacterium* infection on media containing auxin with or without cytokinins is a good method of inducing cells to undergo dedifferentiation and may serve as rejuvenating treatment to the explant. Juvenile plant cells may be more susceptible to

Agrobacterium infection than differentiated old cells. Pre-culture treatment has recently been shown to improve the efficiency of *Agrobacterium*-mediated transformation (Chakrabarty et al., 2002; Wu et al., 2003).

Another approach of reprogramming plant cell development is wounding. Wounding is the most effective biological trigger for shifting cells potentially competent for regeneration to the competent state (Potrykus, 1990). Wound healing has earlier been shown to trigger a sequence of reactions at the cellular level that is important for T-DNA-induced transformation (Binns and Thomashow, 1988). The ability of wounded plant cells to enter and carry out one or more cell cycles may be absolutely required for successful transformation (Binns, 1990; Sangwan et al. 1992). Apart from stimulating dedifferentiation, wounding also leads to the excretion of chemical signal that induces *Agrobacterium* infection (Citovsky et al., 1982; Kahl, 1982; Bolten et al., 1986; Potrykus, 1990; Shaw et al., 1991). The wound-exuded chemical signals are phenolic compounds, like acetosyringone. However, monocots do not show the wound response characteristic of the dicot species (Binns, 1990). Moreover, Escudero and Hohn (1997) demonstrated that the competence of plant cells for *Agrobacterium*-mediated DNA transfer is not necessarily linked to wounding. In this case, exogenous acetosyringone, added in the inoculation and co-cultivation media, replaced the need for wounding.

Apart from inducing vir genes in the *Agrobacterium* (Cangelosi et al., 1990), it is possible that acetosyringone also perturb the *Agrobacterium*-induced defense signal transduction events in plant cells, leading to reprogramming of *Agrobacterium*-incompetent cell to a competent one. In several experiments with monocots and other 'recalcitrant' species, addition of acetosyringone to the inoculation and co-cultivation media improved the efficiency of *Agrobacterium*-mediated transformation (May et al., 1995; Kumria et al., 2001; Mahmoudian et al., 2002; Wu et al., 2003).

It appears, therefore, that with proper optimization of transformation parameters, like duration of pre-culture and addition of acetosyringone (Wu et al., 2003), plant cells of any species could be made competent for *Agrobacterium*-mediated transformation. It was, for example, previously suggested that cereals cannot be transformed by *Agrobacterium* (Potrykus, 1990), but with proper manipulation of transformation parameters, large number of fertile transgenic rice were produced through *Agrobacterium*-mediated DNA transfer system (Hiei et al., 1994; Rashid et al., 1996).

CONCLUSION

This review shows that *Agrobacterium*-induced necrosis often observed in target plant tissues is linked to

hypersensitive defense reaction in plants to *Agrobacterium* infection. The plant defense mechanisms against *Agrobacterium* involve successful recognition of some sort of signals from the *Agrobacterium* which triggers oxidative burst at the infection site. The HR-inducing factor in *Agrobacterium* is yet to be fully understood. However, Zheng et al. (2003) have recently discovered a gene in *Agrobacterium vitis* that is associated with *Agrobacterium*-induced HR. This gene, *aviR*, is said to be homologous to *luxR*, which implies that the *Agrobacterium*-induced HR is regulated by a quorum-sensing mechanism. The *Agrobacterium*-induced HR could lead to rapid and large generation of reactive oxygen radicals in target plant cells, resulting to plant cell death (necrosis), oxidative stress to the invading *Agrobacterium* cells, production of toxic antibacterial substances and the deleterious effects on DNA molecules, especially at the site of oxidative burst. All these factors significantly reduce the efficiency of stable transformation of plants. Therefore, detoxification of the ROS with carefully selected mixture of compounds with high antioxidative activity, and, reprogramming a non-susceptible plant cell into an *Agrobacterium*-competent cell by wounding, addition of acetosyringone and optimization of pre-culture conditions, are possible methods for improving the efficiency of *Agrobacterium*-mediated transformation, especially in 'recalcitrant' crops.

ACKNOWLEDGEMENT

The first author is grateful to USDA and IITA for awarding him a Visiting Fellow position at IITA-IBADAN, which made this review possible.

REFERENCES

- Bhalla PL, Smith N (1998). *Agrobacterium tumefaciens*-mediated transformation of cauliflower, Brassica oleracea var. botrytis. Mol. Breed 4: 531–541.
- Binns AN (1990). *Agrobacterium*-mediated gene delivery and the biology of host range limitations. Physiol. Plant 79: 135–139.
- Binns AN, Thomashow M (1988). Cell Biology of *Agrobacterium* infection and transformation of plants. Ann. Rev. Microbiol. 42: 575–606.
- Blumwald E, Aharon GS, Lam BCH (1998). Early signal transduction pathways in plant–pathogen interactions. Trends in Plant Sci. 3: 342–346.
- Bolten GW, Nester EW, Gordon MP (1986). Plant phenolic compounds induce expression of *Agrobacterium tumefaciens* loci needed for virulence. Science 232: 981–985.
- Cangelosi GA, Ankenbauer RA, Nester EW (1990). Sugars induce the *Agrobacterium* virulence genes through a periplasmic binding protein and a transmembrane signal protein. Proceedings of the National Academy of Sciences 87: 6708–6712.
- Chakrabarty R, Viswakarma N, Bhat SR, Kirti PB, Singh BD, Chopra VL (2002). *Agrobacterium*-mediated transformation of cauliflower: optimization of protocol and development of Bt-transgenic cauliflower. J. Biosci. 27: 495–502.
- Chilton MD (1993). *Agrobacterium* gene transfer: progress on a “poor man’s vector” for maize. Proc. Natl. Acad. Sci. USA 90, 3119–3120.
- Citovsky V, McLean BG, Greene E, Howard E, Kulsau G, Thornstenson Y, Zupan J, Zambryski, PC (1982). *Agrobacterium*-plant cell interaction: induction of vir genes and T-DNA transfer. In: Molecular signals in Plant-microbe communications (Verma, DPS, ed.), Boca Raton, FL: CRC Press. pp.169–190.
- Dangl JL, Dietrich RA, Richberg MH (1996). Death don't have no mercy: Cell death programs in plant–microbe interactions. Plant Cell 8: 1793–1807.
- Das DK, Reddy MK, Upadhyaya KC, Sopory SK (2002). An efficient leaf-disk culture method for the regeneration via somatic embryogenesis and transformation of grape (*Vitis vinifera* L.). Plant Cell Rep. 20: 999–1005.
- Deng W, Pu X-A, Goodman RN, Gordon MP, Nester EW (1995). T-DNA genes responsible for inducing a necrotic response on grape vines. Mol. Plant-Microbe Interact. 8: 538–548.
- Ditt R, Nester EW, Comai L (2001). The plant gene expression response to *Agrobacterium tumefaciens*. Proc. Natl. Acad. Sci. USA. 98: 10954–10954.
- Filippone E, Penza R (1992). *Agrobacterium*-mediated gene transfer. In Biotechnology: Enhancing Research on Tropical Crops in Africa (Thottappilly G, Monti L, Mohan Raj DR, Moore AW, eds.), CTA/IITA co-publication. IITA, Ibadan, pp.197–202.
- Escudero J, Hohn B (1997). Transfer and integration of T-DNA without cell injury in the host. The Plant Cell 9: 2135–2142.
- Finnegan J, McElroy D (1994). Transgene inactivation: plants fight back! Biotechnology 12: 883–888.
- Ganapathi TR, Higgs NS, Balint-Kurti PJ, Arntzen CJ, May GD, van Eck JM (2001). *Agrobacterium*-mediated transformation of embryogenic cell suspensions of the banana cultivar Rasthali (AAB). Plant Cell Rep. 20: 157–162.
- Gheysen G, Angenon G, Van Montague M (1998). *Agrobacterium*-mediated plant transformation: A scientifically intriguing story with significant application. In: K Lindsey (ed) Transgenic plant research. Harwood Academic Press, The Netherlands. pp.1–33
- Goodman RN, Novacky AJ (1994). The hypersensitive reaction in plants to pathogens. A resistant phenomenon. APS PRESS, St. Paul, Minnesota, p. 244.
- Grimsley NH (1990). Agroinfection. Physiol. Plant 79: 147–153.
- Gustavo AR, Gonzalez-Cabrera J, Vazquez-Padron R, Ayra-Pardo C (1998). *Agrobacterium tumefaciens*: A natural tool for plant transformation. Electronic J. Biotechnol. V.1 <http://www.ejb.org/content/vol1/issue3/full/1>.
- Hammond-Kosack KE, Jones JDG (1996) Resistance gene–dependent plant defense responses. Plant Cell 8: 1773–1791.
- Hansen G (2000). Evidence for *Agrobacterium*-induced apoptosis in maize cells. Mol. Plant-Microbe Interact. 13: 649–657.
- Hansen G, Wright MS (1999). Recent advances in transformation of plants. Trends Plant Sci. 4: 226–231.
- Hiei Y, Komari T, Kubo T (1997). Transformation of rice mediated by *Agrobacterium tumefaciens*. Plant Mol. Biol. 35: 205–218.
- Hiei Y, Ohta S, Komari T, Kumashiro T (1994). Efficient transformation of rice (*Oryza sativa* L.) mediated by *Agrobacterium* and sequence analysis of the boundaries of the T-DNA. Plant J. 6: 271–282.
- Jordan MC, Hobbs SLA (1994). The transformation of legumes using *Agrobacterium tumefaciens*. In Biotechnological applications of plant cultures (Shargool, PD, Ngo TT, eds.), CRC press, Boca Raton. pp. 61–76.
- Kahl G (1982). Molecular Biology of Wound Healing: The conditioning phenomenon. In Molecular biology of plant tumors (Kahl, G., and Schell, J., eds.), New York, pp. 211–268.
- Kamoun S (2003). Agrosuppression: A bioassay for the hypersensitive response suited to high-throughput screening. Mol. Plant-Microbe Interact. 16: 7–13.
- Komari T, Hiei Y, Saito Y, Murai N, Kumashiro T (1996). Vectors carrying two different T-DNAs for co-transformation of higher plants mediated by *Agrobacterium tumefaciens* and segregation of transformants free from selection markers. Plant J. 10: 165–174.
- Kumria R, Waie B, Rajam MV (2001). Plant regeneration from transformed embryogenic callus of an elite Indica rice via *Agrobacterium*. Plant Cell Tissue and Organ Cult. 67: 63–71.

- Lam E, Pontier D, del Pozo O (1999). Die and let live - programmed cell death in plants. *Current Opin. Plant Biol.* 2: 502-507.
- Lamb CJ, Lawton MA, Dron M, Dixon, RA (1989). Signals and transduction mechanisms for activation of plant defenses against microbial attack. *Cell* 56: 215—224.
- Mahmoudian M, Yucel M, Oktem HA (2002). Transformation of Lentil (*Lens culinaris* M.) cotyledonary nodes by vacuum infiltration of *Agrobacterium tumefaciens*. *Plant Mol. Biol. Rep.* 20, 251--257.
- Matthews PR, Wang MB, Waterhouse PM, Thornton S, Fieg SJ, Gubler F, Jacobsen JV (2001). Marker gene elimination from transgenic barley, using co-transformation with adjacent 'twin T-DNAs' on standard *Agrobacterium* transformation vector. *Mol Breed* 7: 195—202.
- May GD, Afza R, Mason HS, Wiecko AF, Novak J, Arntzen CJ (1995). Generation of transgenic banana (*Musa acuminata*) via *Agrobacterium*-mediated transformation. *Biotechnology* 13, 486--492.
- McKersie B.D. 1996 Oxidative stress. <http://www.agronomy.psu.edu/Courses/AGRO518/Oxygen.htm>.
- Mehdy MC (1994). Active oxygen species in plant defense against pathogens. *Plant Physiol.* 105, 467—472.
- Mercuri A, Benedetti LD, Burchi G, Schiva T (2000). *Agrobacterium*-mediated transformation of African violet. *Plant Cell Tissue and Organ culture* 60: 39—46.
- Owens LD, Cress DE (1985). Genotypic variability of soybean response to *Agrobacterium* strains harboring Ti or Ri plasmids. *Plant Physiol.* 77: 87.
- Perl A, Lotan O, Abu-Abied M, Holland D (1996). Establishment of an *Agrobacterium*-mediated transformation system for grape (*Vitis vinifera* L.): The role of antioxidants during grape-*Agrobacterium* interactions. *Natur Biotechnol.* 14: 624-628.
- Potrykus I (1990). Gene transfer to cereals: an assessment. *Bio/Technology* 8: 535—542.
- Pu X-A, Goodman RN (1992) Induction of necrosis by *Agrobacterium tumefaciens* on grape explants. *Physiol. Mol. Plant Pathol.* 41: 245--254.
- Rashid H, Yokoi S, Toriyama K, Hineta K (1996). Transgenic plant production mediated by *Agrobacterium* in indica rice. *Plant Cell Rep.* 15: 727--773.
- Richter TE, Ronald PC (2000). The evolution of disease resistance genes. *Plant Mol. Biol.* 42: 195--204.
- Robinette D, Matthyse AG (1990). Inhibition by *Agrobacterium tumefaciens* and *Pseudomonas savastanoi* of development of the hypersensitive response elicited by *Pseudomonas syringae* pv. phaseolicola. *J. Bacteriol.* 172: 5742- 5749.
- Sangwan RS, Bourgeois Y, Brown S, Yasseur G, Sangwan-Norreel B (1992). Characterization of competent cells and early events of *Agrobacterium*-mediated genetic transformation in *Arabidopsis thaliana*. *Planta* 188: 439—456.
- Shaw CH, Loake GJ, Brown AP, Garrett CS (1991). The early events in *Agrobacterium* infection. In *Biochemistry and Molecular Biology of Plant-pathogen interactions* (Smith CJ, ed.), Clarendon press, Oxford. pp.197—209.
- Shibata D, Liu YG (2000). *Agrobacterium*-mediated plant transformation with large DNA fragments. *Trends Plant Sci.* 5: 354—357.
- Somssich IE, Hahlbrock K (1998). Pathogen defense in plants – a paradigm of biological complexity. *Trends in Plant Science* 3, 86--90.
- Wojtaszek P (1997). Oxidative burst: an early plant response to pathogen infection. *Biochem. J.* 322: 681 - 691.
- Wu H, Sparks C, Amoah B, Jones HD (2003). Factors influencing successful *Agrobacterium*-mediated genetic transformation of wheat. *Plant Cell Rep.* 21: 659--668.
- Zambryski P (1988). Basic processes underlying *Agrobacterium*-mediated DNA transfer to plant cells. *Annu. Rev. Genet.* 22:1-30.
- Zheng D, Zhang H, Carle S, Hao G, Holden MR, Burr TJ (2003). A luxR homolog, aviR, in *Agrobacterium vitis* is associated with induction of necrosis on grape and a hypersensitive response on tobacco. *Mol. Plant-Microbe Interact.* 16: 650--658.