

Full Length Research Paper

# Metabolic engineering and thermodynamic characterization of an extracellular $\beta$ -glucosidase produced by *Aspergillus niger*

Sana Zahoor<sup>1</sup>, Muhammad Mohsin Javed<sup>1\*</sup>, Saima Aftab<sup>1</sup>, Farooq Latif<sup>2</sup> and Ikram-ul-Haq<sup>1</sup>

<sup>1</sup>Institute of Industrial Biotechnology, GC University, Lahore, Pakistan.

<sup>2</sup>National Institute of Biotechnology and Genetic Engineering (NIBGE), Faisalabad, Pakistan.

Accepted 3 June, 2011

The production of an extracellular  $\beta$ -glucosidase by *Aspergillus niger* NRRL 599 was optimized using submerged fermentation technique. Effect of different media, different carbon sources, initial pH of the fermentation medium, temperature, incubation period and inoculum size on the production of  $\beta$ -glucosidase enzyme was investigated. *A. niger* NRRL 599 produced maximum extracellular  $\beta$ -glucosidase (4.48 U/mg) in Eggins and Pugh medium with 1% wheat bran (w/v) at pH 5.5 inoculated with 4% conidial suspension after 96 h of incubation at 30°C. Purified  $\beta$ -glucosidase gave  $K_m$  and  $V_{max}$  values of 3.11 mM and 20.83 U/mg respectively for pNPG hydrolysis. The enzyme was optimally active at pH 4.8 and at temperature of 60°C. Thermodynamic parameters,  $E_a$ ,  $\Delta H$  and  $\Delta S$  were found to be 52.17 KJ/mol, 49.90 J/mol.K and -71.69 KJ/mol, respectively. The  $pK_{a1}$  and  $pK_{a2}$  of ionizable groups of active site residues involved in  $V_{max}$  were calculated to be 4.1 and 6.0 respectively.

**Key words:**  $\beta$ -Glucosidase, *Aspergillus niger*, kinetics, thermodynamics.

## INTRODUCTION

$\beta$ -Glucosidase (EC 3.2.1.21) catalyzes the hydrolysis of  $\beta$ -1,4 glucosidic bonds in disaccharides, oligosaccharides, alkyl and aryl  $\beta$ -D-glucosides by attacking on the non-reducing terminal and releases  $\beta$ -D glucose (Iwashita et al., 1999; Wallecha and Mishra, 2003).  $\beta$ -Glucosidase is produced by humans, insects, plants, bacteria, fungi and yeast (Villena et al., 2007). This enzyme has been exploited in a variety of biotechnological and biochemical industries. Moreover,  $\beta$ -glucosidase component plays a pivotal role in the final step of cellulose degradation (Seidle and Huber, 2005).  $\beta$ -Glucosidase is generally

responsible for the regulation of the whole cellulolytic process and is a rate-limiting factor during enzymatic hydrolysis of cellulose, as both endoglucanase and exoglucanase activities are often inhibited by cellobiose. Thus,  $\beta$ -glucosidase not only produces glucose from cellobiose, but also reduces cellobiose inhibition, allowing endoglucanase and exoglucanase enzymes to function more efficiently (Kaur et al., 2007). Supplementing  $\beta$ -glucosidase in commercial cellulase preparations has been reported to result in increased rate of cellulose hydrolysis and ethanol production (Wallecha and Mishra, 2003).

*Aspergillus niger* is by far the most efficient producer of  $\beta$ -glucosidase among the microorganisms investigated. It is generally recognized as safe by the Food and Drug Administration (Tailor and Richardson, 1979).  $\beta$ -Glucosidase can be produced both by solid-state and submerged fermentation techniques (Christakopoulos et al., 1995; lembo et al., 2002; Fawzi, 2003; Garcia-Kirchner et al., 2005; Daroit et al., 2007; Leite et al., 2007). Sub-

\*Corresponding author. E-mail: mmj\_bot@yahoo.com. Tel: +92 42 99211634. Fax: +92 42 99211634.

**Abbreviations:** CMC, Carboxymethylcellulose; FPLC, fast protein liquid chromatography; pNPG, *p*-nitrophenyl- $\beta$ -D-glucopyranoside.

merged fermentation offers several advantages, among them a possibility to automate many operations, the ease with which various parameters can be monitored (by periodic sampling of broth) and controlled if necessary by the addition of further nutrients or reagents and easier removal of heat generated during the microbial growth in a large scale reactor (Jakubikova et al., 2006). Extra-cellular production of  $\beta$ -glucosidase using submerged fermentation technique had been investigated by different researchers. Gunata and Vallier (1999) studied the production of extracellular  $\beta$ -glucosidase from *A. niger* and *Aspergillus oryzae* using submerged fermentation technique. Jager et al. (2001) investigated the extra-cellular production of  $\beta$ -glucosidase from different *Aspergillus* strains using submerged fermentation and Garcia-Kirchner et al. (2005) studied the effect of media composition and growth conditions on the production of  $\beta$ -glucosidase by *A. niger* C-6 under submerged culture.

This study was initiated with the aim to exploit *A. niger* NRRL 599 for maximum yield of  $\beta$ -glucosidase and the subsequent characterization of  $\beta$ -glucosidase.

## MATERIALS AND METHODS

### Organism

*A. niger* NRRL 599 was obtained from the stock culture of Institute of Industrial Biotechnology, GC University, Lahore. The culture was maintained on 4% potato dextrose agar medium (pH: 5.5). The slants were incubated at 30°C for maximum growth and stored at 4°C. Sub-culturing was carried out every 15 days.

### Inoculum

The conidial suspension was used as an inoculum. The number of conidia was counted with the help of haemocytometer slide bridge (Neubauer improved, Precidder HBG, Germany). Ten milliliters of sterilized distilled water was transferred to each slant having profused conidial growth on its surface. The sterilized inoculating needle was used to break the clumps of conidia. The tube was shaken vigorously to make a homogenous mixture of conidial suspension.

### Fermentation

$\beta$ -Glucosidase fermentation was carried out using submerged technique in 250 ml Erlenmeyer flasks containing 25 ml of Eggins and Pugh (1962) medium consisting of:  $(\text{NH}_4)_2\text{SO}_4$  (0.5 g/l), KCl (1.0 g/l),  $\text{KH}_2\text{PO}_4$  (0.5 g/l),  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  (0.2 g/l), L-asparagine (0.5 g/l),  $\text{CaCl}_2$  (0.1 g/l), yeast extract (0.5 g/l) with 1% carbon source (w/v) at pH 5.5 inoculated with 4% conidial suspension. The flasks were incubated in an orbital shaking incubator (SANYO, Gallenkamp, PLC, UK) at 30°C (200 rpm) for 96 h. After the incubation, the fermented broth was centrifuged at 6,000 rpm for 10 min and the supernatant was used for analytical purposes.

### Optimization of fermentation parameters

Eight different types of fermentation media, (M-I to M-VIII) already reported by Eggins and Pugh (1962), Mandel and Reese (1960), Romanelli et al. (1975), Coutts and Smith (1976), Saddler (1982), Macris and Panayatou (1989), Almin et al. (1975) and Riou et al. (1998) respectively, were screened for the production of  $\beta$ -glucosidase. Different carbon sources, for example, carboxymethyl cellulose (CMC), cellulose powder, lactose, maltose, sucrose, wheat bran and cellobiose (1%, w/v) were added to fermentation medium to test their effect on enzyme production. Time course study for the production of  $\beta$ -glucosidase was carried up to 144 h. Effect of different incubation temperature (25 to 35°C), initial pH of the fermentation medium (4.0 to 7.0) and conidial inoculum (1 to 5%) was also investigated on the production of  $\beta$ -glucosidase. All the experiments were carried out in triplicate.

### Analysis

$\beta$ -Glucosidase activity was estimated according to the method used by Rajoka and Malik (1997). "One unit of  $\beta$ -glucosidase activity is defined as the amount of enzyme required to release 1.0  $\mu\text{mol}$  of p-nitrophenol per min under the assay conditions". Total protein was estimated according to the method of Bradford (1976) using bovine serum albumin as standard.

### Purification of $\beta$ -glucosidase


Crude  $\beta$ -glucosidase was purified to homogeneity by using ammonium sulfate precipitation and fast protein liquid chromatography (FPLC) by using RESOURCE S (AMERSHAM BIOSCIENCES, USA). Solid ammonium sulfate was slowly added to crude enzyme to 80% saturation at 0°C to precipitate the  $\beta$ -glucosidase with stirring at an interval of 5 min. The pellet was collected by centrifugation at 12000  $\times g$  for 20 min (4°C) to recover precipitated proteins. The pellet was dissolved in 3.0 ml of phosphate buffer (pH: 7.2) and dialyzed against 0.05 M sodium phosphate buffer (pH: 7.5) by constant stirring for 24 h and periodic change of buffer. Dialysis membrane with a molecular weight cut-off of 12000 to 14000 was used. The dialyzed sample, filtered through 0.4  $\mu\text{m}$  Millipore filter, was loaded on an anion exchange column RESOURCE-S (6.0 ml, 16  $\times$  30 mm; AMERSHAM BIOSCIENCES, USA) equilibrated with 50 mM sodium phosphate buffer (pH: 7.5) with a flow rate of 1.0 ml/min. The protein was eluted with a linear gradient of NaCl (0 to 1 M) in the buffer. The elution profile was examined at 280 nm. The fractions were collected and assayed for enzyme activity.

### Enzyme characterization

The purified enzyme was characterized as a function of temperature and pH. The optimal temperature for enzyme activity was determined over a temperature range of 20 to 65°C using the standard assay conditions. To determine the optimum pH for the enzyme,  $\beta$ -glucosidase activity was measured over a pH range of 3.0 to 7.0 using the standard assay conditions.

### Kinetic and thermodynamic studies

The kinetic constants, Michaelis constant ( $K_m$ ), and maximum



**Figure 1.** Screening of fermentation media for  $\beta$ -glucosidase production by *Aspergillus niger* NRRL 599 in shake flasks.  $\pm$  indicates the standard deviation among the three parallel replicates. Incubation at 30°C for 96 h at pH 5.5 using wheat bran as carbon source.

velocity ( $V_{max}$ ), were evaluated by determining the  $\beta$ -glucosidase activity at 60°C against different *p*-nitrophenyl- $\beta$ -D-glucopyranoside (pNPG) concentrations ranging from 1-15mM and keeping the enzyme concentration constant.  $K_m$  and  $V_{max}$  were calculated from Lineweaver-Burk double reciprocal plot using Equation 1 (Lineweaver and Burk, 1934):

$$1/v = (1/V_{max}) + (K_m/V_{max}) \cdot 1/[S] \quad (1)$$

Thermodynamic constants, activation energy ( $E_a$ ), enthalpy of activation ( $\Delta H$ ) and entropy of activation ( $\Delta S$ ), were calculated from Arrhenius plot using Equations 2 to 4 after Siddiqui et al. (1997a):

$$E_a = -\text{slope} (R) \quad (2)$$

$$\Delta H = E_a - RT$$

$$\ln(V_{max}/T) = \ln(K_B/h) + \Delta S/R - (\Delta H/R)1/T \quad (3)$$

where slope =  $\Delta H/R$  and intercept =  $\ln(K_B/h) + \Delta S/R$

$$\Delta S = R [\text{intercept} - \ln(K_B/h)] \quad (4)$$

Where,  $T$ ,  $K_B$ ,  $h$ ,  $R$ ,  $E_a$ ,  $\Delta H$  and  $\Delta S$  are absolute temperature (K), Boltzmann constant ( $1.38 \times 10^{-23}$  J/K), Planck's constant ( $6.626 \times 10^{-34}$  Js), Gas constant (8.314 J/K.mol), activation energy, enthalpy of activation and entropy of activation, respectively.  $pK_{a1}$  and  $pK_{a2}$  values of ionizable groups of active site residues were


calculated using Dixon plot (Dixon and Webb, 1979) (Figure 10).

### Statistical analysis


Treatment effects were compared after Snedecor and Cochran (1980) using computer software Costat, cs6204W.exe. Significant difference among replicates has been presented as Duncan's multiple range tests in the form of probability ( $p$ ) values.

## RESULTS AND DISCUSSION

The composition of fermentation media has a marked influence on enzyme production. Eight different type of fermentation media were tested for the production of  $\beta$ -glucosidase. M-I medium reported by Eggins and Pugh (1962) was found to be the best medium for the production of  $\beta$ -glucosidase (Figure 1). In M-I, the presence of L-asparagine and  $(NH_4)_2SO_4$  in addition to other essential components might have resulted in the increased production of  $\beta$ -glucosidase. Other media gave relatively less amount of enzyme as they might be lacking in any of the components essential for the growth of fungi and enzyme production. Grajek (1987) also reported  $(NH_4)_2SO_4$  as the best nitrogen source for the production



**Figure 2.** Effect of different carbon sources on  $\beta$ -glucosidase production by *Aspergillus niger* NRRL 599 in shake flasks.  $\pm$  indicates the standard deviation among the three parallel replicates. Incubation at 30°C for 96 h at pH 5.5 in M-I medium.




**Figure 3.** Effect of incubation time on  $\beta$ -glucosidase production by *Aspergillus niger* NRRL 599 in shake flasks.  $\pm$  indicates the standard deviation among the three parallel replicates. Incubation at 30°C and pH 5.5 using wheat bran as carbon source in M-I medium.

of  $\beta$ -glucosidase from *Sporotrichum (Chrysosporium) thermophile*.

$\beta$ -Glucosidase is an inducible enzyme (Busto et al., 1995; Daroit et al., 2007). In order to get maximum production, wheat bran used in Eggins and Pugh medium was replaced with different carbon sources but no positive change was observed with other carbon sources (Figure 2). It is reported that cellobiose as a carbon source is the best for the induction of  $\beta$ -glucosidase (Busto et al., 1995) but in this study CMC, cellulose powder, lactose, maltose, sucrose and cellobiose gave less  $\beta$ -glucosidase production than wheat bran. Wheat bran

contain plenty of cello-oligosaccharides that results in the increased activity of extracellular  $\beta$ -glucosidase (Sun et al., 2008). Jager et al. (2001) and Rajoka et al. (2006) found wheat bran as the best substrate for the production of  $\beta$ -glucosidase from different *Aspergillus* strains using solid state fermentation.

The optimization of time course for the production of  $\beta$ -glucosidase by *Aspergillus niger* NRRL 599 was investigated up to 144 h after inoculation. Maximum  $\beta$ -glucosidase was obtained after 96 h of fermentation (Figure 3). However, further increase in the incubation time reduced the enzyme production. It might be due to



**Figure 4.** Effect of incubation temperature on  $\beta$ -glucosidase production by *Aspergillus niger* NRRL 599 in shake flasks.  $\pm$  indicates the standard deviation among the three parallel replicates. Incubation at pH 5.5 for 96 h using wheat bran as carbon source in M-I medium.


the depletion of macro- and micronutrients in the fermentation medium with the lapse in time, which stressed the fungal physiology resulting in the inactivation of secretory machinery of the enzymes. In addition, the substances were initially more susceptible, making a rapid rise in enzymes biosynthesis. But with the prolongation of cultural time, the susceptible portions were completely hydrolyzed by microorganisms (Haq et al., 2006). Tsao et al. (2000) also obtained maximum  $\beta$ -glucosidase production from *A. niger* after 96 h of fermentation while Garcia-Kirchner et al. (2005) obtained maximum  $\beta$ -glucosidase titer from *A. niger* within 72 h of fermentation.

Effect of different incubation temperature (25 to 35°C) on the production of  $\beta$ -glucosidase was evaluated. Maximum  $\beta$ -glucosidase was obtained at 30°C (Figure 4). Further increase or decrease in temperature resulted in gradual decrease in enzyme production. At lower temperature, the transport of nutrients in the cells is hindered and lower product yields are attained, while at high temperature, the maintenance energy requirement increases owing to inactivation of proteins of the metabolic pathway and less product formation occurs (Rajoka et al., 2004). Gunata and Vallier (1999) also obtained maximum production of  $\beta$ -glucosidase from *A. niger* and *Aspergillus oryzae* at incubation temperature of 30°C.


The initial pH of the medium has great influence on cell membrane function, cell morphology and structure, the uptake of various nutrients, and metabolite biosynthesis (Li et al., 2008). Effect of different initial pH (4.0 to 7.0) of

the fermentation medium was investigated for the production of  $\beta$ -glucosidase. Optimum  $\beta$ -glucosidase production was obtained at a pH of 5.5 (Figure 5). Most of the filamentous fungi prefer acidic pH for maximum growth and subsequently for high production of enzyme. This might be due to the fact that alkaline pH has inhibitory effect on the growth of the fungus and enzyme production. In addition, cultivation of fungi at an unfavorable pH value may result in reduced enzyme production by reducing accessibility of the substrate (Bakri et al., 2008). Garcia-Kirchner et al. (2005) also optimized pH 5.5 for the production of  $\beta$ -glucosidase from *A. niger*. In contrast, Abdel-Fattah et al. (1997) obtained maximum  $\beta$ -glucosidase activity from *A. niger* at a pH of 4.0.

The effect of different size of conidial inoculum (1.0, 2.0, 3.0, 4.0, and 5.0%) on  $\beta$ -glucosidase production by *A. niger* NRRL 599 in shake flasks for 48 to 120 h was investigated. Maximum  $\beta$ -glucosidase was obtained after 96 h of fermentation with 4% conidial inoculum (Figure 6). Increase or decrease in the size of conidial inoculum resulted in the decreased production of the enzyme. Low inoculum levels might be inadequate for enzyme production while the higher inoculum level might result in rapid depletion of nutrients in the media and reduce the enzyme yield. Also, higher cell density would lead to the production of inhibitory metabolites that may be interfering with the enzyme production (Niladevi and Prema, 2008). In contrast, Abdel-Naby et al. (1999) obtained maximum  $\beta$ -glucosidase activity from *A. niger* with 2%



**Figure 5.** Effect of initial pH on  $\beta$ -glucosidase production by *Aspergillus niger* NRRL 599 in shake flasks.  $\pm$  indicates the standard deviation among the three parallel replicates. Incubation at 30°C for 96 h using wheat bran as carbon source in M-I medium.



**Figure 6.** Effect of inoculum size on  $\beta$ -glucosidase production by *Aspergillus niger* NRRL 599 in shake flasks.  $\pm$  indicates the standard deviation among the three parallel replicates. Incubation at 30°C for 96 h at pH 5.5 using wheat bran as carbon source in M-I medium.


conidial inoculum.

### Enzyme characterization

The enzyme was purified using ammonium sulfate precipitation and FPLC with a purification fold of 4.36.

The purified enzyme was then used for enzyme characterization.

The kinetic constants,  $K_m$  and  $V_{max}$ , for  $\beta$ -glucosidase from *A. niger* were determined at 60°C using pNPG as a substrate and the data was analyzed using Lineweaver-Burk double reciprocal plot (Figure 7).  $\beta$ -Glucosidase exhibited Michaelis-Menten type kinetics and  $K_m$  and  $V_{max}$




**Figure 7.** Lineweaver-Burk double reciprocal plot of  $\beta$ -glucosidase from *Aspergillus niger* NRRL 599. Error bars of standard deviation among the three parallel replicates were too small to be visible.

were found to be 3.11 mM and 20.83 U/mg respectively.  $K_m$  is a measure of affinity of an enzyme for a substrate and low values of  $K_m$  indicate high affinity of the enzyme for the substrate (Hamilton et al., 1998). Seidle et al. (2004) calculated  $K_m$  and  $V_{max}$  of 1 mM and 40 U/mg respectively for pNPG hydrolysis with a  $\beta$ -glucosidase from *A. niger*.  $K_m$  and  $V_{max}$  values of 3.3 mM and 43.68  $\mu$ mol/min. mg respectively were calculated for pNPG hydrolysis by  $\beta$ -glucosidase from the *Melanocarpus* sp. by Kaur et al. (2007). The  $K_m$  and  $V_{max}$  values for pNPG hydrolysis for *Humicola insolens*  $\beta$ -glucosidase were calculated to be 0.16 mM and 18.1 U/mg respectively (Souza et al., 2010).


The effect of different temperature (20 to 65°C) was also investigated on the activity of  $\beta$ -glucosidase. Maximum  $\beta$ -glucosidase activity was obtained at 60°C. Further increase in temperature resulted in decreased  $\beta$ -glucosidase activity. Kimura et al. (1999) and Kaur et al. (2007) also obtained maximum  $\beta$ -glucosidase activity at 60°C from *Aspergillus sojae* and *Melanocarpus* sp. respectively. However, Christakopoulos et al. (1995) and Jeya et al. (2010) obtained maximum  $\beta$ -glucosidase activity at 65°C from *Fusarium oxysporum* and *Penicillium Purpurogenum* respectively. Thermodynamic studies of  $\beta$ -glucosidase were carried out using Arrhenius plot and activation energy ( $E_a$ ), change in enthalpy ( $\Delta H$ ) and change in entropy ( $\Delta S$ ) were calculated to be 52.17 KJ/mol, 49.90 KJ/mol and -71.69 J/mol.K respectively

(Figures 8 and 9). Low values of enthalpy and negative values of entropy indicate the formation of a more efficient and ordered transition state complex between enzyme and substrate (Akolkar and Desai, 2010). Kvesitadze et al. (1990) reported  $\Delta H$  and  $\Delta S$  of 125 KJ/mol and 65 J/mol.K respectively for  $\beta$ -glucosidase of *Aspergillus wentii*.

The activity of the purified  $\beta$ -glucosidase was examined over a pH range of 3.0 to 7.0 at 60°C using sodium acetate buffer. Maximum  $\beta$ -glucosidase activity was achieved at pH 4.8. Further increase or decrease in pH resulted in decrease in activity. Pericin and Jarak (1995) obtained maximum  $\beta$ -glucosidase activity from *Diaporthe (Phomopsis) helianthi* at pH 4.8. In contrast, Leite et al. (2007) obtained maximum activity of  $\beta$ -glucosidase from *Thermoascus aurantiacus* at pH 4.5 while Hang and Woodams (1994), Karnchanatat et al. (2007), Dhake and Patil (2005) and Zanoelo et al. (2004) obtained maximum  $\beta$ -glucosidase activity from *Aspergillus Foetidus*, *Daldinia eschscholzii*, *Penicillium purpurogenum* and *Scytalidium thermophilum* at pH 4.6, 5.0, 5.5 and 6.5 respectively. The change in pH affects the ionization of essential active site amino acid residues, which are involved in substrate binding and catalysis, that is, breakdown of substrate into products. Some ionizable residues may be located on the periphery of the active site, commonly known as non-essential residues. The ionization of these residues may cause distortion of active site cleft and hence indirectly



**Figure 8.** Arrhenius plot for the determination of activation energy ( $E_a$ ) and enthalpy of activation ( $\Delta H$ ) of  $\beta$ -glucosidase from *Aspergillus niger* NRRL 599. Error bars of standard deviation among the three parallel replicates were too small to be visible.




**Figure 9.** Arrhenius plot for the determination of entropy of activation ( $\Delta S$ ) of  $\beta$ -glucosidase from *Aspergillus niger* NRRL 599. Error bars of standard deviation among the three parallel replicates were too small to be visible.

affect the enzyme activity (Amin et al., 2010). Dixon analysis was carried out to evaluate pKa of ionizable groups of active site amino acid residues at 60°C which were involved in  $V_{max}$  (Fig 10). pKa<sub>1</sub> and pKa<sub>2</sub> of  $\beta$ -glucosidase were found to 4.1 and 6.0 respectively. Siddiqui et al.

(1997b) reported pKa<sub>1</sub> and pKa<sub>2</sub> values of 5.5 and 7.9 respectively for  $\beta$ -glucosidase of *Cellulomonas biazotea*. pKa<sub>1</sub> and pKa<sub>2</sub> of 3.3 and 6.9 respectively were reported for  $\beta$ -glucosidase of *Schizophyllum commune* by Clarke (1990).





**Figure 10.** Dixon plot of  $\beta$ -glucosidase from *Aspergillus niger* NRRL 599 at 60°C for the determination of pKa of active site residues that control the  $V_{max}$ .  $\pm$  Indicates the standard deviation among the three parallel replicates.

## ACKNOWLEDGEMENTS

Authors are thankful to HEC Pakistan and GC University, Lahore for all kind of facilities.

## REFERENCES

- Abdel-Fattah AF, Osman MY, Abdel-Naby MA (1997). Production and immobilization of cellobiase from *Aspergillus niger* A 20. Chem. Eng. J. 68: 189-197.
- Abdel-Naby MA, Osman MY, Abdel-Fattah AF (1999). Purification and properties of three cellobiases from *Aspergillus niger* A 20. Appl. Biochem. Biotechnol. 76: 33-44.
- Akolkar AV, Desai AJ (2010). Catalytic and thermodynamic characterization of protease from *Halobacterium* sp. SP1(1). Res. Microbiol. 161: 355-362.
- Almin KE, Eriksson KE, Petersson B (1975). Extracellular enzyme system utilized by the fungus *Sporotrichum pulverulentum* (*Chrysosporium lignorum*) for the breakdown of cellulose. Eur. J. Biochem. 51: 207-211.
- Amin F, Bhatti HN, Asgher M (2010). Partial purification and characterization of an acid invertase from *Saccharum officinarum* L. Pak. J. Bot. 42(4): 2531-2540.
- Bakri Y, Jawahar M, Arabi MIE (2008). Improvement of xylanase production by *Cochliobus sativus* in submerged culture. Food Technol. Biotechnol. 46(1): 116-118.
- Bradford MM (1976). A rapid and sensitive method for the quantitation of the microgram quantities of the protein utilizing the principle of protein-dye binding. Anal. Biochem. 72: 248-254.
- Busto MD, Ortega N, Mateos MP (1995). Induction of  $\beta$ -glucosidase in fungal and soil bacterial cultures. Soil Biol. Biochem. 27(7): 949-954
- Christakopoulos P, Kekos D, Kolisis FN, Macris BJ (1995). Controlling simultaneous production of endoglucanase and beta-glucosidase by *Fusarium oxysporum* in submerged culture. Biotechnol. Lett. 17(8): 883-888.
- Clarke AJ (1990). Chemical modification of a  $\beta$ -glucosidase from *Schizophyllum commune*: evidence for essential carboxyl groups. Biochem Biophys. Acta, 1040: 145-152.
- Coutts AD, Smith RE (1976). Factors influencing the production of cellulase by *Sporotrichum thermophile*. Appl. Environ. Microbiol. 31: p. 819.
- Daroit DJ, Silveira ST, Hertz PF, Brandelli A (2007). Production of extracellular  $\beta$ -glucosidase by *Monascus purpureus* on different growth substrates. Process Biochem. 42: 904-908.
- Dhake AB, Patil MB (2005). Production of  $\beta$ -glucosidase by *Penicillium purpurogenum*. Braz. J. Microbiol. 36: 170-176.
- Dixon M, Webb EC (1979). Enzyme kinetics: Enzymes. New York Academic Press, pp. 44-467.
- Eggs HOW, Pugh GJF (1962). Isolation of cellulose decomposing fungi from soil. Nature, 193: 94-95.
- Fawzi EM (2003). Production and purification of  $\beta$ -glucosidase and protease by *Fusarium proliferatum* NRRL 26517 grown on *Ficus nitida* wastes. Ann. Microbiol. 53(4): 463-476. Garcia-Kirchner O, Sengura-Granados M, Rodriguez-Pascual P (2005). Effect of media composition and growth conditions on production of  $\beta$ -glucosidase by *Aspergillus niger* C-6. Appl. Environ. Microbiol. 121-124: 347-359.
- Grajek W (1987). Hyperproduction of thermostable  $\beta$ -glucosidase by *Sporotrichum (Chrysosporium) thermophile*. Enzyme Microb. Technol. 9(12): 744-748.
- Gunata Z, Vallier MJ (1999). Production of a highly glucose-tolerant extracellular  $\beta$ -glucosidase by three *Aspergillus* strains. Biotechnol. Lett. 21(3): 219-223.
- Hamilton LM, Kelly CT, Fogarty MW (1998). Raw starch degradation by the non raw starch-adsorbing bacterial  $\alpha$ -amylase of *Bacillus* sp. IMD 434. Carbohydr. Res. 314: 251-257.
- Hang YD, Woodams EE (1994). Apple pomace: A potential substrate for production of  $\beta$ -glucosidase by *Aspergillus Foetidus*. Lebensm.-Wiss. Technol. 27(6): 587-589.
- Haq I, Javed MM, Khan TS (2006). An innovative approach for

- hyperproduction of cellulolytic and hemicellulolytic enzymes by consortium of *Aspergillus niger* MSK-1 and *Trichoderma viride* MSK-10. Afr. J. Biotechnol. 5(8): 609-614.
- Iembo T, Da-Silva R, Pagnocca FC, Gomes E (2002). Production, characterization and properties of beta-glucosidase and beta-xylosidase from a strain of *Aureobasidium* sp. Prikl. Biokhim. Mikrobiol. 38(6): 639-643.
- Iwashita K, Nagahara T, Kimura K, Takano M, Shimoi H, Ito K (1999). The *bgIA* gene of *Aspergillus kawachii* encodes both extracellular and cell wall bound  $\beta$ -glucosidases. Appl. Environ. Microbiol. 65(12): 5546-5553.
- Jager S, Brumbauer A, Feher E, Reczey K, Kiss L (2001). Production and characterization of  $\beta$ -glucosidases from different *Aspergillus* strains. World J. Microbiol. Biotechnol. 17(5): 455-461.
- Jakubikova L, Farkas V, Kolkarova N, Nemcovic M (2006). Conidiation of *Trichoderma atroviride* isolate during submerged cultivation in a laboratory stirred-tank fermenter. Folia Microbiol. 51(3): 209-213.
- Jeya M, Joo AR, Lee KM, Tiwari MK, Lee KM, Kim SH, Lee JK (2010). Characterization of  $\beta$ -glucosidase from a strain of *Penicillium Purpurogenum* KJS506. Appl. Microbiol. Biotechnol. 86: 1473-1484.
- Karnchanat A, Petsom A, Sangvanich P, Piaphukiew J, Whalley AJS, Reynolds CD, Sihanonth P (2007). Purification and characterization of an extracellular beta glucosidase from the wood-decaying fungus *Daldinia eschscholzii*. FEMS Microbiol. Lett. 270(1): 162-170.
- Kaur J, Chadha BS, Kumar BA, Kaur GS, Saini HS (2007). Purification and characterization of  $\beta$ -glucosidase from *Melanocarpus* sp. MTCO 3922. Electron. J. Biotechnol., 10(2): 260-270.
- Kimura I, Yoshioka N, Tajima S (1999). Purification and characterization of a  $\beta$ -glucosidase with  $\beta$ -xylosidase activity from *Aspergillus sojae*. J. Biosci. Bioeng. 87(4): 538-541.
- Kvesitadze GI, Svanidze RS, Tsuprun VL, Nizharadze DN, Chirgadze LT, Buachidze TS (1990). Quaternary structure and properties of  $\beta$ -glucosidase isolated from a thermophilic culture of *Aspergillus wentii*. Bioorg. Khim. 16: 881-888.
- Leite RSR, Gomes E, Da Silva R (2007). Characterization and comparison of thermostability of purified  $\beta$ -glucosidases from a mesophilic *Aureobasidium pullulans* and a thermophilic *Thermoascus aurantiacus*. Process Biochem. 42: 1101-1106.
- Li J, Li P, Liu F (2008). Production of theanine by *Xerocomus badius* (mushroom) using submerged fermentation. LWT- Food Sci. Technol. 41: 883-889.
- Lineweaver H, Burk D (1934). The determination of enzyme dissociation constants. J. Am. Chem. Soc. 56: 658-666.
- Macris BJ, Panayatou MG (1989). Enhanced cellobiohydrolase production from *Aspergillus ustus* and *Trichoderma harzianum*. Enzyme Microb. Technol. 8: 141-144.
- Mandel M, Reese ET (1960). Fungal cellulases and the microbial decomposition of cellulosic fabric. Dev. Microbiol. 5: 5-19.
- Niladevi KN, Prema P (2008). Effect of inducers and process parameters on laccase production by *Streptomyces psammoticus* and its application in dye decolourization. Bioresour. Technol. 99: 4583-4589.
- Pericin D, Jarak M (1995). Production and some characteristics of beta glucosidase in *Diaporthe (Phomopsis) helianthi*. Acta Microbiol. Immunol. Hung. 42(1): 29-37.
- Rajoka MI, Akhtar MW, Hanif A, Khalid AM (2006). Production and characterization of a highly active cellobiase from *Aspergillus niger* grown in solid state fermentation. World J. Microbiol. Biotechnol. 22(9): 991-998.
- Rajoka MI, Khan S, Latif F, Shahid R (2004). Influence of carbon and nitrogen sources and temperature on hyperproduction of a thermotolerant  $\beta$ -glucosidase from synthetic medium by *Kluyveromyces marxianus*. Appl. Biochem. Biotechnol. 117(2): 75-92.
- Rajoka MI, Malik KA (1997). Cellulase production by *Cellulomonas biazotea* cultured in media containing different cellulosic substrates. Bioresour. Technol. 59(1): 21-27.
- Riou C, Salmon JM, Vallier MJ, Gunata Z, Barre P (1998). Purification, characterization and substrate specificity of a highly glucose tolerant  $\beta$ -glucosidase from *Aspergillus oryzae*. Appl. Environ. Microbiol. 64(10): 3607-3614.
- Romanelli RA, Houston CW, Barnett SM (1975). Studies on thermophilic cellulolytic fungi. Appl. Microbiol. 30: 276-281.
- Saddler JN (1982). Screening of high cellulolytic fungi and the action of their cellulase enzyme systems. Enzyme Microb. Technol. 4: 414-418.
- Seidle HF, Huber RE (2005). Transglucosidic reactions of the *Aspergillus niger* family 3  $\beta$ -glucosidase: Qualitative and quantitative analyses and evidence that the transglucosidic rate is independent of pH. Arch. Biochem. Biophys. 436: 254-264.
- Seidle HF, Marten I, Shoseyov O, Huber RE (2004). Physical and kinetic properties of the family 3  $\beta$ -Glucosidase from *Aspergillus niger* which is important for cellulose breakdown. Protein J. 23(1): 11-23.
- Siddiqui KS, Rashid MH, Ghauri TM, Durrani IS, Rajoka MI (1997a). Purification and characterization of an intracellular  $\beta$ -glucosidase from *Cellulomonas biazotea*. World J. Microbiol. Biotechnol. 13: 245-247.
- Siddiqui KS, Rashid MH, Rajoka MI (1997b). Kinetic analysis of the active site of an intracellular  $\beta$ -glucosidase from *Cellulomonas biazotea*. Folia Microbiol. 42: 53-58.
- Snedecor G, Cochran WG (1980). Statistical Methods. 7<sup>th</sup> Edition, Iowa State Univ., pp. 80-86.
- Souza FHM, Nascimento CV, Rosa JC, Masui DC, Leone FA, Jorge JA, Furriel RPM (2010). Purification and biochemical characterization of a mycelial glucose- and xylose-stimulated  $\beta$ -glucosidase from the thermophilic fungus *Humicola insolens*. Process Biochem. 49: 272-278.
- Sun X, Liu Z, Qu Y, Li X (2008). The effects of wheat bran composition on the production of biomass-hydrolyzing enzymes by *Penicillium decumbens*. Appl. Biochem. Biotechnol. 146: 119-128.
- Taylor MJ, Richardson T (1979). Application of microbial enzymes in food systems and in biotechnology. Adv. Appl. Microbiol. 25: 7-35.
- Tsao GT, Xia L, Cao N, Gong CS (2000). Solid state fermentation with *Aspergillus niger* for cellobiase production. Appl. Biochem. Biotechnol. 84-86: 743-749.
- Villena MA, Iranzo JFU, Parez AIB (2007).  $\beta$ -glucosidase activity in wine yeasts: Application in enology. Enzyme Microb. Technol. 40: 420-425.
- Wallecha A, Mishra S (2003). Purification and characterization of two  $\beta$ -glucosidases from a thermo-tolerant yeast *Pichia etchellsii*. Biochim. Biophys. Acta, 16(49): 74-84.
- Zanoelo FF, Polizeli MLTM, Terenzi HF, Jorge JA (2004).  $\beta$ -glucosidase activity from the thermophilic fungus *Scytalidium thermophilum* is stimulated by glucose and xylose. FEMS Microbiol. Lett. 240(2): 137-143.