

Full Length Research Paper

# Effects of phosphate solubilizing microorganisms and *Rhizobium* sp. on the growth, nodulation, yield and root-rot disease complex of chickpea under field condition

Mohd Sayeed Akhtar<sup>1,2\*</sup> and Zaki A. Siddiqui<sup>1</sup>

<sup>1</sup>Department of Botany, Aligarh Muslim University, Aligarh-202 002, India.

<sup>2</sup>Botanical Institute, University of Basel, Hebelstrasse 1, 4056, Basel, Switzerland.

Accepted 8 May, 2009

Effects of Phosphate solubilizing microorganisms (*Glomus intraradices*, *Pseudomonas putida*, *P. alcaligenes*, *P. aeruginosa* (Pa28), *A. awamori*) and *Rhizobium* sp. was observed on the growth, nodulation yield and root-rot disease complex of chickpea under field condition. Inoculation of *Rhizobium* sp. caused a greater increase in growth and yield than *P. putida*, *P. aeruginosa* or *G. intraradices*. The number of nodules per root system was significantly higher in plants inoculated with *Rhizobium* sp. compared to plants without *Rhizobium* sp. Inoculation of *P. putida* caused highest reduction in galling followed by *P. aeruginosa*, *P. alcaligenes*, *G. intraradices* and *A. awamori* while *Rhizobium* sp. caused almost similar reduction in galling as caused by *P. putida*.

**Key words:** Chickpea, *Glomus* sp., phosphate solubilizing microorganisms, disease complex.

## INTRODUCTION

India is the largest producer and importer of the legume-crop (Shakya et al., 2008). Amongst the leguminous crops, chickpea (*Cicer arietinum* L.) occupy an important position due to its nutritive values (17-23% protein) in large vegetarian population of the country (Ali and Kumar, 2006). The major chickpea producing states are Andhra Pradesh, Madhya Pradesh, Maharashtra, Rajasthan and Uttar Pradesh contributes 86% of the total production of the country. During last decade the production of chickpea has been declined in the western Uttar Pradesh. The main cause behind this is the root knot nematode *Meloidogyne incognita* (Kofoid and White) Chitwood and *Macrophomina phaseolina* (Tassi) Goid (Akhtar and Siddiqui, 2007a). The synergistic interaction of these pathogens together causes more loss in term of yield than the sum of their individuals to this important pulse crop (Siddiqui and Husain, 1992; Akhtar and Siddiqui, 2008a, b).

The rhizosphere is the zone of intense microbial activity

and provides a front line defense against pathogen attack to root (Weller, 1988). The rhizoplane and surrounding rhizosphere soil are colonized or occupied by a wide range of microorganisms. Of the various microorganisms present, AM (arbuscular mycorrhizal) fungi improves the plant growth, nutrient cycling and protection against biotic and abiotic stresses and soil structure through aggregate formation (Bethlenfalvay and Linderman, 1992; Gianinazzi and Schuepp, 1994; Kapulnik and Douds, 2000; van der Heijden and Sanders, 2002; Jeffries et al., 2003; Barea et al., 2005; Smith and Read, 2008). The AM symbiosis increases the supply of mineral nutrients particularly phosphorus and other minerals like ammonium, zinc and copper. The AM association also improves plant health through increased protection against soil-borne microbial pathogens (Allen, 1996; Gianinazzi et al., 2002; Jefferies et al., 2003; Barea et al., 2005; Akhtar and Siddiqui, 2008a).

Some other microorganisms that are associated with the roots of crop plants and play an important role in mobilization and immobilization of phosphorus are known as phosphate solubilizing microorganisms (PSM). PSM largely include bacteria and fungi viz. some of the

\*Corresponding author. E-mail: sayeedbot@yahoo.co.in. Tel.: +91-9719066915. Fax: +91-571-2402016.

species of *Bacillus*, *Pseudomonas*, *Penicillium* and *Aspergillus* (Tilak, 1991; Tilak et al., 2005). PSM can solubilize and mineralize P from inorganic and organic pools of total soil P and may be used as inoculants to increase P-availability to plants (Richardson, 2001; Illmer et al., 1995; Whitelaw et al., 1999) and also have the capacity to increase the growth and yield of crop plants (Gupta and Namdeo, 1997; Ozgonen et al., 1999) besides reducing disease severity (Weller, 1988; Siddiqui and Mahmood, 1999). Similarly, presence of rhizobia in the rhizosphere may also protect the host root from damage caused by pathogens (Siddiqui and Mahmood, 1995; Elbardy et al., 2006; Huang and Erickson, 2007).

The consistency in performance of different biocontrol agent have been tested in our previous experiments (Siddiqui and Akhtar, 2007; Akhtar and Siddiqui, 2007a, b; Akhtar and Siddiqui, 2008 a, b). On the basis of the results drawn from the earlier experiments the biocontrol agents which perform best under pot trials were selected for the field trial. The aim of the present study was to investigate the effects of PSM (found best in pot trial in our previous experiments) and *Rhizobium* sp. on the growth, nodulation yield and on the root-rot disease complex of chickpea in field condition and also to find out the best biocontrol agent under field condition.

## MATERIALS AND METHODS

### Site location and field history

Aligarh is situated at 27°53' N latitude, 78°4' E longitude and 187.45 m altitude with an area of 3431 sq km. Its climate is sub-tropical, with hot dry summer and cold winters. The winter extends from the middle of October to the end of March. The mean temperature during this season ranges from 23°C and 12°C respectively.

The average rainfall is 847.3 mm. More than 85% of the total rainfall occurs during June to September and some 10% in the winter. The winter rainfall is useful for Rabi crops. The relative humidity of the winter season ranges between 56 and 77% with an average of 66.5% that of the summer, between 37 and 49% with an average of 43% and that of the monsoon season, between 63 and 73% with an average 68%.

A field experiment was conducted at University Fort, Aligarh Muslim University. Aligarh generally available for the field research work (soil sandy loam, pH 7.5, porosity 39%, water holding capacity 42%, electrical conductivity 0.67, available N 95.90 mg/kg soil, available P 11.4 mg/kg soil and available K 160.89 mg/kg soil). The site at Fort was selected with a view suitable for chickpea crop.

Maize (*Zea mays*) was harvested September, 2006. The field was prepared by ploughing and removing grasses and plant debris.

### Raising and maintenance of plants

Healthy, uniform size seeds of chickpea (*Cicer arietinum* L.) cv. Avarodhi were surface sterilized with 0.1% sodium hypochlorite (NaOCl) and washed 3 times with distilled water. Sowing was done by the usual 'behind the plough' method at the rate of 60 kg / ha on October 25, 2006. After germination, the distance between seedlings and rows was maintained at 20 cm and 40 cm respectively. The plots were free from weed, thrice. The plants were irrigated thrice at 40, 80 and 120 days after sowing and were harvested at 150 days.

### Nematode Inoculum

*Meloidogyne incognita* was collected from chickpea field soil identified with the help perineal pattern (Taylor and Sasser, 1978). The pure culture of nematode was multiplied on egg plant (*Solanum melongena* L.) using a single egg mass placed in a glass house at the Department of Botany, A.M.U., Aligarh, India. Egg masses were hand-picked using sterilized forceps and placed in 9 cm diameter sieves of 1 mm pore size, which were previously mounted with cross layered tissue paper and placed in a petri plates containing water just deep enough to contact the egg masses. The sieves were placed in petri dishes with distilled water for hatching and incubated at 27°C. The hatched second stage juveniles ( $J_2$ ) were collected from the petri plates every 24 h and fresh water was added to the petri plates. The concentration of  $J_2$  of *M. incognita* in the water was adjusted so that each ml contained  $200 \pm 5$  nematodes. 10 ml of this suspension (that is, 2000 freshly hatched  $J_2$ ) was added to each pot containing a chickpea seedling.

### Fungus Inoculum

*Macrophomina phaseolina* was isolated from infected chickpea root samples collected from Kasimpur, Aligarh (KA1) on potato dextrose agar (PDA). On confirmation its identity, the pure culture of the fungus was maintained on PDA. Fungal inoculum was prepared by culturing the isolates in Richard's medium (Riker and Riker, 1936) for 15 days at 25°C. Mycelium was collected on blotting sheets to remove excess water and nutrients. 100 g of mycelium was macerated in 1 l distilled water. 10 ml of this suspension (containing 1 g fungus) was applied to each plant.

### Preparation of PSM inoculum

The AM fungus, *Glomus intraradices* Schenck and Smith, was isolated from the soil of chickpea field of Aligarh, India. The species was identified using the synoptic keys of Trappe (1982) and Schenck and Pervez (1990) and the identification was reconfirmed from Curator, INVAM, USA (<http://invam.caf.wvu.edu/>). For inoculum, the fungus was produced on *Chloris gayana* Kunth (Rhodes grass) grown in sandy loam soil. The population of *G. intraradices* isolate AI03 (Aligarh, India) in the inoculum was assessed by the most probable number method (Porter, 1979). 50 g of inoculum with soil was added around the seed to provide 500 infective propagules of *G. intraradices* per pot (1 g inoculum contains 10 infective propagules). The crude inoculum consisted of soil, extra metrical spores and spore carps, hyphal fragments and infective rhodes grass segments.

*Paenibacillus polymyxa* (MTCC No. 122), *Pseudomonas putida* (MTCC No. 3604), and *Pseudomonas alcaligenes* (MTCC No. 493) were obtained from Microbial Type of Culture Collection and Gene Bank, Institute of Microbial Technology, Chandigarh, India while the *P. aeruginosa* (Pa28) isolated from chickpea soil. These isolates were sub culture on nutrient broth (Hi-Media Laboratories, Mumbai, India) incubated at  $37 \pm 1^\circ\text{C}$  for 72 h. 1 ml nutrient broth suspension contains about  $1.5 \times 10^7$  cfu /ml. 10 ml of this suspension was inoculated into each pot around the pea seedling. Another PSM, *Aspergillus awamori* Nakazawa (ITCCF 4680) which was obtained from Indian Agricultural Research Institute, New Delhi and cultured in Richard's medium at  $25 \pm 1^\circ\text{C}$  for 15 days. 10 ml (equivalent to 1 g) suspension was inoculated into each pot around the chickpea seedling.

### Rhizobium inoculum

100 g commercial culture of *Rhizobium* Jordan (chickpea strain AP)

**Table 1.** Effect of phosphate solubilizing microroganisms (*G. intraradices*, *P. putida*, *P. alcaligenes*, *P. aeruginosa* (Pa28), *A. awamori*) and *Rhizobium* sp. on the growth of chickpea under field condition.

Treatments		Shoot dry weight (g)	No. of pods/plant	No. of nodules/ root system
Control	C	26.30 ± 0.36 e	26 ± 3.16 g	48 ± 3.16 fg
	Gi	27.91 ± 0.41 d	30 ± 3.74 efg	54 ± 2.74 bcd
	Pp	29.92 ± 0.48 b	34 ± 4.69 e	58 ± 3.74 ab
	Pa	28.61 ± 0.24 c	31 ± 4.36 ef	56 ± 4.00 bc
	A	27.66 ± 0.31 d	29 ± 5.48 fg	53 ± 3.81 cde
	R	30.74 ± 0.15 a	81 ± 5.92 a	62 ± 4.00 a
	Pa28	28.92 ± 0.33 c	32 ± 4.00 ef	56 ± 3.16 bc
<i>M. incognita</i>	C	18.76 ± 0.65 p	9 ± 2.24 jklm	33 ± 3.24 m
	Gi	22.10 ± 0.22 kl	11 ± 2.00 ijkl	40 ± 3.74 j kl
	Pp	24.18 ± 0.48 g	13 ± 2.55 hij	49 ± 2.23 efg
	Pa	22.87 ± 0.45 ij	10 ± 2.54 ijkl	41 ± 2.55 ijk
	A	21.24 ± 0.38 m	12 ± 2.45 hijk	46 ± 3.16 gh
	R	24.88 ± 0.33 f	56 ± 5.48 c	51 ± 2.92 def
	Pa28	23.58 ± 0.40 h	11 ± 3.08 ijkl	46 ± 4.47 gh
<i>M. phaseolina</i>	C	18.88 ± 0.26 p	12 ± 1.87 hijk	36 ± 4.00 lm
	Gi	22.50 ± 0.28 jk	13 ± 2.55 hij	42 ± 3.16 hij
	Pp	24.92 ± 0.48 f	6 ± 1.87 h	51 ± 2.91 def
	Pa	23.34 ± 0.45 hi	12 ± 2.45 hijk	42 ± 3.08 hij
	A	21.67 ± 0.46 l m	12 ± 1.87 hijk	48 ± 2.45 fg
	R	25.25 ± 0.30 f	62 ± 4.00 b	53 ± 3.81 cde
	Pa28	24.19 ± 0.52 g	14 ± 2.34 hi	46 ± 3.08 gh
<i>M. incognita</i> + <i>M. phaseolina</i>	C	12.55 ± 0.17 r	5 ± 1.41 m	21 ± 2.73 n
	Gi	18.76 ± 0.52 p	8 ± 2.12 klm	33 ± 2.74 m
	Pp	20.54 ± 0.49 n	10 ± 2.55 ijkl	43 ± 3.61 hij
	Pa	19.56 ± 0.17 v o	7 ± 2.34 l m	37 ± 2.00 klm
	A	18.14 ± 0.19 q	9 ± 2.24 jklm	40 ± 3.16 jkl
	R	21.48 ± 0.18 m	40 ± 5.10 d	45 ± 3.31 ghi
	Pa28	20.10 ± 0.30 n	8 ± 2.12 klm	40 ± 3.74 jkl
L.S.D. p = 0.05		0.47	4	4

C = Control; Gi = *G. intraradices*; Pp = *P. putida*; Pa = *P. alcaligenes*; A = *A. awamori*; Pa = *P. aeruginosa*; R = *Rhizobium* sp.

\*Values within each column followed by same letter are not significantly different ( $p = 0.05$ ).  $\pm$  = Standard deviation.

was suspended in 1000 ml distilled water and 10 ml (equivalent to 1 g inoculum) into each pot around chickpea seedling after thinning.

#### Inoculation technique

For inoculation of *M. incognita*, *M. phaseolina*, PSM (*G. fasciculatum*, *P. putida*, *P. alcaligenes*, *P. aeruginosa* (Pa28), *A. awamori*) and *Rhizobium* sp. and soil around the roots was carefully moved aside without damaging the roots. The inocula of these microorganisms were poured or placed around the roots and the soil was replaced. An equal volume of sterile water was added to control treatments.

#### Experimental design

Experiment was conducted in randomized block design and the

size of each plot was 9 square meter. There were 7 treatments comprising of *G. intraradices*, *P. putida*, *P. alcaligenes*, *A. awamori*, *P. aeruginosa* (isolate 28), *Rhizobium* sp. and control. These 7 treatments were also tested against 4 pathogen combination ( $7 \times 4 = 28$  treatments) and each set was replicated 3 times ( $7 \times 4 \times 3 = 84$  pots) as listed in Table 1.

#### Observations

The plants were harvested 150 days after inoculation. Data were recorded on shoot dry weight, number of pods, number of nodules, number of galls, grain weight, yield and root-rot index. Each plant was cut with a knife above the base of the root-emergence zone to separate shoot and root. The numbers of galls per root system were counted. For dry-weight determination, shoots were kept in envelopes at 60°C for 2-3 days and weight was recorded in g. A root-rot index was determined by scoring on a scale ranging from 0 (no

disease) to 5 (severe root-rot).

### Statistical analysis

The entire dataset was analyzed as a single two-factor experiment (pathogens × PSM / *Rhizobium* sp.) as described by Dospekhov (1984). All analyses were carried out using Stat view 5.0 (SAS institute, Cary, NC, USA). Least significant differences (L.S.D.) were calculated at  $p = 0.05$  and Duncan's multiple-range test was used to test significant differences between treatments. Standard deviation was also calculated for the variance. The graph for galling was prepared using the program sigma plot version 9.0 with error bars showing standard deviation.

## RESULTS

Treatment of plants without pathogens with *G. intraradices*, *P. putida*, *P. alcaligenes*, *A. awamori*, *P. aeruginosa* (Pa28) and *Rhizobium* caused a significant increase shoot dry weight over uninoculated ones (Table 1). Inoculation of *Rhizobium* to plants without pathogens caused a greater increase in shoot dry weight to that caused by *P. putida*, *P. aeruginosa* or *G. intraradices*. However, increase in shoot dry weight of plants with *P. alcaligenes* was similar to that caused *P. aeruginosa* while *A. awamori* increased shoot dry weight similar to that caused by *G. intraradices* (Table 1).

Inoculation of plants with *M. incognita* and *M. phaseolina* alone and in combination caused a significant reduction in shoot dry weight over uninoculated ones (Table 1). Reduction in shoot dry weight was greater when *M. incognita* and *M. phaseolina* were inoculated together than the inoculation of either of them. Inoculation of *G. intraradices*, *P. putida*, *P. alcaligenes*, *A. awamori*, *P. aeruginosa* and *Rhizobium* sp. significantly increased shoot dry weight of plants with pathogens. Inoculation of *Rhizobium* sp. to plants with pathogens caused a greater increase in shoot dry weight than by *P. putida*, *P. aeruginosa* or *G. intraradices*. However, inoculation of *P. alcaligenes* to plants with pathogens caused almost similar increase in shoot dry weight as by *P. aeruginosa* while *A. awamori* increased shoot dry weight similar to that caused by *G. intraradices* (Table 1).

Inoculation of *M. incognita* / *M. phaseolina* or both significantly reduced the number of pods per plant. Inoculation of *G. intraradices*, *P. putida*, *P. alcaligenes*, *A. awamori*, *P. aeruginosa* and *Rhizobium* sp. significantly increased the number of pods per plant both in pathogens inoculated and uninoculated ones. The number of nodules per root system was significantly higher in plants inoculated with *Rhizobium* sp. compared to uninoculated one (Table 1).

Inoculation of plants without pathogens with *G. intraradices*, *P. putida*, *P. alcaligenes*, *A. awamori*, *P. aeruginosa* and *Rhizobium* sp. significantly increased seed weight (per 100 seeds) and yield over uninoculated ones (Table 2). Inoculation of *Rhizobium* sp. to plants without pathogens caused a greater increase in seed weight and

yield than by *P. putida* or *P. aeruginosa* or *G. intraradices*. However, increase in seed weight and yield of plants with *P. alcaligenes* was similar to that caused *P. aeruginosa* while *A. awamori* increased seed weight and yield similar to that caused by *G. intraradices* (Table 2).


Inoculation of plants with *M. incognita* and *M. phaseolina* alone and in combination caused a significant reduction in seed weight and yield over uninoculated controls (Table 2). Inoculation of *G. intraradices*, *P. putida*, *P. alcaligenes*, *A. awamori*, *P. aeruginosa* and *Rhizobium* sp. significantly increased seed weight and yield of plants with pathogens. Inoculation of *Rhizobium* caused a greater increase in seed weight and yield than by *P. putida*, *P. aeruginosa* or *G. intraradices*. However, inoculation of *P. alcaligenes* to plants with pathogens caused almost similar increase in seed weight and yield as caused by *P. aeruginosa* while *A. awamori* increased seed weight and yield similar to that caused by *G. intraradices* (Table 2).

The number of galls per root system was high when *M. incognita* was inoculated alone (Figure 1). In presence of *M. phaseolina* root galling were reduced. Inoculation of *P. putida* caused highest reduction in galling followed by *P. aeruginosa*, *P. alcaligenes*, *G. intraradices* and *A. awamori*. *Rhizobium* sp. caused almost similar reduction in galling as caused by *P. putida* (Figure 1).

Root-rot index was 4 when *M. incognita* and *M. phaseolina* were inoculated together (Figure 2). Index was reduced to 3 when *M. incognita* plus *M. phaseolina* inoculated plants were treated with *G. intraradices* or *A. awamori* while index was found to 2 when *M. incognita* plus *M. phaseolina* inoculated plants were treated singly with *P. putida*, *P. alcaligenes*, *P. aeruginosa* and *Rhizobium* sp. Index was reduced to 1 when *M. phaseolina* inoculated plants was treated singly with *G. intraradices*, *P. putida*, *P. alcaligenes*, *A. awamori*, *Rhizobium* sp. and *P. aeruginosa* (Figure 2).

## DISCUSSION

Among the 6 biocontrol agents tested under field condition, *Rhizobium* sp. performed better in the management of root-rot disease complex followed by *P. putida*, Pa28, *P. alcaligenes*, *G. intraradices* and *A. awamori*. Root nodules formed by *Rhizobium* sp. under field condition were larger than the nodules formed in the plants under pot condition (Akhtar and Siddiqui, 2008a). Higher nitrogen fixation by large size nodules may account for reduced development (Barker and Huisling, 1970). *P. putida* had a greater siderophore production than *P. alcaligenes* (Siddiqui et al., 2007). This may be a reason that *P. putida* was better than *P. alcaligenes* under field condition. Pa28 also showed greater siderophores production among pseudomonads isolates from chickpea fields and performed better than *G. intraradices* and *A. awamori*. Least effectiveness of *G. intraradices* and *A. awamori* may be attributed to their inability to adapt in the fields condition with other soil microflora.



**Figure 1.** Effect of PSM (*G. intraradices*, *P. putida*, *P. alcaligenes*, *P. aeruginosa* (Pa28), *A. awamori*) and *Rhizobium* sp. on the galling of chickpea under field condition. C = Control; Gi = *G. intraradices*; Pp = *P. putida*; Pa = *P. alcaligenes*; A = *A. awamori*; Pa = *P. aeruginosa*; R = *Rhizobium* sp. \*Different letters within 1 parameter are significantly different at  $p = 0.05$ . Error bars represent Standard deviation.

In this study, *G. intraradices* improved plant growth of nematode-infected plants by reducing nematode multiplication. The root-rot severity of plants inoculated with *M. phaseolina* was also reduced by *G. intraradices* (Akhtar and Siddiqui, 2007b). Bodker et al. (1998) observed a reduction in root-rot of pea caused by *Aphanomyces euteiches*, while Akkopru and Demir (2005) observed about 17% reduction in Fusarium wilt of tomato after inoculation of plants with *G. intraradices*. In addition, the changes in nutrient uptake in the root system, a mycorrhizosphere effect and activation of plant defense mechanisms are thought to be responsible for disease inhibition by AM fungi (Linderman, 1994; Demir and Akkopru, 2005). Moreover, treatment with *Glomus* sp. is also reported to increase phenylalanine and serine in tomato roots (Suresh, 1980), these amino acids have an

inhibitory effect on nematodes (Reddy, 1974).

*Pseudomonads* may also improve plant growth by suppressing parasitic and non-parasitic root pathogens (Oostendrop and Sikora, 1989) through the production of biologically active substances (Gamliel and Katan, 1993) or the conversion of unavailable minerals and organic compounds into forms that are available to plants (Broadbent et al., 1977; Siddiqui and Mahmood, 1999). *Pseudomonas* can synthesize enzymes that can modulate plant hormone levels, may limit the available iron via siderophore production and can also kill the pathogen with antibiotics (Siddiqui, 2006). In addition, induced systemic resistance by *Pseudomonas* is also thought to be a biocontrol mechanism against plant pathogens (Wei et al., 1996). Similarly *Bacillus* isolates have been reported to promote the growth of a wide range of plants

**Table 2.** Effect of phosphate solubilizing microroganisms (*G. intraradices*, *P. putida*, *P. alcaligenes*, *P. aeruginosa* (Pa28), *A. awamori*) and *Rhizobium* sp. on the yield and of chickpea under field condition.

Treatment		Grain (100 seeds) weight (g)	Yield (q/ha)
<b>Control</b>	C	14.76 ± 0.43 d	16.04 ± 0.26 e
	Gi	15.60 ± 0.14 c	17.02 ± 0.33 d
	Pp	16.54 ± 0.26 a	18.25 ± 0.38 b
	Pa	15.92 ± 0.23 b	17.45 ± 0.44 c
	A	15.34 ± 0.17 c	16.87 ± 0.18 d
	R	16.72 ± 0.11 a	18.75 ± 0.26 a
	Pa28	16.20 ± 0.32 b	17.64 ± 0.36 c
<b><i>M. incognita</i></b>	C	10.94 ± 0.30 p	11.44 ± 0.22 p
	Gi	13.06 ± 0.36 k	13.48 ± 0.18 kl
	Pp	14.36 ± 0.26 ef	14.74 ± 0.42 g
	Pa	13.64 ± 0.17 ij	13.95 ± 0.31 ij
	A	12.68 ± 0.13 l	12.95 ± 0.29 m
	R	14.47 ± 0.08 def	15.17 ± 0.24 f
	Pa28	14.04 ± 0.33 gh	14.38 ± 0.30 h
<b><i>M. phaseolina</i></b>	C	11.06 ± 0.24 op	11.51 ± 0.42 p
	Gi	13.40 ± 0.24 j	13.72 ± 0.33 jk
	Pp	14.58 ± 0.22 de	15.20 ± 0.19f
	Pa	13.82 ± 0.23 hi	14.23 ± 0.25 hi
	A	12.79 ± 0.19 kl	13.21 ± 0.29 lm
	R	14.60 ± 0.25 de	15.40 ± 0.37 f
	Pa28	14.26 ± 0.08 fg	14.75 ± 0.21 g
<b><i>M. incognita</i> + <i>M. phaseolina</i></b>	C	9.15 ± 0.32 q	7.65 ± 0.24 r
	Gi	11.72 ± 0.30 n	11.44 ± 0.36 p
	Pp	12.62 ± 0.35 l	12.52 ± 0.28 n
	Pa	12.05 ± 0.31 m	11.93 ± 0.24 o
	A	11.35 ± 0.23 o	11.06 ± 0.21 q
	R	12.85 ± 0.18 kl	13.10 ± 0.44 m
	Pa28	12.15 ± 0.18 m	12.26 ± 0.38 no
L.S.D. p = 0.05		0.31	0.34

C = Control; Gi = *G. intraradices*; Pp = *P. putida*; Pa = *P. alcaligenes*; A = *A. awamori*; Pa = *P. aeruginosa*; R = *Rhizobium* sp.


\*Values within each column followed by same letter are not significantly different (p = 0.05).

± = Standard deviation

(De Freitas et al., 1997; Kokalis-Burelle et al., 2002). *B. pumilus* induces callose and pectin in close association with phenolic compounds in newly formed wall appositions in pea roots in response to attack by *Fusarium oxysporum* (Benhamou et al., 1996). However, treatment with *B. pumilus*, induced a rapid lignification in cucumber plants in response to ingress of *Colletotrichum orbiculare*, and total peroxidase and superoxide dismutase activities increased more than those in the buffer control (Jetiyanon et al., 1997). These responses may be due to the production of siderophores, antibiotics, wall apposi-

tions and defense enzymes, which adversely affect on the pathogens.

*G. intraradices*, *P. alcaligenes* and *B. pumilus* were used individually and concomitantly to control the root-rot disease complex of chickpea (Akhtar and Siddiqui, 2008b). Phosphate solubilizing bacterial isolate that promote and stimulate colonization by the AM fungus are called mycorrhiza helper bacteria (Barea et al., 1998) and they also stimulate the germination of AM spore and mycelial development (Meyer and Linderman, 1986). The present study demonstrated that PSM can coexist without



**Figure 2.** Effect of PSM (*G. intraradices*, *P. putida*, *P. alcaligenes*, *P. aeruginosa* (Pa28), *A. awamori*) and *Rhizobium* sp. on the on the root-rot indices under field condition. C = Control; Gi = *G. intraradices*; Pp = *P. putida*; Pa = *P. alcaligenes*; A = *A. awamori*; Pa = *P. aeruginosa*; R = *Rhizobium* sp.

adversely affecting each other. In fact, suitable combinations of these biocontrol agents may be increase plant growth and resistance to pathogens. In future, more detail investigations are needed on the performance of PSM on different hosts of various pathosystems.

## REFERENCES

- Akhtar MS, Siddiqui ZA (2007a). Biocontrol of a chickpea root-rot disease complex with *Glomus intraradices*, *Pseudomonas putida* and *Paenibacillus polymyxa*. Aust. Plant Pathol. 36: 175-180.
- Akhtar MS, Siddiqui ZA (2007b). Effects of *Glomus fasciculatum* and *Rhizobium* sp. on the growth and root-rot disease complex of chickpea. Arch. Phytopathol. Plant Protec. 40: 37-43.
- Akhtar MS, Siddiqui ZA (2008a). Biocontrol of a root-rot disease complex of chickpea by *Glomus intraradices*, *Rhizobium* sp. and *Pseudomonas straita*. Crop Protec. 23: 410-417.
- Akhtar MS, Siddiqui ZA (2008b). *Glomus intraradices*, *Pseudomonas alcaligenes*, *Bacillus pumilus* as effective biocontrol agents for the root-rot disease complex of chickpea (*Cicer arietinum* L.). J. Gen. Plant Pathol. 74: 53-60.
- Ali M, Kumar S (2006). Pulse production in India. Yojana, Sept. pp. 13-15.
- Akkopru A, Demir S (2005). Biological control of Fusarium wilt in tomato caused by *Fusarium oxysporum* f. sp. *lycopersici* by AMF *Glomus intraradices* and some rhizobacteria. J. Phytopathol. 153: 544-550.
- Allen MF (1996). The ecology of arbuscular mycorrhizas: a look back into 20<sup>th</sup> century and a peak into the 21<sup>st</sup> century. Mycorrhizal Res. 100: 769-782.
- Barea JM, Andrade G, Bianciotto V, Dowling D, Lohrke S, Bonfante P, O'Gara F, Azcon-Aguilar C (1998). Impact on arbuscular mycorrhiza formation of *Pseudomonas* strains used as inoculants for the biocontrol of soil-borne plant fungal pathogens. Appl. Environ. Microbiol. 64: 2304-2307.
- Barea JM, Pozo MJ, Azcon R, Azcon-Aguilar C (2005). Microbial cooperation in the rhizosphere. J. Exp. Bot. 56: 1761-1778.
- Barker KR, Huisingh D (1970). Histopathological investigations of the antagonistic interaction between *Heterodera glycines* and *Rhizobium japonicum* on soybean. Phytopathology, 60: 1282-1283.
- Benhamou N, Kloepper JW, Quadt-Hallman A, Tuzun S (1996). Induction of a defense-related ultrastructural modifications in pea root tissues inoculated with endophytic bacteria. Plant Physiol. 112: 919-929.
- Bethlenfalvay GJ, Linderman RG (1992). Mycorrhizae in sustainable agriculture. Madison, Wisconsin, ASA Special publication No. 54.
- Bodker L, Kjoller R, Rosendahl S (1998). Effect of phosphate and arbuscular mycorrhizal fungus *Glomus intraradices* on disease severity of root rot of peas (*Pisum sativum*) caused by *Aphanomyces euteiches*. Mycorrhiza, 8: 169-174.
- Broadbent P, Baker KFM, Franks N, Holland J (1977). Effect of *Bacillus* sp. on increased growth of seedlings in steamed and non treated soil. Phytopathology, 67: 1027-1034.
- De Freitas JR, Banerjee MR, Germida JJ (1997). Phosphate solubilizing rhizobacteria enhance the growth and yield buy not phosphorus uptake of canola (*Brassica napus* L.). Biol. Fert. Soils 24: 358-364.
- Demir S, Akkopru A (2005). Use of arbuscular mycorrhizal fungi for biocontrol of soilborne fungal plant pathogens. In: Chincholkar SB,

- Mukerji KG (eds) Biological control of plant diseases. Howarth Press, New York, pp. 17-37.
- Dosppekhnov BA (1984). Field experiment-tation. Statistical procedures. Mir Publishers, Moscow.
- Elbardy M, Taha RM, Eldougdoug KA, Gamal Eldin H (2006). Induction of systemic resistance in faba bean (*Vicia faba* L.) to bean yellow mosaic potyvirus (BYMV) via seed bacterization with plant growth promoting rhizobacteria. *J. Plant Dis. Protec.* 113: 247-251.
- Gamlial A, Katan J (1993). Suppression of major and minor pathogens by fluorescent pseudomonads in solarized and non-solarized soil. *Phytopathology*, 83: 68-75.
- Gianinazzi S, Schuepp H (1994). Impact of arbuscular mycorrhizas on sustainable agriculture and natural eco-systems. Basel, Switzerland: ALS, Birkhauser Verlag.
- Gianinazzi S, Schuepp H, Barea JM, Haselwandter K (2002). Mycorrhizal technology in agriculture: From genes to bioproducts. Basel, Switzerland, Birkhauser Verlag.
- Gupta SC, Namdeo SL (1997). Effect of *Rhizobium*, phosphate solubilizing bacteria and FYM on nodulation, grain yield and quality of chickpea. *Indian J. Pulse Res.* 10: 171-174.
- Huang HC, Erickson RS (2007). Effect of seed treatment with *Rhizobium leguminosarum* on Pythium damping-off, seedling height, root nodulation, root biomass, shoot biomass, and seed yield of pea and lentil. *J. Phytopathol.* 155: 31-37.
- Illmer P, Barbato A, Schinner F (1995). Solubilization of hardly soluble  $AlPO_4$  with P-solubilizing microorganisms. *Soil Biol. Biochem.* 27: 265-270.
- Jeffries P, Gianinazzi S, Perotto S, Turnau K, Barea JM (2003). The contribution of arbuscular mycorrhizal fungi in sustainable maintenance of plant health and soil fertility. *Biol. Fertil. Soil.* 37: 1-16.
- Jetiyanon, K, Tuzun S, Kloepper JW (1997). Lignification, peroxidase and superoxide dismutases as early plant defense reactions associated with PGPR-mediated induced systemic resistance. In: Ogoshi A, Kobayashi K, Homma Y, Kodama F, Kondo N, Akino S (eds.), *Plant Growth-Promoting Rhizobacteria-Present Status and Future Prospects*. Nakanishi Printing, Sapporo, Japan, pp. 265-268.
- Kapulnik Y, Douds Jr DD (2000). Arbuscular mycorrhizas: physiology and function. Dordrecht, The Netherlands: Academic Publishers.
- Kokalis-Burelle N, Vavrina CS, Roskopf EN, Shelby RA (2002). Field evaluation of plant growth-promoting rhizobacteria amended transplant mixes and soil solarization for tomato and pepper production in Florida. *Plant Soil* 238: 257-266.
- Linderman RG (1994). Role of VAM fungi in biocontrol. In: Bethelenfalvay GJ, Linderman RG (eds.) *Mycorrhizae and Plant Health*. APS Press, St Paul, MN, USA, pp. 1-26.
- Meyer JR, Linderman RG (1986). Selective influence on populations of rhizosphere or rhizoplane bacteria and actinomycetes by mycorrhizas formed by *Glomus fasciculatum*. *Soil Biol. Biochem.* 18: 191-196.
- Oostendrop M, Sikora RA (1989). Seed treatment with antagonistic rhizobacteria for the suppression of *Heterodera schachtii* early root infection with sugar beet. *Rev. de Nematol.* 12: 77-83.
- Ozgonen H, Bicici M, Erkilic A (1999). The effect of salicylic acid and endomycorrhizal fungus *G. intraradices* on plant development of tomato and Fusarium wilt caused by *Fusarium oxysporum* f. sp. *lycopersici*. *Turk. J. Agric. For.* 25: 25-29.
- Porter WM (1979). The most probable number method for enumerating infective propagules of vesicular-arbuscular mycorrhizal fungi in soil. *Aust. J. Soil Res.* 17: 515-519.
- Reddy PP (1974). Studies on the action of amino acids on the root-knot nematode *Meloidogyne incognita*. Ph.D. dissertation, University of Agricultural Sciences Bangalore, Indian.
- Richardson AE (2001). Prospects for using soil microorganisms to improve the acquisition of phosphorus by plants. *Aust. J. Plant Physiol.* 28: 897-906.
- Riker AJ, Riker RS (1936). Introduction of Research of Plant diseases. John's Swift Co. Inc. St. Louis, Chicago, New York, Indian polis p. 117.
- Schenck NC, Pervez Y (1990). A manual of vesicular-arbuscular mycorrhizal fungi INVAM 3<sup>rd</sup> edn., University of Florida, Gainesville, Florida, p. 286.
- Shakya MS, Patel MM, Singh VB (2008). Knowledge level of chickpea growers about chickpea production technology. *Indian Res. J. Ext. Edu.* 8: 65-68.
- Siddiqui ZA (2006). PGPR: prospective biocontrol agents of plant pathogens. In: Siddiqui ZA (ed) *PGPR: Biocontrol and Biofertilization*. Springer, The Netherlands, p. 318.
- Siddiqui ZA, Akhtar MS (2007). Biocontrol of a chickpea root-rot disease complex with phosphate solubilizing microorganisms. *J. Plant Pathol.* 89: 66-77.
- Siddiqui ZA, Baghel G, Akhtar MS (2007). Biological control of *Meloidogyne javanica* by *Rhizobium* and plant growth promoting rhizobacteria on lentil. *World J. Microbiol. Biotechnol.* 23: 435-441.
- Siddiqui ZA, Husain SI (1992). Interaction of *Meloidogyne incognita* race 3 *Macrophomina phaseolina* and *Bradyrhizobium* sp. in a root-rot disease complex of chickpea, *Cicer arietinum*. *Fundam. Appl. Nematol.* 15: 491-494.
- Siddiqui ZA, Mahmood I (1995). Biological control of *Heterodera cajani* and *Fusarium udum* by *Bacillus subtilis*, *Bradyrhizobium japonicum* and *Glomus fasciculatum* on pigeon pea. *Fundam. Appl. Nematol.* 18: 559-566.
- Siddiqui ZA, Mahmood I (1999). Role of bacteria in the management of plant parasitic nematodes. A Review. *Bioresour. Technol.* 69: 167-179.
- Smith SE, Read DJ (2008). Mycorrhizal symbiosis, 3<sup>rd</sup> ed. Academic Press, London, UK.
- Suresh CK (1980). Interaction between vesicular arbuscular mycorrhizal and root knot nematode in tomato. M.Sc. (Agric) Thesis, University of Agricultural Sciences Bangalore, Indian.
- Taylor AL, Sasser JN (1978). Biology, Identification and Control of root-knot nematodes (*Meloidogyne* spp.) Dept. Pl. Pathol. Raleigh N.C. North St. Univ. and USAID, North Carolina State Graphics.
- Tilak KVBR (1991). Bacterial fertilizers. Publications and information Division ICAR, New Delhi, pp. 66.
- Tilak KVBR, Ranganayak N, Pal KK, De R, Saxena AK, Nautiyal CS, Mittal S, Tripathi AK, Johri BN (2005). Diversity of plant growth and soil health supporting bacteria. *Curr. Sci.* 89: 136-150.
- Trappe JM (1982). Synoptic keys to genera and species of Zygomycetes Mycorrhizal fungi. *Phytopathology*, 72: 1102-1108.
- van der Heijden MGA, Sanders IR (2002). Mycorrhizal ecology. Berlin, Heidelberg, Springer-Verlag.
- Wei G, Kloepper JW, Tuzun S (1996). Induced systemic resistance to cucumber diseases and increased plant growth by plant growth promoting rhizobacteria under field conditions. *Phytopathology*, 86: 221-224.
- Weller DM (1988). Biological control of soil borne plant pathogens in the rhizosphere with bacteria. *Ann. Rev. Phytopathol.* 26: 379-407.
- Whitelaw MA, Harden TJ, Helyar KR (1999). Phosphate solubilisation in solution culture by the soil fungus *Penicillium radicum*. *Soil Biol. Biochem.* 31: 655-665.