

## Short Communication

# Combined usage of testosterone and nandrolone may cause heart damage

Erdal Tasgin<sup>1\*</sup>, Sefa Lok<sup>2</sup> and Nagehan Demir<sup>3</sup>

<sup>1</sup>Health Science Institute, Selcuk University, 42075, Konya, Turkey.

<sup>2</sup>Physical Education and Sports Academy, Selcuk University, 42075, Konya, Turkey.

<sup>3</sup>Department of Biochemistry, Veterinary Faculty, Selcuk University, 42075, Konya, Turkey.

Accepted 18 March, 2011

The aim of this study is to determine the effects of combined application of testosterone and nandrolone on male rabbits during adolescence period for biochemical values which are indicators of damage to heart, liver and kidney. Seven male New Zealand white rabbits, 60-days old, were used in this study. Testosterone (10 mg/kg) + nandrolone deconoate (10 mg/kg) were injected simultaneously to the rabbits in an intramuscular manner once in 1 week for a period of 12 weeks. Blood samples were taken in 0, 4, 8 and 12 weeks. Serum creatine kinase-MB, lactate dehydrogenase, alkaline phosphatase, alanine aminotransferase, aspartate aminotransferase, gamma-glutamyl-transferase, creatinine and blood urea nitrogen were measured with an auto-analyzer. Combined use of testosterone and nandrolone caused ( $p < 0.05$ ) increase in serum creatine kinase-MB activity, but no statistical changes ( $p > 0.05$ ) were determined in other parameters. As a result, it can be concluded that the use of testosterone plus nandrolone simultaneously during early ages may cause heart damage.

**Key words:** Testosterone, nandrolone, creatine kinase-MB.

## INTRODUCTION

The term testosterone has first been used as sexuality hormone during the 1930s as the combination of 'testo↔testes', 'ster↔sterol' and 'one↔ketone' (Handelsman, 2006; Dotson, 2007). Testosterone is produced in leydig cells (Hartgens, 2004) and it is effective in spermatogenesis, development of secondary sexual characters, release of gonadotropine and stimulation of protein synthesis in an anabolic manner (Wu, 1997; Basaria et al., 2001; Hartgens, 2004). Nandrolone (19-nortetosterone, 17 $\beta$ -hydroxyestr-4en3-one) has been synthesized during the 1950s for the first time. It displays a high affinity to androgen receptors and it has both ana-

bolic and androgenic qualities (Ayotte, 2006; Lumia et al., 2010).

Testosterone and nandrolone are used in the evaluation of the clinical potential of male fertility, kidney failure, aplastic anemia, delayed adolescence, asthenia based on Acquired immune deficiency syndrome (AIDS), protection of immune system after cancer, treatment of hemodialysis patients, osteoporosis, cachexia, trauma or the post-surgery recovery period (Ayotte, 2006; Kerr, 2007).

It is seen that the use of anabolic androgenic steroids (AAS), particularly by young men, have dramatically increased in recent years. It has been used to increase the mass and weight of the body and power to compete, to better their appearances and to improve self-esteem. One of the most common misuses of ASS is testosterone which is an endogen androgen. Testosterone does not only have an endogen effect but it also has an excellent anabolic effect (Lumia et al., 2010). The study that has been conducted has shown that 1 to 2% of young girls and 4 to 6 % of young boys have taken steroids, even if it is just once in a lifetime (Marshall et al., 2009). Misuse by sportsmen cause ethical problems as well (Bhasin et al.,

\*Corresponding author. E-mail: [erdaltasgin@selcuk.edu.tr](mailto:erdaltasgin@selcuk.edu.tr). Tel: +90 332 2233161.

**Abbreviations:** AIDS, Acquired immune deficiency syndrome; AAS, anabolic androgenic steroids; WADA, world anti doping agency; CK-MB, creatine kinase-MB; LDH, lactate dehydrogenase; ALP, alkaline phosphates; ALT, alkaline aminotransferase; AST, aspartate aminotransferase; GGT, gamma glutamyl transferase; BUN, urea.

1996). World Anti Doping Agency (WADA) has stated that sportsmen use testosterone, nandrolone and stanazolol (Handelsman 2006).

AAS may cause bile flow disorders, sensitivity against insulin, disorders in lipid metabolism, hyper-tension, testicular atrophy, gynecomastia, liquid retention, tendon injuries, nose bleeds, hepatic and renal dysfunction, sleep disorders, frequent cold, tachycardia, cardiomegaly, muscle pain, increase in atherosclerosis, acne, increase in sebum production in men, hirsutizm, virilization in women and sudden death (Bhasin et al., 1996; Basaria et al., 2001; Modlinski et al., 2006; Casavant, 2007; Marshall, 2009).

Values of serum biochemistry provide us information about organ and system damages in a creature. Creatine kinase-MB (CK-MB) (Dickerman et al., 1999) increases in heart damage, whereas lactate dehydrogenase (LDH) increases in myocardium, liver and musculoskeletal diseases (Turgut, 2000). Liver damage indicators are alkaline phosphates (ALP), alkaline aminotransferase (ALT), aspartate aminotransferase (AST) and gamma glutamyl transferase (GGT), whereas urea (BUN) and creatinine are kidney damage indicators (Bartley, 1989; Finco, 1989; Kramer, 1989). No information regarding the effects of long term combined use of testosterone and nandrolone during youth on serum biochemical values have been obtained in the researches that have been made.

The aim of this study is to examine effects of applying testosterone and nandrolone on male rabbits in adolescence on the damage indicators regarding heart, liver and kidney.

## MATERIALS AND METHODS

Seven New Zealand white rabbit of 60 days old (1000 to 13000 g, SÜDAM, Konya, Turkey) were used in the study. Animals were fed *ad libitum* during the experiment. The research has been confirmed by the Veterinary Faculty Ethic committee. Once in a week, 10 mg/kg testosterone (Sustanon® 250 amp, Organon ilaçları A.S, Istanbul, Turkey) + 10 mg/kg nandrolone deaconate (DECA 1000 flk. Gen-shi labs.co. Osaka, Japan) was injected into the animals in an intramuscular manner. Blood was taken from auricular artery in weeks 0, 4, 8 and 12. Levels of CK-MB, LDH, ALP, ALT, AST, GGT, creatinine and BUN were determined in auto-analyzer (IL-300 Instrumentation laboratory Milano, Italy) from the serums that have been obtained.

### Statistical analysis

Results of the study were evaluated by analysis of variance (ANOVA) and Tukey test (SPSS 10.0).  $P < 0.05$  value has been accepted as significant in statistical terms.

## RESULTS AND DISCUSSION

Effect of testosterone + nandrolone application on adolescent male rabbits on serum biochemical values is

given in Table 1. Testosterone and nandrolone application causes increase in CK-MB level ( $P < 0.05$ ). Changes in LDH, ALP, ALT, AST, GGT, CREAT and BUN levels have been determined to be within the reference range.

Testosterone and its derivatives, which have been used for treatment since 1940s, are being misused by some sportsmen (Kerr et al., 2007; Handelsman, 2006). Testosterone and nandrolone application on adolescent rabbits in this study caused an increase ( $P < 0.05$ ) in serum CK-MB level (Table 1).

Similarly, while an increase is observed in CK-MB with the male rats to which testosterone has been applied (Lok et al., 2010), no increase has been observed in CK-MB levels with female rats (Tasgin et al., 2010). Serum CK-MB levels are being used in the diagnosis of cardiovascular system disorders (Brancaccio et al., 2006). It has not been observed in literature reviews that combined use of testosterone and nandrolone has effects on serum CK-MB levels. Combined use of testosterone and nandrolone may cause heart damage especially for adolescent males.

LDH levels in the current study have been determined to be above ( $P < 0.05$ ) the reference range during first sampling time (Table 1). In this study, in which enzyme levels from childhood to advanced ages have been studied, it is observed that LDH levels have decreased for young and adolescent people (Lockitch et al., 1988). It may be said also that the reason why LDH levels are high at the beginning is because of the early ages. Also, the high ( $P < 0.05$ ) levels determined for GGT in the day 0 may result from the fact that it has been high during early ages, as it is the situation in LDH.

Serum ALP, ALT, AST and GGT levels increases in liver damage (Turgut, 2000; Er and Yazar, 2010). Although it has been stated that these values increased in the studies which reported the effect of AAS on liver damage (Hartgens et al., 2004; Vieira et al., 2008; Lok et al., 2010; Tasgin et al., 2010), these values have been determined to be within reference range in the current study (Altunok et al., 2002; Hernandez-Divers, 2004; Yazar et al., 2004) (Table 1). The result of this study may be as a result of the applied medicine, dose and animal breed differences.

Although there have been some changes in creatinine and BUN levels in the current study (Table 1), these differences have been determined to be within reference range (Elmas et al., 2006; Elmas et al., 2008).

## Conclusion

As a conclusion, it can be said that long term testosterone and nandrolone use in early ages may cause heart damage in young ages. However, this study should be supported especially with histopathological researches.

**Table 1.** Effects of testosterone + nandrolone on serum organ damage markers (mean±SE).

Weeks	0	4	8	12	Reference range
CK-MB (U/L)	3114±327 <sup>b</sup>	1895±343 <sup>b</sup>	2695±592 <sup>b</sup>	4737±166 <sup>a</sup>	-----
LDH (U/L)	1188±122 <sup>a</sup>	355±30.3 <sup>c</sup>	226±17.6 <sup>c</sup>	754±101 <sup>b</sup>	34-500
ALP (U/L)	183±15.7 <sup>a</sup>	156±14.2 <sup>ab</sup>	109±12.0 <sup>bc</sup>	60.8±7.33 <sup>c</sup>	16-231
ALT (U/L)	51.8±3.37 <sup>a</sup>	58.0±6.55 <sup>a</sup>	81.2±15.4 <sup>a</sup>	52.3±5.13 <sup>a</sup>	22-87
AST (U/L)	59.6±7.53 <sup>a</sup>	30.1±5.32 <sup>b</sup>	29.2±6.16 <sup>b</sup>	58.2±7.54 <sup>a</sup>	14-113
GGT (U/L)	16.2±1.72 <sup>a</sup>	12.3±2.22 <sup>ab</sup>	9.62±0.77 <sup>b</sup>	8.87±0.93 <sup>b</sup>	0-7
Creatinine (mg/dL)	0.64±0.06 <sup>c</sup>	0.77±0.05 <sup>bc</sup>	0.88±0.04 <sup>ab</sup>	1.04±0.06 <sup>a</sup>	0.6-1.97
BUN (mg/dL)	32.7±1.88 <sup>b</sup>	40.6±4.40 <sup>ab</sup>	40.8±1.39 <sup>ab</sup>	46.2±3.49 <sup>a</sup>	17-48

Different letters on the same line are statistically significant (Tukey test, P < 0.05).

## REFERENCES

- Altunok V, Yazar E, Elmas M, Tras B, Bas AL, Col R (2002). Investigation of hematological and biochemical side effects of tilmicosin in healthy New Zealand rabbits. *J. Vet. Med. B.* 49: 68-70.
- Ayotte C (2006). Significance Of 19-norandrosterone in athletes' urine samples. *Br. J. Sports. Med.* 40: 25-29.
- Bartley JC (1989). Lipid Metabolism and Its Diseases, In; *Clinical Biochemistry of Domestic animals*. Editor: Kaneko JJ. 4th Edition Academic Press, London, UK, pp.106-135.
- Basaria S, Wahlstrom JT, Dobs AS (2001). Anabolic-androgenic Steroid Therapy in the Treatment of Chronic Diseases. *J. Clin. Endocrinol. Metab.* 86: 5108-5117.
- Bhasin S, Storer TW, Berman N, Callegari C, Clevenger B, Phillips J, Bunnell TJ, Tricker R, Shirazi A, Casaburi R (1996). The effects of supraphysiologic doses of testosterone On muscle size and strength in normal men. *N. Engl. J. Med.* pp. 335:1-7.
- Brancaccio P, Maffulli N, Limongelli FM (2006). Monitoring of serum enzymes in sport. *Br J sports Med.* 40:96-97. <http://bmb.oxfordjournals.org/cgi/reprint/ldm014v1>
- Casavant MJ, Blake K, Griffith J, Yates A, Copley LM (2007). Consequences of use on anabolic androgenic steroids. *Pediatr. Clin. N. Am.* 54: 677-690.
- Dickerman RD, Petrusi RM, Zachariah NY, Dufour DR, McConathy WJ (1999). Anabolic steroid-induced hepatotoxicity: Is it overstad?. *Clin. J. Spor. Med.* 9: 34-19.
- Dotson JL, Brown RT (2007). the history of the development of anabolic-androgenic steroids. *Pediatr. Clin. N. Am.* 54: 761-769.
- Elmas M, Yazar E, Uney K, Er A, Tras B (2008). Pharmacokinetics of enrofloxacin and flunixin meglumine and interactions between both drugs after intravenous co-administration in healthy and endotoxaemic rabbits. *Vet. J.* 177: 418-424.
- Elmas M, Yazar E, Uney K, Karabacak A (2006). Pharmacokinetics of flunixin after Intravenous administration in healthy and endotoxaemic rabbits. *Vet. Res. Commun.* 30: 73-81.
- Er A, Yazar E (2010). Effects of macrolide antibiotics on blood inflammatory mediators and organ damage markers in lipopolysaccharide-induced pulmonary damage rats. *Eurasian J. Vet. Sci.* 26: 7-13.
- Finco DR (1989). Kidney Function, in; *Clinical Biochemistry of Domestic animals*. Editor: Kaneko JJ. 4th Edition Academic Press, London, UK, pp. 524-537.
- Handelsman DJ (2006). Testosterone: use, misuse and abuse. *Med. J. Aust.* 185: 436-439.
- Hartgens F, Kuipers H (2004). Effects of androgenic-anabolic steroids in athletes. *Sport Med.* 34: 513-554.
- Hernandez-Divers SJ (2004). Rabbits, in; *Exotic Animal Formulary*, Ed; Carpenter JW. Third Edition, Elsevier Saunders, Pennsylvania, USA, pp. 411-446.
- Kerr JM, Congeni JA (2007). Anabolic-androgenic steroids: use and abuse in pediatric patients. *Pediatr. Clin. N. Am.* 54: 771-785.
- Kramer JW (1989). *Clinical Enzymology*, in; *Clinical Biochemistry of Domestic Animals*. Editor: Kaneko JJ. 4th Edition Academic Press, London, UK, pp. 352-360.
- Lockitch G, Halstead AC, Albersheim S, Maccallum C, Quigley G (1988). Age-and Sex-Specific pediatric reference intervals for biochemistry analytes as measured with the ektachem-700 analyzer. *Clin. Chem.* 34: 1922-1625.
- Lok S, Tasgin E, Demir N, Ozdemir M (2010). Long term used testosterone may cause heart and liver damage. *J. Anim. Vet. Adv.* 9: 2343-2345.
- Lumia AR, McGinnis MY (2010). Impact of anabolic androgenic steroids on adolescent males. *Physiol. Behav.* 100: 199-204.
- Marshall-Grandisnik S, Green R, Brenu EW, Weatherby RP (2009). Anabolic androgenic steroids effects on the immune system: a review. *Cent. Eur. J. Biol.* 4: 19-33.
- Modlinski R, Fields KB (2006). The effect of anabolic steroids on the gastrointestinal system, kidneys and adrenal glands. *Curr. Sports Med. Rep.* 5: 104-910
- Tasgin E, Lok S, Demir N, Ozdemir M (2010). The Effect of Testosterone Used in Sportsmen on Routine Biochemical Parameters. *J. Anim. Vet. Adv.* 9: 2038-2040.
- Turgut K (2000). *Veteriner Klinik Laboratuar Teşhis. Bahçivanlar basım sanayi.* ISBN 975-94595-1-5: 180-198-371.
- Vieira RP, Franca RF, Damaceno-Rodrigues NR, Dolhnikoff M, Caldini EG, Carvalho CRF, Ribeiro W (2008). Dose-dependent hepatic response to subchronic administration of nandrolone decanoate. *Med. Sci. Sport Exer.* 40: 842-847.
- Yazar E, Col R, Konyalioglu S, Birdane YS, Elmas M, Bas AL (2004). Effect of vitamin and prednisolone on biochemical and heametological parameters in endotoxaemic New Zealand white rabbits. *Bull. Vet. Inst. Pulawy.* 48: 105-108.
- Wu FCW (1997). Endocrine aspects of anabolic steroids. *Clin. Chem.* 43: 1289-1292.