

## Review

# Role of inhibitors and biodegradable material in mitigation of nitrogen losses from fertilized lands

Nasima Junejo<sup>1\*</sup>, Khanif M.Y.<sup>1</sup>, Hanfi M.M.<sup>2</sup>, W. M. Z. Wan Yunus<sup>3</sup> and Dharejo K. A<sup>1</sup>

<sup>1</sup>Department of Land Management, Faculty of Agriculture, Universiti Putra Malaysia, Serdang, Selangor, Malaysia.

<sup>2</sup>Institute of Tropical Agriculture, Universiti Putra Malaysia, Serdang, Selangor, Malaysia.

<sup>3</sup>Department of Chemistry, Faculty of Science, Universiti Putra Malaysia, Serdang, Selangor, Malaysia.

Accepted 25 January, 2011

Loss of N, occurring mainly through NH<sub>3</sub> volatilization, biological denitrification, and NO<sub>3</sub><sup>-</sup> leaching, has both economic and environmental implications. Therefore, the economic benefits of reduced environmental pollution and future damage to our environment as a result of the use of urease inhibitors are of higher significance to the production of gains over the long-term. The literature review discussed the role, effectiveness and mechanism of urease inhibitors as well as coating of urea to reduce ammonia volatilization loss from urea fertilizer. Coating with urease inhibitors can improve the bioavailability of N, resulting in increased dry matter yield and N uptake. Such increases result from delayed urea hydrolysis by urease inhibitors and coating materials. The value of inhibitors in mitigating N loss would depend on their rate of biodegradation and persistence in soils. Previous studies has proven that micronutrients could be competent urease inhibitors, but the information about their effects on soil, nutrient uptake of plants and dry matter yield is poorly documented. Some nutrients such as Cu and Zn as urease inhibitors and natural biodegradable material such as agar, gelatin and palm stearin were found effective in reduction of nitrogen losses from nitrogen fertilizer especially from urea fertilizer. The article deals with the importance of use of above mentioned materials and micronutrient.

**Key words:** Nitrogen, urea, fertilizer, inhibitors, biodegradation, land, nitrogen loss.

## INTRODUCTION

Nitrogen is a mobile nutrient, thus, it easily moves and losses quickly from soils. Agricultural soils need a continuous supply of N from various sources to maintain productivity. Most agricultural land is deficient in nitrogen (N) for the growth of crops.

There are a variety of nitrogenous fertilizers available in the market; however, urea consumption is 38%, which is higher than other nitrogenous fertilizers due to the relatively low manufacturing cost and high concentration of N (Bouwman et al., 1997). Urea is a widespread, major


form of N produced and used by agricultural world. Urea consumption ranged from about 67,146,000 tons during 2008 and 2009 (IFA, 2008) (Figure 1); however, there is concern about the efficiency of using urea-N for agricultural crops because farmers' practices in Asia commonly result in recoveries of <40% of the applied N (Malhi et al., 2001).

The addition of urea to soil not only increases plant productivity but also results in increased nitrate (NO<sub>3</sub><sup>-</sup>) leaching as well as the release of gaseous N, such as ammonia (NH<sub>3</sub>) and nitrous oxide (N<sub>2</sub>O), which are detrimental to the environment. Urea volatilization is a function of several factors which includes:

\*Corresponding author. E-mail: [neelum85@hotmail.com](mailto:neelum85@hotmail.com).

**Abbreviations:** N, Nitrogen; NIs, nitrification inhibitors; UIs, urease inhibitors; CRF, controlled release fertilizers; SRF, slow release fertilizers.

1. Hydrolysis rate
2. Equilibrium reactions
3. NH<sub>3</sub> exchange between the soil and the atmosphere
4. Exchange between NH<sub>4</sub> in solution and exchange sites in the soil.



**Figure 1.** Global consumption of mineral fertilizers (IFA, statistics 2008 to 2009).

Gaseous emissions of N, via ammonia (NH<sub>3</sub>) volatilization and denitrification, have been identified as the dominant mechanisms of fertilizer N loss in many different agricultural systems (Peoples et al., 1995). There has been a concern to improve the effectiveness of urea fertilizers via the use of coatings or modifications using various urease inhibitors and synthetic materials to mitigate N loss.

## NITROGEN DYNAMICS IN SOIL


It is important to understand the process of N transformation in soil and the biochemical mechanism involved in alleviating N loss through the use of inhibitors (Singh et al., 2004).

### Nitrogen mineralization

In nitrogen mineralization, plant in unavailable organic forms is converted into plant-available inorganic forms through the activity of soil micro-organisms. The process includes two reactions, which include the following: ammonization and ammonification (Figure 3).


Ammonization is a microbial process, in which micro-organisms first hydrolyze macromolecules of organic N compounds. In ammonification, micro-organisms convert amines and amino acids into NH<sub>4</sub><sup>+</sup> ions. For example, urea (CO(NH<sub>2</sub>)<sub>2</sub>) in fertilizer undergoes an ammonification reaction, releasing NH<sub>4</sub><sup>+</sup> ions. This process is also known as 'urea hydrolysis' and is carried out in the presence of urease enzymes in soil. Urease is an enzyme produced by some microbial and plant species. The ammonification

process also releases hydroxyl (OH<sup>-</sup>) ions and increases soil pH around urea granules, resulting in alkaline conditions. The high NH<sub>4</sub><sup>+</sup> ion concentration and the elevated pH during the ammonification reaction provide ideal conditions for ammonia volatilization to occur (Singh et al., 2008).




### Ammonia volatilization loss

The total ammonical N in a soil solution comprises of two principle forms of N (NH<sub>4</sub> and NH<sub>3</sub>). When NH<sub>3</sub> is present in a soil solution, the physical process of NH<sub>3</sub> transfer across a water surface (in contact with the atmosphere) is described by the simple reaction:




Some of the most important factors regulating NH<sub>3</sub> loss are the concentration of ammonical N, the temperature and the pH of the soil solution or irrigation water; moreover, all three variables markedly affect the partial pressure of NH<sub>3</sub> (Singh et al., 2008).

The pH in particular, affects the equilibrium between ammonium and NH<sub>3</sub> so that the relative concentration of NH<sub>3</sub> increases from 0.1 to 1, 10 and 50% as the pH changes from 6 to 7, 8 and 9, respectively (Freney et al., 1983). The high soil pH (> 7) favors ammonia volatilization. In the case of urea application, the initial increase



**Figure 2.** World production of urea (IFA, statistics 2008 to 2009).

in soil pH through the ammonification process results in  $\text{NH}_3$  volatilization (Figure 2).



The lost  $\text{NH}_3$  is consequently deposited on land or in water, which causes eutrophication and acidification of natural ecosystems on a regional scale (Sommer and Hutchings, 2001).

As temperatures rises, the relative proportions of  $\text{NH}_3$  to ammonium present at a given pH increases, while the solubility of  $\text{NH}_3$  in water decreases (Figure 4), which increases the diffusion of  $\text{NH}_3$  through the soil solution and affects the rate of microbial transformations. The rate of urea hydrolysis in soil and  $\text{NH}_3$  volatilization are greatly affected by soil temperature (Vlek and Carter, 1983; Carmona et al., 1990). Wind speed is another major factor that controls volatilization (via its effect on mixing in the liquid phase) as well as the rate of transport of  $\text{NH}_3$  away from the air-water or air-soil interface (Freney et al., 1981; Denmead et al., 1982; Fillery et al., 1984).

Other variables that influence  $\text{NH}_3$  volatilization include the pH-buffer capacity, cation exchange capacity of the soil, levels of urease activity, and the availability of moisture, soil texture, nitrification rate, and the presence of plants or plant residues (Freney and Black, 1988).

Large losses of  $\text{NH}_3$  from applied fertilizer have been detected from soil, floodwater and irrigation water in many agro ecosystems (Freney and Black, 1988; Peoples et al., 1995). Losses of  $\text{NH}_3$  measured from different upland and lowland cropping systems have ranged from negligible amounts to > 50% of the fertilizer N applied, depending upon fertilizer practices and environmental conditions (Bacon et al., 1986; Keller and Mengel, 1986; Black et al., 1989; Freney et al., 1992). In flooded rice,  $\text{NH}_3$  volatilization can account for 20 to > 80% of the total N lost from fertilizer sources (Simpson et al., 1984; De Datta et al., 1989; Freney et al., 1990; Mosier et al., 1989;

Zhu, 1992). Ammonia emission can be reduced by reduction of ammonical N in soil solutions through microbial processes, such as leaching, cation exchange complex, immobilization and nitrification.

### Immobilization

Immobilization is a microbial process, in which the plant-available in  $\text{NH}_4^+$  and  $\text{NO}_3^-$  ions are converted to plant unavailable organic N. For instance, the addition of carbon (C)-rich substances in arable soils promotes immobilization and reduces N availability to plants. When the N content of the decomposing organic matter is small, a reduction in N availability occurs because microbes become deprived of N and compete with plants for available N in soil. Thus, the addition of organic matter with high C:N ratios causes immobilization of soil N by micro-organisms, thereby decreasing the amount of plant-available soil N (Singh et al., 2008).

### Nitrification and denitrification

Nitrification is the oxidation of ammonium to nitrate. In soil, the process is mediated by some bacteria. The effect of soil water content on N transformations can be observed by its effect on oxygen diffusion.

The nitrification process is dependent upon the movement of ammonium into the oxidized zone to be processed by nitrifying organisms. The transport of  $\text{NH}_4^+$  by diffusion is affected by both the organic matter status and the cation exchange capacity of the soil as well as the presence of reduced iron and manganese, the bulk density and the rate of nitrification in the oxidized soil layer (Peoples et al., 1995).

The mechanism of denitrification involves the transformation of nitrate and nitrite to  $\text{NO}$ ,  $\text{N}_2\text{O}$  and  $\text{N}_2$ . Soil factors that significantly influence denitrification are oxygen (which is controlled primarily by soil water content), nitrate concentration, pH, temperature and organic carbon (Peoples et al., 1995). Soil water tends to moderate oxygen diffusion in soil, and denitrification occurs only when the soil water content is > 60% of the air-filled pore space in flooded rice fields (Linn and Doran, 1984). Soil organic matter provides energy for denitrified growth as well as supplies of protons and electrons for the reduction process. In crops, the effect of application timing of fertilizer on N loss by denitrification may be more related to precipitation rate.

### Ammonia fixation

The influence of the  $\text{NH}_4^+$  fixation capacity of soil on gains and losses has not been fully ascertained; however, there is some evidence that the quantities of fixed  $\text{NH}_4^+$  do not change greatly over long periods of time when concurrent




Figure 3. Schematic representation of nitrogen transformation (Singh et al., 2008).





Figure 4. Effect of pH and temperature in a soil solution of total ammoniacal N (Sommer et al., 2002).

changes on soil organic N have occurred (Jaiyebo and Bouldin, 1997). Fixation can influence  $\text{NH}_4\text{-N}$  transformation by reducing  $\text{NH}_3$  loss and nitrification rates. In a  $^{15}\text{N}$  balance study, Kowalenko (1998) found that 59% of the  $^{15}\text{NH}_4$  applied to the 0 to 15 cm layer of an ammonium-fixed clay loam soil was immediately fixed by clay minerals.

### Leaching losses

Leaching is very common in Malaysian soils due to the high annual rate of rainfall. Losses occur mainly as  $\text{NO}_3$ , the movement of which is closely related to water movement. Major losses of N occur when soil  $\text{NO}_3$  content is high and water movement is large.



**Figure 5.** Ribbon diagram, Urease enzyme, the blue green and red presenting the sub units (Stefano et al., 1998).

Leaching losses are also strongly influenced by seasonal effects, such as water and temperature. In humid regions, mineralization rates are low in water, but leaching of residual  $\text{NO}_3^-$  from the previous season often occurs. In the spring,  $\text{NO}_3^-$  accumulates as nitrification rates increase, and N fertilizers are applied. Nitrate leaching is least likely to take place during the summer. Leaching not only depletes  $\text{NO}_3^-$  N but also takes away clay, soil and organic matter, which leads to low chemical and low plant available water reserves (Allison, 1993).

#### Temporal ammonia loss pattern from urea fertilizer

When urea is applied on the soil, it absorbs water and is hydrolyzed, producing ammoniacal nitrogen and bicarbonate ( $\text{HCO}_3^-$ ). The rate of hydrolysis is related to the amount of soil moisture and temperature. The  $\text{NH}_3$  loss

rate decreases after 5 to 10 days due to a reduction in  $\text{NH}_3$  concentration through various processes, such as volatilization, leaching, denitrification and immobilization (Black et al., 1985; Haynes and Williams, 1992). The maximum  $\text{NH}_3$  volatilization loss rates may occur within 1 to 10 days after application, depending on the soil and environmental conditions (Stevens et al., 1989). The pattern of  $\text{NH}_3$  volatilization from urea is affected by applied forms of urea despite whether it is applied in pellets, in solution, or crushed to a fine powder because the release and hydrolysis of urea into the soil solution is different. From urea applied in different forms, such as powder or in solution, the ammonia volatilization occurs earlier than after application of urea. This delay in emission from urea is related to hydrolysis of the applied urea.

#### UREA HYDROLYSIS

Urea, an odorless, white, crystalline water-soluble solid, is a diamide of carbonic acid. In soil, urea decomposes enzymatically to  $\text{CO}_2$  and  $\text{NH}_4$ . The urease enzyme (Figure 5) present in soil, plants and plant litter is responsible for the urea hydrolysis process (Frenay and Black, 1988).

The urease active site consists of two nickel (II) atoms linked by a carbamate bridge. Two imidazole N atoms bind to each Ni atom, with a carboxylate group and water molecule filling the remaining coordination sites of the metal ion. The capacity to hydrolyse urea varies from 17 to 70% for soil bacteria and from 78 to 98% for soil fungi (Lloyd and Sheaffe, 1973). However, soil urease is considered to be of microbial origin, and there is indication that some soil urease activity may be derived from plants (Frankenberg and Tabatabai, 1982).

#### Factors affecting urea hydrolysis

Urease activity of soils is associated with organic matter, soil pH, temperature, moisture and urea concentrations (O'Toole et al., 1982; Reynolds et al., 1985; Kissel and Cabrera, 1988). Urease activity decreases when the organic matter content of soil decreases with depth (Bremner and Mulvaney, 1978; Mulvaney and Bremner, 1981).

#### Effect of pH on urea hydrolysis

Urease enzymes are responsible for the hydrolysis of urea fertilizer (into  $\text{NH}_3$  and  $\text{CO}_2$ ) applied to the soil, with the concomitant rise in soil pH (Byrnes and Amberger, 1989) resulting in rapid N loss to the atmosphere through  $\text{NH}_3$  volatilization (Fillery and Frenay, 1988).

The rate of urea hydrolysis was measured under laboratory conditions using soils with a range of pH values (from 2.2 to 8.0). It has been reported that soil pH

**Table 1.** Fatty acid content of Palm stearin.

Type of fatty acid	Colour	Content (%)
Lauric saturated C12		48.2
Myristic saturated C14		16.2
Palmitic saturated C16		8.4
Capric saturated C10		3.4
Caprylic saturated C8		3.3
Stearic saturated C18		2.5
Oleic monounsaturated C18		15.3
Linoleic polyunsaturated C18		2.3
Other/Unknown		0.4

Red, saturated; orange, monounsaturated; blue, polyunsaturated (Hyes and Khosla, 1992).

increases as the rate of urea hydrolysis increases, and the highest rate of hydrolysis was observed at pH 8.0 (Ahmed et al., 2006) Table 1.

### Effect of temperature on urea hydrolysis

Hydrolysis of urea is temperature dependent and increases with increasing soil temperature over the range from 0 to 40°C (Vlek and Carter, 1983); however, slight hydrolysis has been detected at sub-zero temperatures as well (Bremner and Mulvaney, 1978). It was found that as temperature increases, the rate of dissolution of urea in water increases. Urea hydrolysis is accelerated by increasing temperatures in part because the rate of urea diffusion is positively correlated with temperature.

Moreover, urease is an extra cellular enzyme produced by microorganisms, and high temperatures can increase both microorganism growth and urease production. Moyo et al. (1989) found that increasing temperature (from 5 to 45°C) greatly increases urease activity. Urease activity reaches a maximum between 60 and 70°C.

### Effect of urea concentration on urea hydrolysis

Urea hydrolysis is also influenced by urea concentration, soil water and soil pH. The optimum pH for urea hydrolysis is between 6.0 to 7.0 (Kissel and Cabrera, 1988). When urea is applied to the soil, the concentration may range from 0.01 to 10 M. There is a possible existence of two reactions, one with high affinity and one with low affinity for urea. The high affinity reaction is responsible for most of the urea hydrolysis at urea concentrations lower than 0.1 M. In general, the urea N concentration, at which both affinity enzyme reactions contribute equally, is 0.5 M. Cabrera et al. (1991) found that when urea concentration is > 6 M, the rate of urea hydrolysis decreases, which is possibly due to enzyme

denaturizing (Kistiakowsky and Rosenberg, 1952).

### Effect of moisture on urea hydrolysis

Soil moisture content is one of the important factors affecting urea hydrolysis because urea is a hygroscopic material and absorbs water through contact with liquid water as well as from water –vapor in the air (Wahl et al., 2006). After application of urea to soil, the process of dissolution of urea initiates immediately because urea is a compound that is highly soluble in water (about 1080 gL<sup>-1</sup> can be dissolved at 20°C) (Wahl et al., 2006).

### Strategies to reduce ammonia volatilization loss from urea fertilizer

There are several strategies that have been adopted to reduce ammonia volatilization loss, such as fertilizer placement and time of applications as well as amendment of acidic materials and the use of polymers and other chemicals as coatings.

### Modification in placement, rate and method of fertilizer application

Placement of urea in the lower zone is related to soil properties, and this strategy is not very useful for sandy and peat soils because in sandy soil, urea diffuses to the surface very fast and is lost (Blaise et al., 1996), and in peat soils, it may cause toxicity to plant roots due to high concentrations of NH<sub>4</sub>.

The placement of urea is directly related to the rate of application in calcareous soils. High application rates should be placed at a greater depth than low application rates of fertilizer due to heavy N losses in these soils (Fenn and Miyamoto, 1981; Ismail et al., 1991).

In contrast on acidic soils, NH<sub>3</sub> emission has increased due to the high rate of fertilizer applications because the addition of high rates of fertilizer increases the fertilizer micro site pH, which results in a high accumulation of NH<sub>4</sub> on soil surface (Rachhpal and Nye, 1984).

Methods of fertilizer application have also been considered to reduce ammonia loss. It has been suggested that application of urea in split doses with band applications instead of broadcasting could be helpful in reducing ammonia loss (Rachhpal and Nye, 1984).

### Amendment and coating of urea with chemicals

Amendments or applications of urea mixed with inorganic salts and acidic materials, such as humic acid, CaCl<sub>2</sub> and KCl, may reduce NH<sub>3</sub> emission significantly (Fenn et al.,



1982; Ahmed, 2008).

### Controlled/Slow release fertilizers (CRF/SRF)

Controlled release fertilizers (CRF) have been shown to increase plant yields in some but not all studies (Shoji et al., 2001; Chen et al., 2008), and in some studies, reduced N<sub>2</sub>O emissions have been observed (Shoji and Kanno, 1994). Controlled release fertilizers are used only to a small extent in agriculture which is mainly due to their high cost (Chen et al., 2008).

CRFs, in which a physical barrier is used to reduce dissolution rate, are commonly prepared by coating (or matrix formation) of water soluble, granular plant nutrients with low permeability and hydrophobic membranes (Shaviv and Mikkelsen, 1993). The nutrient in CRF is made to release its contents gradually and to coincide with the requirements of the plant. This fertilizer can be physically prepared from granules of the soluble fertilizers by coating them with materials, which reduce their dissolution rate, and it also causes an increase in the fertilizer efficiency, reduces soil toxicity, minimizes the potential negative effects associated with over dosage and reduces the frequency of application (Tomaszewska et al., 2002).

The control of fertilizer release keeps its' concentration at effective levels in soil and releases nutrients when they are most needed by the plant. As a result, maximum utilization of the fertilizer from the plant system reduces nutrient loss, decreases fertilizer rate and applications, prevents seedling damage, and provides full protection of the environment.

By adopting specific fertilizer formulations and techniques to improve fertilizer efficiency, it should be possible to provide sufficient N in a single application to satisfy plants' needs. If this could be done, any N loss event would be small because of the limited amount of N in the substrate.

Various slow-release forms of N have been suggested, including the following: coated fertilizers (Shoji et al., 1991), complex organic N compounds that are much less soluble in water than urea (Oertli, 1980; Allen, 1984) and urea super granules (Youngdahl et al., 1986).

The effect of slow-release forms on levels of soil mineral N as well as the recovery of fertilizer N have been assessed, and the use of these products has generally decreased total loss of fertilizer N (Chauhan and Mishra, 1989; Malhi and Nyborg, 1992). Sulfur coated urea has the potential to reduce ammonia emission, but it may not be very significant in all environments. For example, in a study by Black et al. (1985), NH<sub>3</sub> emission from sulfur-coated urea was lower than from untreated urea (10.1% as compared with 12.6%). Sediments of pyrite (FeS) have been shown to reduce emission from urea by 54% (Blaise et al., 1996), and a 10% reduction in hydrolysis rates was reported by amending urea with thiosulfate under drying conditions (Black et al., 1985).

## USE OF INHIBITORS

Urea N use efficiency could be enhanced if the hydrolysis of urea to ammonium by soil urease could be retarded by the use of urease inhibitors. UIs (Urease Inhibitors) slow the conversion of urea to NH<sub>4</sub><sup>+</sup> by inhibiting the urease enzyme, which reduces the NH<sub>4</sub><sup>+</sup> concentration in the soil solution and hence, lowers the potential for NH<sub>3</sub> volatilization and seedling damage; slow urea hydrolysis allows more time for it to release nitrogen from the fertilizer micro site.

### Mechanism of Inhibition of Urease

A number of chemicals have been tested as potential inhibitors of soil urease activity for use with urea fertilizers. These chemicals can be divided into three groups according to their structures and binding modes with urease, including the following; (1) reactive organic or inorganic compounds, (2) chelating compounds that cause inhibition due to complex formation with one of the Ni atoms at the active site of urease, and (3) competitive inhibitors that resemble urea molecules and bind to the active site of the urease enzyme (Amtul et al., 2002).

### Bioavailability of N with inhibitors

Plants take N both as NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup>. Synchronization of plant N uptake with released NH<sub>4</sub><sup>+</sup> or NO<sub>3</sub><sup>-</sup> by controlling the rate at which urea (in applied urea) is hydrolyzed to NH<sub>4</sub><sup>+</sup> and its subsequent oxidation to NO<sub>3</sub><sup>-</sup> as well as the temporary rise in soil pH is critical to minimize gaseous and leaching losses of N. Most plants prefer NO<sub>3</sub><sup>-</sup> over NH<sub>4</sub><sup>+</sup>; however, the rate of uptake of NH<sub>4</sub><sup>+</sup> is often found to be greater than that of NO<sub>3</sub><sup>-</sup>, especially at low temperatures. Since plant roots can absorb both NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> ions, ammonification and nitrification processes markedly influence N absorption efficiency by plants, mainly by controlling the concentrations of these ions in soil solution. It has often been observed that while UIs decrease the concentration of NH<sub>4</sub><sup>+</sup> ions, Nis (nitrification inhibitors) increase the concentration of NH<sub>4</sub><sup>+</sup> ions and decrease NO<sub>3</sub><sup>-</sup> ions. Urease inhibitors slow down urea hydrolysis and, thus, keep N in urea form (Bremner et al., 1991; Wang et al., 1991; Watson et al., 1994b).

In the presence of sufficient sun light as a source of energy, N assimilatory enzymes (NO<sub>3</sub><sup>-</sup> reductase) in plants rapidly reduce NO<sub>3</sub><sup>-</sup> to NH<sub>3</sub>, which is then assimilated into glutamine and glutamate. Glutamine, glutamate and organic acids produced from carbohydrate metabolism then serve as N donors in the biosynthesis of amines, amides and essentially all amino acids and nucleic acids. The amino acids serve as building blocks for the synthesis of proteins; thus, NO<sub>3</sub><sup>-</sup> reduction occurs both in aerial portions (shoots and leaves) as well as in the roots of plants, with most reduction occurring in shoots. The

relative importance of these two sites of  $\text{NO}_3^-$  conversion is considered most important.

It is obvious that the rate of  $\text{NH}_4^+$  assimilation is faster than that of  $\text{NO}_3^-$  because the former is directly incorporated into organic compounds. To maintain a charge balance, plant uptake of  $\text{NH}_4^+$  and  $\text{NO}_3^-$  affects the pH of rhizosphere by releasing either hydrogen ( $\text{H}^+$ ) or hydroxyl ( $\text{OH}^-$ ) ions. The release of such ions by plants also affects the uptake of other anions and cations. For example,  $\text{NH}_4^+$  absorption reduces the uptake of cations such as calcium ( $\text{Ca}^{++}$ ), magnesium ( $\text{Mg}^{++}$ ) and potassium ( $\text{K}^+$ ) and increases the uptake of anions, such as phosphate and sulphate.  $\text{NO}_3^-$  uptake reduces the absorption of anions.

Many studies have been carried out in different agricultural systems that have stated that the application of N fertilizer with UIs or NIs improves the bioavailability of N, resulting in increased plant production and N uptake (Watson et al., 1998; Xu et al., 2002; Zaman et al., 2005). Such increases are always attributed to slow urea hydrolysis by UIs. Increase of 20, 17 and 15% in pasture production have been reported from urea applied with UI, NI and UI+NI, respectively (Zaman et al., 2005). Increase in dry matter production have been obtained from urea applied with UIs. This high, dry matter yield and N uptake with inhibitors can be attributed to the retention of applied N as mineral N or organic N in the soil profile, which consequently becomes available for plants.

### Effect of inhibitors on N loss

Many research studies have confirmed that inhibitors are effective in delaying the conversion of either urea to  $\text{NH}_4^+$  (UIs) or  $\text{NH}_4^+$  to  $\text{NO}_3^-$  (NIs). Most research has shown that the application of UIs to soil with urea reduces  $\text{NH}_3$  volatilization, while the application of NIs reduces  $\text{NO}_3^-$  leaching as well as  $\text{N}_2\text{O}$  emissions. Some studies also have shown that NIs increases  $\text{NH}_3$  volatilization (Davies and Williams, 1995; Nastri et al., 2000).

Treating urea with urease inhibitors reduces  $\text{NH}_3$  loss from surface applications (Bremner et al., 1991; Clay et al., 1990; Carmona et al., 1990). Laboratory (Carmona et al., 1990; Vittori-Antisari et al., 1996) and field studies (Rawluk et al., 2001; Watson et al., 1994a) have reported increased inhibition of urease activity with an increasing rate of UI, which followed the law of diminishing returns (Watson et al., 1994b).

The environmental concern over the use of toxic and hazardous chemicals is increasing worldwide. There are many such chemicals polluting the environment, with their damage depending on their exposure and persistence in the ecosystem as well as the characteristics of the affected organisms (Hashim et al., 1992).

In general, soil microorganisms readily degrade natural polymers because they contain chemical bonds that may be broken down through enzymatic hydrolysis in soil. Synthetic polymers are not degradable and are more

resistant to biological breakdown; however recently, there have been some degradable synthetic polymers introduced (Devassine et al., 2002).

### Use of micronutrients as urease inhibitors

Some micronutrients (Cu and Zn) were investigated as competent urease inhibitors. These micronutrients have a crucial value in the growth of plant. Before describing their inhibitory effects, it is important to understand their role in plant production and status in soil. Copper is an essential element for all crops, and it influences both carbohydrate and nitrogen metabolism in plants (Mengle and Kirby, 1987). Cu is present in soils as oxides, carbonates, silicates and sulfides. Copper availability decreases in soil solutions due to adsorption of Cu in soil exchange complexes, as well as due to chemical fixation of Cu as sulfides (Lea et al., 1993). Soil Cu availability depends on soil pH; it is lower in the alkaline range and higher in the acidic range. Most Cu deficiencies occur in sandy soils, which have a high pH, or on soils with a 5% higher organic matter content (Johne, 1983).

Zinc is an essential nutrient that plays an important role in plant growth. It is an important part of protein that works as a synthesizer of sugars and starch (Marschner, 1995; Sharma, 2006). Zinc is the most common crop micronutrient deficiency, particularly in high-pH soils. Notably, 50% of cultivated soils in the world are classed as Zn-deficient (Graham et al., 1992; White and Zasoski, 1999; Cakmak, 2002, 2004; Alloway, 2004).

Cu and Zn were firstly tested as urease inhibitors in 1971 by Bremner and Douglas. The authors observed that  $\text{NH}_3$  volatilization loss from urea, that is surface applied to soil can be mitigated by 30.3 and 24.6%, and N-use efficiency improved by 28.3 and 23.9% if urea was amended with  $\text{CuSO}_4$  (Reddy and Sharma, 2000). In their study, copper amended urea was applied to reduce urea volatilization loss and to provide Cu as a micronutrient. This approach has been shown to be effective (Leong, 2002). The amendment of urea with  $\text{ZnSO}_4$  can improve the agronomic efficiency of fertilizer by 11% (Purakayastha and Katyal, 1998).

### FATTY ACIDS

Natural materials, such as natural oils and fatty acids, have been used to reduce nitrification loss and avoid environmental pollution. Some natural products, such as karanjin, neem, tea waste and mint oil, can reduce nitrification significantly (Sahrawat, 1987; Patra et al., 2009). It was observed that the constituents of fatty acid are linolenic acid, and lanoleic acid, which have 98% inhibitory effects on nitrosamine successfully reduce nitrification loss (Subbarao, 2008).

Palm stearin is one of the preferred natural and cost effective fatty acids used by many manufacturers in



formulations of consistent fats, such as margarines, bakery fats and frying fats. It was also found that different demethylated oils and terpenes are potential retardants of urease activity and nitrification (Patra et al., 2009); palm stearin could be one of them. In addition, the coating by palm stearin enables Cu and urea to come together at the micro site.

### Use of agar and gelatin as biodegradable polymers

Agar, the sea weed well known in commerce, is derived from several species of aquatic plants and is made principally from algae. The algae grow on rocks along the coast and are collected by divers between May and October, with the best months for collection being July and August. The weed is dried on the shore, and during the process, it is partially bleached and is then ready to be manufactured into the commercial product. The chemical formula of agar is  $C_{11}H_{16}O_{10}$ , which is composed of crude protein, carbohydrates and crude fiber (Whittaker, 1910). Gelatin is the secondary product of collagen and is subjected to degradative processes. The properties of gelatin are similar to rigid-chain synthetic polymers.

Gelatin macromolecules assume, at high temperatures, the conformation of a statistical coil. Under specific conditions (temperature, solvent and pH), gelatin macromolecules can display flexibility sufficient to realize a wide variety of conformations (Kozlov, 1983). Gelatin mostly is a constituent of the shells of pharmaceutical capsules to make them easy to swallow.

These materials (agar and gelatin) were considered to be natural polymers (Ward and Courts, 1977) but were never used as a coating for fertilizer.

In the presented study, these materials have been used as adhesive agents of urease inhibitors, and a source of coating to release urea-N slowly from fertilizer.

### CONCLUSION

Loss of N, occurring mainly through  $NH_3$  volatilization, biological denitrification and  $NO_3^-$  leaching has both economic and environmental implications. Therefore, the economic benefits of reduced environmental pollution and future damage to our environment as a result of the use of urease inhibitors are of higher significance to the production of gains over the long-term.

The literature review discussed the role, effectiveness and mechanism of urease inhibitors as well as coating of urea to reduce ammonia volatilization loss from urea fertilizer. Coating with urease inhibitors can improve the bioavailability of N, resulting in increased dry matter yield and N uptake. Such increases result from delayed urea hydrolysis by urease inhibitors and coating materials.

The value of inhibitors in mitigating N loss would depend on their rate of biodegradation and persistence in

soils. Previous studies has proven that micronutrients could be competent urease inhibitors, but the information about their effects on soil, nutrient uptake of plants and dry matter yield is poorly documented

### REFERENCES

- Ahmed OH, Husni AMH (2006). Reducing ammonia loss from urea and improving soil-exchangeable ammonium retention through mixing triple superphosphate, humic acid and zeolite. *Soil Use Manage.* 22: 315-319.
- Ahmed OH, Hussain, AH Ahmad MH (2008). Ammonia volatilization and ammonium accumulation from urea mixed with zeolite and triple super phosphate. *ACTA Agr. Scand. B-S P* 58: 182-186.
- Alloway BJ (2004). Zinc in soils and crop nutrition. International Zinc Association Belgium, Brussels.
- Amtul Z, Atta-ur-Rahman, Siddiqui RA and Choudhary MI (2002). Chemistry and mechanism of urease inhibition. *Current Medicinal Chemistry* 9: 1323-1348.
- Bacon PE, Houlst EH and McGarity JW (1986). Ammonia volatilization from fertilizers applied to irrigated wheat soils. *Fert. Res.* 10: 27-42
- Black AS, Sherlock RR, Smith NP, Cameron KC, Goh KM (1985). Effects of form of nitrogen, season, and urea application rate on ammonia volatilization from pastures. *New Zeal. J. Agr. Res.* 28: 469-474.
- Blaise D, Tyagi PC, Khola OPS (1996) Ammonia volatilisation from urea as affected by the addition of iron pyrites and method of application. *Nutr. Cyc.I Agroecosys.* 46: 97-101.
- Bouwman AF, Lee DS, Asman WAH, Dentener FJ, Van Der Hoek KW, Olivier JGJ. (1997) A global high-resolution emission inventory for ammonia. *Global Biogeochem. Cy.* 11: 561-587.
- Bremner JM, Mulvaney RL (1978) Urease activity in soils. In "Soil Enzymes" R. G. Burns, ed., pp. 149-196. Academic Press, London.
- Bremner JM, McCarty GW, Higuchi T (1991) Persistence of the inhibitory effects of phosphoramides on urea hydrolysis in soils. *Commun. Soil Sci. Plan.* 22: 1519-1526.
- Byrnes BH, Amberger A (1989) "Fate of broadcast urea in a flooded soil when treated with N-(n-butyl) thiophosphoric triamide, a urease inhibitor. *Fert. Res.* 18: 221-231.
- Cabrera RI (1997). Comparative evaluation of nitrogen release patterns from controlled release fertilizers by nitrogen leaching analysis. *Hort. Sci.* 32(4): 669-673.
- Cakmak I (2002). Plant nutrition research: priorities to meet human needs for food in sustainable ways. *Plant Soil* 247: 3-24.
- Cakmak I (2004). Identification and correction of widespread zinc deficiency in Turkey – a success story (a NATO-Science for Stability Project). *Proceedings of the International Fertiliser Society* 552: 1-26.
- Carmona G, Christianson CB, Byrnes BH (1990) Temperature and low concentration effects of the urease inhibitor N-(n-butyl) thiophosphoric triamide (Nbtpt) on ammonia volatilization from urea. *Soil Biol. Biochem.* 22: 933-937.
- Chauhan HS, Mishra B (1989) Ammonia volatilization from a flooded rice field fertilized with amended urea materials. *Fert. Res.* 19: 57-63.
- Chen D (2008). Enhanced efficiency fertilizers for agricultural sustainability and environmental quality in Australia. IFA cross roads Asia Pacific report, University of Melbourne Australia.
- Clay DE, Malzer GL, Anderson JL (1990). Ammonia volatilization from urea as influenced by soil temperature, soil water content, and nitrification and hydrolysis inhibitors. *Soil Sci. Soc. Am. J.* 54: 263-266.
- De Datta SK, Trevitt ACF, Freney JR, Obcemea WN, Real JG and Simpson JR (1989). Measuring nitrogen losses from lowland rice using bulk aerodynamic and nitrogen-15 balance methods. *Soil Sci. Soc. Am. J.* 53: 1275-1281.
- Denmead OT, Freney JR, Simpson JR (1982) Dynamics of ammonia volatilization during furrow irrigation of maize. *Soil Sci. Soc. Am. J.* 46: 149-155.
- Devassine M, Henry F, Guein P and Briand X (2002). Coating of fertilizer by degradable polymeric. *J. Pharmaceutics.* 22: 399-404.
- Fenn LB, Miyamoto S (1981) Ammonia loss and associated reactions of

- urea in calcareous soils. *Soil Sci. Soc. Am. J.* 45: 537–540.
- Fernery JR, Black AS (1988) Importance of ammonia volatilization as a loss process. In *Advances in Nitrogen Cycling in Agricultural Ecosystems*. J. R. Wilson, ed. pp. 156-173. CAB International, Wallingford.
- Fernery JR, Simpson JR, Denmead OT (1981). Ammonia volatilization. In: *Terrestrial Nitrogen Cycles: Processes, ecosystem strategies and management impacts*. Eds. F.E. Clark, T Roswell. pp. 291-302.
- Fillery IRP, Simpson JR, De Datta SK (1984) Influence of field environment and fertilizer management on ammonia loss from flooded rice. *Soil Sci. Soc. Am. J.* 48: 914-920.
- Frankenberger WT, Tabatabai MA (1982). Amidase and urease activities in plants. *Plant Soil* 64: 153-166.
- Freney JR, Trevitt ACF, Muirhead WA, Denmead OT, Simpson JR, Obcemea WN (1988). Effect of water depth on ammonia loss from lowland rice. *Fert. Res.* 16: 97-107.
- Freney JR, Black AS (1988) Importance of ammonia volatilization as a loss process. In: *Advances in Nitrogen Cycling in Agricultural Ecosystems*, J.R. Wilson (ed.). C.A.B International, Wallingford, pp. 156-173.
- Freney JR, Denmead OT, Watanabe I, Craswell ET (1981). Ammonia and nitrous oxide losses following applications of ammonium sulfate to flooded rice. *Aust. J. Agr. Res.* 32: 37-45.
- Freney JR, Denmead OT, Wood AW, Saffigna PG, Chapman LS, Ham GJ, Hurney AP, Stewart RL (1992). Factors controlling ammonia loss from trash covered sugarcane fields fertilized with urea. *Fert. Res.* 31: 341-349.
- Freney JR, Keerthisinghe DG, Chaiwanakupt P, Phongpan S (1993). Use of urease inhibitors to reduce ammonia loss following application of urea to flooded rice fields. *Proceedings of the Plant Nutrition Colloquium Perth Australia*, pp. 491-493.
- Freney JR, Simpson JR, Denmead OT (1983). Volatilization of ammonia. In: *Gaseous Loss of Nitrogen from Plant-Soil Systems*, J.R. Freney, and J.R. Simpson, (eds.). Martinus Nijhoff/Dr. W. Junk Publishers, The Hague, pp. 1-32.
- Freney JR, Trevitt ACF, De Datta SK, Obcemea WN, Real JG (1990). The interdependence of ammonia volatilization and denitrification as nitrogen loss processes in flooded rice fields in the Philippines. *Bio. Fert. Soils* 9: 31-36.
- Graham RD, Ascher JS, Hynes SC (1992). Selecting zinc-efficient cereal genotypes for soils of low zinc status. *Plant Soil* 146: 241–250.
- Hashim MA, Kuldandai J, Hassan RS (1992). Biodegradability of Branched Alkylbenzene Sulphonates. *J. Chem. Technol. Biot.* 54: 207 - 214
- Haynes RJ and Williams PH (1992). Changes in soil solution composition and pH in urine affected areas of pasture. *Soil Sci.* 43: 323–334.
- IFA Statistics (2008). International Fertilizer Industry Association: <http://www.fertilizer.org/ifa/statistics/STATSIND/tablen.asp>
- Ismail KM, Wheaton FW, Douglass LW, Potts W (1991). Modeling ammonia volatilization from loamy sand soil treated with liquid urea. *Trans. ASAE* 34: 756–763.
- Jaiyebo EO, Bouldin DR (1997). Influence of fertilizer and manure additions and crop rotations on nonexchangeable ammonium content of two soils. *Soil Sci.* 103: 16-22
- Johan JM (1983). Impact of acid deposition on micronutrient cycling in agro ecosystem. *Env.Exp. Botany* 23: 243-249
- Keller GD, Mengel DB (1986). Ammonia volatilization from nitrogen fertilizers surface applied to no-till corn. *Soil Sci. Soc. Am. J.* 50: 1060-1063.
- Kissel DE, Cabrera ML (1988) Factors affecting urea hydrolysis. In "Ammonia volatilization from urea fertilizers" (B. R. Bock and D. E. Kissel, eds.), Bulletin Y-206, pp. 53-66. National Fertilizer Development Center, Tennessee Valley Authority, Muscle Shoals, Alabama.
- Kowalenko CG (1998). Nitrogen transformation and transport over 17 months in field fallow microplots using <sup>15</sup>N. *Can. J. Soil Sci.* 58: 69-76.
- Kozlov PV (1983). The structure and properties of solid gelatin and the principles of their modification. *Polymer* 24: 651-666.
- Lea P (1993). Nitrogen metabolism. In "Plant Biochemistry and Molecular Biology" (P. J. Lea and R. C. Leegood, eds.), pp. 155-180. John Wiley and Sons, New York.
- Leong TK (2002). The development of cooper-coated urea for rice production. A Master Thesis, UPM, Serdang, Malaysia
- Linn DM, Doran JW (1984). Effect of water-filled pore space on carbon dioxide and nitrous oxide production in tilled and nontilled soils. *Soil Sci. Soc. of Am. J.* 48: 1267-1272.
- Lloyd AB, Sheaffe MJ (1973) Urease activity in soils. *Plant Soil* 39: 71-80.
- Malhi SS, Grant CA, Johnston AM, Gill KS (2001) Nitrogen fertilization management for no-till cereal production in the Canadian Great Plains: a review. *Soil. Till. Res.* 60: 101–122.
- Malhi SS, Nyborg M (1992). Recovery of nitrogen by spring barley from ammonium nitrate, urea and sulphur-coated urea as affected by time and method of application. *Fer. Res.* 32: 19-25.
- Marschner H (1995). *Mineral Nutrition of Higher Plants*, 2nd Edn. Academic Press, San Diego
- Mengel K, Kirkby EA (1987) *Principles of Plant Nutrition*. 4 th Edition. International Potash Institute, IPI, Bern, Switzerland. pp. 685.
- Mosier AR, Kroeze C (2000). Potential impact of global atmospheric N<sub>2</sub>O budget of the increased nitrogen input required to meet future global food demands. *Chemosphere-Global Change Science* 2: 465-473.
- Mulvaney RL, Bremner JM (1981). Control of urea transformations in soils. In: Paul EA, Ladd JN (eds). *Soil biochemistry*, New York, Marcel Dekker. pp.153-196.
- Nastri A, Toderi G, Bernati E, Govi G (2000). Ammonia volatilization and yield response from urea applied to wheat with urease (NBPT) and nitrification (DCD) inhibitors. *Agrochimica* 44: 231-239.
- O'Toole P, Morgan MA, McAleese DM. (1982). Effects of soil properties, temperature and urea concentration on patterns and rates of urea hydrolysis in some Irish soils. *Irish J. Agr. Res.* 21: 185-197.
- Patra DD, Kiran U, Chand S (2009). Use of urea coated with natural products to inhibit urea hydrolysis and nitrification in soil. *Biol. Fert. Soils* 45: 617-621.
- Peoples MB, Freney JR, Mosier AR (1995). Minimizing gaseous losses of nitrogen. In: *Nitrogen Fertilization in the Environment*, P.E. Bacon (ed.). Marcel Dekker, Inc. New York, pp. 565-602.
- Purakayastha TJ, Katyal JC (1998). Evaluation of compacted urea fertilizers prepared with acid and non-acid producing chemical additives in three soils varying in pH and cation exchange capacity. *NH<sub>3</sub> volatilization. Nutr. Cyc. Agroecosys.* 51: 117–121.
- Rachhpal S, Nye PH (1984). The effect of soil pH and high urea concentration on urease activity in soil. *Eur. J. Soil Sci.* 35: 519-527.
- Reddy D, Sharma KL (2000). Effect of amending urea fertilizer with chemical additives on ammonia volatilization loss and nitrogen-use efficiency. *Biol. Fert. Soils* 32: 24–27.
- Reynolds CM, Wolf DC, Armbruster JA (1985). Factors related to urea hydrolysis in soils. *Soil Sci. Soc. Am. J.* 49: 104-108.
- Sahrawat KL, Keeney DR, Adams SS (1987). Ability of nitrapyrin, dicyandiamide and acetylene to retard nitrification in a mineral and an organic soil. *Plant Soil* 101: 179-182.
- Sharma CP (2006). *Plant micronutrients*. Enfield, NH, USA: Science Publishers.
- Shaviv AE, Mikkelsen RL (1993). Slow release fertilizers to increase efficiency of nutrient use and minimize environmental degradation: A review. *Fert. Res.* 35: 1-12.
- Shoji S, Delgado J, Mosier A, Miura Y (2001). Use of controlled release fertilizers and nitrification inhibitors to increase nitrogen use efficiency and to conserve air and water quality. *Commun. Soil Sci. Plan.* 32: 1051-1070.
- Shoji S, Gandeza AT, Kimura K (1991). Simulation of crop response to polyolefin-coated urea. II. Nitrogen uptake by corn. *Soil Sci. Soc. Am. J.* 55: 1468-1473.
- Singh J, Bolan NS, Saggarr S, Zaman M (2008). The role of inhibitors in controlling the bioavailability and losses of nitrogen. In: *Chemicals bioavailability in terrestrial environment*. Amsterdam, The Netherlands, Elsevier. pp. 329-362.
- Singh J, Bolan NS, Saggarr S, Zaman M (2008). The role of inhibitors in controlling the bioavailability and losses of nitrogen. In: *Chemicals bioavailability in terrestrial environment*, Amsterdam, The Netherlands, Elsevier. pp. 329-362.
- Singh J, Saggarr S, Bolan N (2004). Mitigating gaseous losses of N from

- pasture soil with urease and nitrification inhibitors. In Proceedings of the Australia New Zealand SuperSoil 2004 Conference, Sydney, Australia. 6–9 December 2004. <http://www.regional.org.au/au/asssi>
- Sommer SG, Hutchings NJ (2001) Ammonia emissions from field applied manure and its reduction. *Eur. J. Agron.* 15: 1-15.
- Subbarao GV, Nakahara K, Ishikawa T (2008). Free fatty acids from the pasture grass *Brachiaria humidicola* and one of their methyl esters as inhibitors of nitrification. *Plant Soil* 313: 89-99
- Tomaszewska M, Jarosiewicz A, Karakulski K (2002). Physical and chemical characteristics of polymer coating in CRF formulation. *Desalination.* 146: 319-323
- Vittori AL, Marzadori C, Gioacchini P, Ricci S, Gessa C (1996). Effects of the urease inhibitor N-(n-butyl) thiophosphoric triamide in low concentrations on ammonia volatilization and evolution of mineral nitrogen. *Biol. Fert. Soils.* 22: 196-201.
- Vlek PLG, Carter MF (1983). The effect of soil environment and fertilizer modifications on the rate of urea hydrolysis. *Soil Sci.* 136: 56-63.
- Wahl MR, Kirsch U, Brockel ST, Bottlinger M (2006). Caking of urea prills. *Chem. Eng. Technol.* 29: 674–678.
- Wang Z, Cleemput V, Liantie O, Baert L (1991). Effect of organic matter and urease inhibitors on urea hydrolysis and immobilization of urea nitrogen in an alkaline Soil. *Biol. Fert. Soils.* 11: 101-104.
- Ward AG, Courts A (1977). *The Science and technology of gelatin.* Academic Press, London.
- Watson CJ (2000). Urease activity and inhibition - principles and practice. In: *Int. Fert. Soc. Proc.* 454.
- Watson CJ, Poland P, Miller H, Allen MBD, Garrett MK, Christianson CB (1994b). Agronomic assessment and N-15 recovery of urea amended with the urease inhibitor Nbtpt (N-(n-Butyl) thiophosphoric triamide) for temperate grassland. *Plant Soil* 161: 167-177.
- White JG, Zasoski RJ (1999). Mapping soil micronutrients. *Field Crop. Res.* 60: 11–26.
- Whittaker BA (1904). The source, manufacture and composition of commercial agar agar. *J. Am. Public Health Assoc.* 24: 137-141
- Xu XK, Boeckx P, Van CO, Zhou LK (2002). Urease and nitrification inhibitors to reduce emissions of CH<sub>4</sub> and N<sub>2</sub>O in rice production. *Nutr. Cyc. I Agroecosys.* 64: 203-211.
- Youngdahl LJ, Lipin MS, Craswell ET (1986) New developments in nitrogen fertilizers for rice. *Fert. Res.* 9: 149-160
- Zaman M, Nguyen L, Blennerhassett JD, Quin BF (2005). Increasing the utilisation of urea fertilisers by pasture. In "Developments in Fertiliser Application Technologies and Nutrient Management" (L. D. Currie and J. A. Hanly, eds.), vol. 18, pp. 276-284, Fertiliser and Lime Research Centre, Massey University, Palmerston North.
- Zhu SL (1992). Efficient management of nitrogen fertilizers for flooded rice in relation to nitrogen transformation in flooded soils. *Pedosphere.* 2: 97-114.