

Full Length Research Paper

Haplotype analysis of Ethiopian bread wheat (*Triticum aestivum*) cultivars and elite lines for yellow rust resistance genes using linked molecular markers

Woubit Dawit^{1*}, Ayele Badebo², Bekele Hunde³, Daniel Kassa³ and M. S. Röder⁴

¹College of Agriculture and Veterinary Science, Ambo University, P. O. Box 19, Ambo, Ethiopia.

²International Maize and Wheat Improvement Center (CIMMYT), P. O. Box 5689, Addis Ababa, Ethiopia.

³Kulumsa Agricultural Research Center, P. O. Box 489, Kulumsa, Ethiopia.

⁴Leibniz Institute of Plant Genetics and Crop Plant Research (IPK), Corrensstr. 3, 06466 Gatersleben, Germany.

Received 5 June, 2018; Accepted 30 August, 2018

Yellow rust, caused by *Puccinia striiformis* f. sp. *tritici*, is one of the most important diseases of wheat in Ethiopia. Poor knowledge of resistance genes deployed in Ethiopian wheat cultivars is one major factor for recurrent epidemics of rust diseases. Molecular marker based gene identification showed the presence of *Yr1*, *Yr4*, *Yr6*, *Yr7*, *Yr9*, *Yr10*, *Yr17* and *Yr18* in various frequencies, whereas *Yr8* was not detected in any of the tested 74 bread wheat genotypes. *Yr7* was the most frequent (74%) followed by *Yr6* (56%) and *Yr18* (37%) whereas *Yr1* and *Yr4* were detected in lower frequency (7 and 14%), respectively. The contribution of each *Yr* gene was evaluated in yellow rust differential lines possessing various genes. The differential lines carrying *Yr9* and *Yr8* had the highest average coefficient of infection (ACI) value (83 each) followed by *Yr6* and *Yr7* with ACI values of 82 and 80, respectively. The lowest ACI value (46.4) was exhibited on a differential line that carried *Yr4*. The number of *Yr* genes identified from the tested genotypes varied from 0 to 5. The ACI value exhibited by varieties possessing the maximum number of five resistance genes was 42. The majority (26 genotypes representing 35%) of the genotypes possessed three genes with an average coefficient of infection of 42. Pyramiding of the identified genes does not provide sufficient protection against yellow rust in Ethiopia. Thus, there is urgent need for searching for more effective resistance genes to be incorporated in Ethiopian bread wheat cultivars.

Key words: Average coefficient of infection (ACI), bread wheat, molecular markers, wheat genotypes, *Yr* genes.

INTRODUCTION

Wheat is one of the staple food crops cultivated by 5 million small scale farmers in Ethiopia. It is ranked fourth in land coverage and total production after teff, maize and sorghum (CSA, 2017). Yellow rust, caused by *Puccinia striiformis* f. sp. *tritici*, is one of the most important

diseases of wheat that incurs 30 to 69% yield loss in Ethiopia (Badebo et al., 2008). Growing yellow rust resistant cultivars is widely recognized as the most eco-friendly and economically feasible approach. Since the 1970s, more than 100 bread wheat varieties have been

*Corresponding author. E-mail: woubit123@yahoo.com. Tel: 251911658400.

Table 1. Details of molecular markers and primer sequences used for *Yr*- gene screening.

Gene	Marker/Primer	Type	Distance to gene (cM)	Primer sequence	Anneling Temperature (°C)	Fragment size (bp)	Reference
<i>Yr1</i>	BU099658	EST-SSR	-	TGAATCAATACAAGGACGCC GTCACAACACTGGCAACCC	54	≥206	Hasancebi et al. (2014)
<i>Yr1</i>	Stm673acag	STM	1.1	TAACTCACAACAGTTCTGGTCGT ACACACACACACAGAGAGAG	56	120-124	Bansal et al. (2009)
<i>Yr4</i>	barc75	SSR	5.3	AGGGTTACAGTTTGCTCTTTAC CCCGACGACCTATCTATACTTCTCTA	50	132-139	Bansal et al. (2009)
<i>Yr5/18</i>	Wms120	SSR	-	GTGAAGCAGACCCAACAC GACGGCTGCGACGTAGAG	46	472	Singh (1992a)
<i>Yr6</i>	Wmc76	SSR	-	CTTCAGAGCCTCTTTCTCTACA CTGCTTCACTTGCTGATCTTTG	51	256	Li and Niu (2007)
<i>Yr6</i>	Wmc276	SSR	-	GACATGTGCACCAGAATAGC AGAAGAACTATTCGACTCCT	47	292	Li and Niu (2007)
<i>Yr7</i>	Xgwm526	SSR	5.3	CAATAGTTCTGTGAGAGCTGCG CCAACCCAAATACACATTCTCA	48	212, 217	Cabuk et al. (2011)
<i>Yr8</i>	Xgwm157	SSR	-	GTCGTCGCGGTAAGCTTG' GAGTGAACACACGAGGCTTG	60	120	-
<i>Yr9</i>	Xgwm582	SSR	-	AAGCACTACGAAAATATGAC TCTTAAGGGGTGTTATCATA	55	-	Cabuk et al. (2011)
<i>Yr9</i>	iag95	Gene specific	-	CTCTGTGGATAGTTACTTGATCGA CCTAGAACATGCATGGCTGTTACA	55	1100	Mago et al. (2002)
<i>Yr9</i>	P6M12-P	STS	0.35	GTACTAGTATCCAGAGGTCACAAG CAGACAAACAGAGTACGGGC	56	56 250/350	Mago et al. (2005)
<i>Yr10</i>	Yr10F & Yr10R	Gene specific	-	TCAAAGACATCAAGAGCCGC TGGCCTACATGAACTCTGGAT	51	543	Liu et al. (2014)
<i>Yr15</i>	Xgwm273	SSR	-	ATTGGACGGACAGATGCTTT AGCAGTGAGGAAGGGGATC	-	167	Yaniv et al. (2015)

Table 1. Contd.

Yr17	VENTRUP-F, LN2	Gene specific	-	AGGGGCTACTGACCAAGGCT TGCAGCTACAGCAGTATGTACACAAAA			Qamar et al. (2008)
Yr17	URIC/LN2	<i>T. ventricosum</i> chromosom specific	-	GGTCGCCCTGGCTTGACCT TGCAGCTACAGCAGTATGTACACAAAA	64	275/285	Jia et al. (2011)
Yr17	SC-385	SCAR	3.4	CTGAATACAAACAGCAAACCAG ACAGAAAGTGATCATTCCATC	50	385	Jia et al. (2011)
Yr18	Wms295	SSR	-	GTGAAGCAGACCCACAACAC GACGGCTGCGACGTAGAG	-	-	Cabuk et al. (2011)
Yr18	L34DINT9F, L34PLUSR	Gene specific	-	TTGATGAAACCAGTTTTTTTCTA GCCATTTAACATAATCATGATGGA	58	517	Krattinger et al. (2009)
Yr16/Yr24	Xgwm18	SSR	-	GTGGTATTTAGGTGGAGTTGTTTTA CGGAGGAGCAGTAAGGAAGG	60	200	Ahmad et al. (2015)
Yr15 & Yr26	Xgwm11	SSR	-	GGATAGTCAGACAATTCTTGTG GTGAATTGTGCTTGTATGCTTCC	55	215	Ma et al. (2001)
Yr65	Xgwm11	SSR	2.1	GATAGTCAGACAATTCTTGTG GTGAATTGTGCTTGTATGCTTCC	50	213 /202	Cheng et al. (2014)
Yr29	Xgwm44	SSR	3.6/10.9	GGTCTTCTGGGCTTTGATCCTG TGTTGCTAGGGACCCGTAGTGG	57	260	Rosewarne et al. (2006)
Yr32	Xgwms198	SSR	-	CACGCTGCCATCACTTTTAC TTGAAGTGGTCATTGTTGCT	51	160	Eriksen (2004)
Yr33	Xgwm111	SSR	-	TCTGTAGGCTCTCTCCGACTG ACCTGATCAGATCCCCTCG	52	184/206	Zahravi et al. (2003)
Yr35	Xgwm508	SSR	7.5	GTTATAGTAGCATATAATGGCC GTGCTGCCATGATATTT	51	135	Dadkhodaie et al. (2011)
Yr46	Xgwm165	SSR	0.4	TGCAGTGGTCAGATGTTTCC CTTTTCTTTCAGATTGCGCC	449	~236 bp	Herrera-Foessel et al. (2014)
Yr46/Yr62	Xgwm192	SSR	0.4/2	GGTTTTCTTTCAGATTGCGC CGTTGTCTAATCTTGCCCTGC	54/51	~130/222	Herrera-Foessel et al. (2014); Liu et al. (2014)

Table 1. Contd.

Yr 49	Xgwm161	SSR	1	GATCGAGTGATGGCAGATGG TGTGAATTACTTGGACGTGG	57	145/154	McIntosh et al. (2016)
Yr52	Barc182	SSR	1.2	CCATGGCCAACAGCTCAAGGTCTC CGCAAAACCGCATCAGGGAAGCACCA AT	59	≥75	Ren et al. (2012)
Yr57	Xgwm389	SSR	2	ATCATGTGATCTCCTTGACG TGCCATGCACATTAGCAGAT	51	117-128	Radhawa et al. (2015)
Yr62	Xgwm251	SSR	3.3	GGTTTTCTTTCAGATTGCGC CGTTGTCTAATCTTGCCCTTGC	51	133	Liu et al. (2014)

released in Ethiopia. However, the resistance is not lasting long due to the development of new virulent races of the pathogen. Poor knowledge of resistant genes deployed in Ethiopian wheat cultivars is one of the major reasons for recurrent epidemics of rust diseases. Those cultivars may be protected by the same resistance gene(s) or combinations, which could increase selection pressure for the corresponding virulent races. To date, more than 67 yellow rust resistant genes have been reported in wheat and its wild relatives (McIntosh et al., 2013). Most of these genes condition race-specific resistance and many have been overcome by the emergence of new races.

The most effective strategy for protecting wheat from rust is to deploy cultivars with combinations of different resistance genes. For this, information on the resistance genes in major cultivars is of paramount importance. The traditional way of gene postulation requires multi-pathotypes testing in which a host cultivar is evaluated against a collection of isolates carrying different avirulence/virulence gene combinations (pathotypes) on the basis of phenotypic expression in the form of infection types (ITs). As an alternative to gene

postulation, presence of resistance genes can be determined by testing host cultivars with molecular markers linked to resistance genes. This approach overcomes gene interactions and plant stage dependent gene expression problems associated with traditional gene postulation (Vanzetti et al., 2011).

Currently, there have been advances in development and mapping of molecular markers that are diagnostic for major *Yr* genes (Mago et al., 2002; Eriksen, 2004; Li and Niu, 2007; Bansal et al., 2009; Krattinger et al., 2009; Jia et al., 2011; Cabuk et al., 2011; Liu et al., 2014; Yaniv et al., 2015; Ahmad et al., 2015). These markers provide an important tool to plant breeders for marker assisted wheat breeding and also for pyramiding resistance genes in the absence of distinguishable rust virulences (Kaur et al., 2008). This study was conducted to identify yellow rust resistant gene (s) that are present in commercial bread wheat cultivars and elite lines using molecular markers linked to *Yr* genes, assess the effectiveness of the identified *Yr* genes to the prevailing races under field conditions in Ethiopia, and to evaluate genetic variations among the

wheat cultivars and elite lines.

MATERIALS AND METHODS

Plant materials

A total of 58 commercial bread wheat cultivars, 16 elite lines and 15 reference lines with known *Yr* genes and a negative check, *Morocco*, with unknown *Yr* genes and *PBW345* were included in this study. All the wheat materials were obtained from the Ethiopian Institute of Agricultural Research (EIAR), KARC, Ethiopia. The list of wheat genotypes together with their additional information is shown in Supplementary Table 1.

Field testing

All wheat genotypes were evaluated for yellow rust reaction in three locations, namely, Mararo, Arsi Robe and Kulumsa under natural infection in 2016 and 2017. The materials were sown in two rows of 1 m length with 0.20 m spacing between rows. For scoring yellow rust severity in the field, the modified Cobb Scale (Peterson et al., 1948) was used to determine the percentage of tissue infected with rust. The host response to infection in the field was scored using "R" or resistant (small uredinia surrounded by

chlorosis or necrosis); “MR” or moderately resistant (medium sized uredinia surrounded by chlorosis or necrosis); “MS” or moderately susceptible (medium large compatible uredinia without chlorosis and necrosis); and “S” or susceptible (large, compatible uredinia without chlorosis and necrosis). Disease severity and host response data were combined in a single value called the coefficient of infection (CI). The average coefficient of infection (ACI) was used for host response: immune = 0.0, R = 0.2, MR = 0.4, MS = 0.8, and S = 1.0.

Molecular markers

Two closely linked (usually flanking) markers of each of the major *Yr* genes were chosen to identify its presence/absence in wheat genotypes/materials, except a few genes for which only one closely linked marker was reported. A total of 31 markers that are linked to yellow rust resistant genes were used to identify *Yr* genes from the test materials as well as for genetic diversity analysis. Primer names, forward and reverse primer sequences, expected amplified fragment size in base pairs, and annealing temperature are shown in Table 1.

DNA extraction

DNA was extracted from 2 weeks old fresh leaves harvested and pooled from five seedlings of each accession and stored cryogenically at -80°C. Extraction from frozen leaves was performed using the modified cetyl trimethylammonium bromide (CTAB) method described by Doyle and Doyle (1990).

PCR amplification and fragment analysis

Polymerase chain reactions (PCR) were performed in Perkin-Elmer (Norwalk, CT) thermocyclers in a total volume of 25 µl containing 50 to 100 ng each template DNA, 250 nM cy5-labelled forward primer, 250 nM unlabelled reverse primer, 0.2 mM dNTPs, 2.5 µl PCR buffer (10x), 1.5 mM MgCl₂, and 1U *Taq* DNA polymerase. After 3 min at 94°C, 45 cycles were performed with 1 min at 94°C, 1 min at either 50, 55, or 60°C (depending on the individual primer), 2 min at 72°C, and a final extension step of 10 min at 72°C. The PCR product was denatured at 94°C for 2 min and placed on a cold block until use. Each sample of 6 µl (4 µl PCR product and 2 µl internal markers) and one external standard marker of 6 µl were loaded in the preheated gel. For SSR markers, fragments were detected by an Automated Laser Fluorescence (ALF express) sequencer (Amersham Biosciences) and fragment sizes were calculated using the computer program Fragment Analyser 1.02 (Amersham Biosciences) by comparison with internal size standards. The EST-SSR, STM and SCAR markers were resolved in 2.0% agarose gels by loading 15 µl PCR products and the amplified fragments were stained with ethidium bromide and photographed.

Data analysis

Presence of *Yr* genes in the wheat genotypes was counted for each yellow rust resistance gene based on presence of the fragment sizes of both flanking markers, except for a few genes where one closely linked marker has been reported. This was explained by comparing with the reference lines with known *Yr* genes. For genetic diversity analysis, amplification profile of markers was recorded with each band representing a different allele, with a particular primer pair. Each allele was scored on the basis of

presence of a band (scored as 1) and its absence (scored as 0) for generating a binary matrix which was further used to calculate Jaccard's similarity coefficients for each pair of parents following NTSYS-PC program (Rohlf, 2000). Allelic Polymorphic Information Content (PIC) for each primer locus and genetic diversity was analysed with Power Marker version 3.25 Software (Liu and Muse, 2005).

RESULTS

Yellow rust reactions in field test

Average coefficient of infection of yellow rust for the wheat genotypes tested at three locations (Kulumsa, Arsi robe and Meraro) during 2016 and 2017 is shown in Table 2. Of the 74 test genotypes, nine cultivars showed resistance (R) with 0 to 10 ACI, 11 cultivars exhibited resistance to moderate resistance (R-MR) with ACI value of 10 to 20 and 14 genotypes showed moderately resistance (MR) with ACI value of 20 to 30, whereas 8, 9 and 23 of the genotypes exhibited moderately resistant to moderately susceptible (MR-MS), moderately susceptible (MS) and susceptible (S) with ACI values in ranges of 30 to 40, 40 to 50, and >50, respectively.

Identification of yellow rust resistance (*Yr*) genes using molecular markers

Yellow rust resistant gene, Yr1

Yr1 is a seedling resistance gene which is located on chromosome 2AL. An EST-SSR marker, BU099658 and STM marker, *Stm673acag* (1.1 cM linkage with *Yr1* gene) were used to identify this gene in 74 wheat genotypes. The expected fragment size for BU099658 marker was 206 bp (Hasancebi et al., 2014). In the present study, the marker amplified DNA fragment sizes of 157 to 206 bp. The amplified fragment at 206 bp was monomorphic and detected in all the tested materials including the reference line (*Yr1/6*Avocet S*). The polymorphic DNA fragment size, 162 bp, which was produced in the reference line was used for identification of *Yr1* gene. In this regard, the presence of *Yr1* was detected in 11 (15%) of the genotypes. The expected fragment size for the second marker, *Stm673acag* was 120 to 124 bp (Bansal et al., 2009). This marker amplified fragment sizes from 118 to 129 bp in this study. The polymorphic fragment size of 129 bp which was produced in the reference line was used to predict the *Yr1* gene. *Yr1* was identified in 11 (15%) of the tested genotypes. Only 5 (6.7%) of the genotypes exhibited similar fragment size to the reference lines for both markers.

Yellow rust resistant gene, Yr4

Yr4 which is synonymous with *Yr4a*, and *Yr4b* was

Table 2. Identified *Yr* gene in released and advanced Ethiopian bread wheat genotypes using linked molecular markers. Numeric values are the fragment sizes (bp) of PCR amplicons for the respective marker and wheat line.

Genotype	YR	<i>Yr1</i>		Genotype	YR	<i>Yr4</i>	Genotype	YR	<i>Yr6</i>	
		BU099658	Stm673acag			Barc075			WMC078	Xgwm278
<i>Yr1/6*Avocet S</i>	62	162	129	<i>Hybrid 46</i>	46.4	105	<i>Yr6/6*Avocet S</i>	82	256	289
Dereselign	-	162	129	Abola	-	105	Pavon 76	40.7	256	292
Digalu	-	162	129	HAR 1018	-	105	Abola	51.7	256	292
Dure	-	162	129	HAR 723	-	105	Alidoro	32.9	256	289
ETBW5890	-	162	129	Hawi	-	105	Bolo	76.7	256	289
ETBW5879	-	162	129	HAR 820	-	105	Bonde	55.2	256	292
Alidoro	-	156	129	Hoggana	-	105	Danda'a	20.4	256	292
Bolo	-	164	129	Jafersson	-	105	Dashen	25.7	256	289
Bonde	-	160	129	Mada Walabu	-	105	Dereselign	63.5	256	289
ETBW6094	-	159	129	PBW343	-	105	Digalu	81.5	256	289
ETBW6098	-	159	129	Simba	-	105	Batu	25.9	256	292
Tura	-	159	129	Sulla	-	105	Enkoy	13.2	256	292
Dinknesh	-	162	NA	Alidoro	-	112	ET13A2	34.7	256	292
ETBW6130	-	162	120	Batu	-	113	ETBW6093	50	256	289, 292
Galil	-	162	120	Bobicho	-	109	ETBW7698	2.7	256	292
Menze	-	162	120	Bolo	-	112	FH4-2-11	14.4	256	289, 292
Shorima	-	162	120	Bonde	-	113	Galema	26.7	256	292
Watera	-	162	124	Danda'a	-	113	Gambo	29.2	256	289
Abola	-	157	120	Dashen	-	113	Gassay	8.9	256	292
Batu	-	159	120	Dereselign	-	109	HAR 1018	9	256	289
Bobicho	-	157	120	Digalu	-	113	HAR 1331	10	256	289
Danda'a	-	156	120	Dinknesh	-	110	HAR 723	68.3	256	292
Dashen	-	157	118	Dodota	-	113	HAR 820	11.4	256	289
Dodota	-	159	120	Dure	-	110	HAR 934	12.1	256	292
Enkoy	-	157	120	Enkoy	-	113	Hidassie	32.7	256	292
ET13A2	-	157	120	ET13A2	-	113	Hoggana	50	256	289
ETBW5800	-	157	120	ETBW5800	-	113	Huluka	26.7	256	289
ETBW6093	-	159	120	ETBW5890	-	112	K6290 Bulk	55	256	292
ETBW6496	-	157	120	ETBW6093	-	112	Kakaba	55.7	256	289
ETBW6647	-	159	123	ETBW6094	-	112	Katar	60	256	289, 292
ETBW6696	-	158	124	ETBW6098	-	112	KBG-01	66.2	256	289
ETBW6861	-	157	123	ETBW6130	-	112	Kulkulu	58.4	256	289
ETBW6939	-	157	124	ETBW6496	-	112	Menze	73.4	256	289
ETBW7698	-	159	120	ETBW6647	-	112	Millennium	42.3	256	289

Table 2. Contd.

FH4-2-11	-	157	124	ETBW6696	-	113	Ogolcho	28.4	256	292
Galema	-	159	120	ETBW6861	-	112	PBW343	70	256	289
Gambo	-	157	124	ETBW6939	-	112	Shorima	28	256	289
Gassay	-	157	125	ETBW7698	-	112	Simba	30.9	256	286
HAR 1018	-	157	120	FH4-2-11	-	NA	Sirbo	30	256	292
HAR 1331	-	157	118	Galema	-	113	Sofumar	65	256	289
HAR 719	-	157	120	Galil	-	113	Sulla	71.7	256	292
HAR 723	-	159	120	Gambo	-	114	Tossa	65	256	292
HAR 727	-	157	118	Gassay	-	112	ETBW5879	30.5	256	292
HAR 820	-	158	120	HAR 1331	-	112	ETBW7255	32.1	256	292
HAR 934	-	158	120	HAR 719	-	112	Bobicho	48.4	NA	292
Hawi	-	157	120	HAR 727	-	112	Dodota	0.3	258	292
Hidassie	-	157	120	HAR 934	-	113	ETBW5890	16.3	258	292
Hoggana	-	156	120	Hidassie	-	113	ETBW6094	53.4	258	292
Huluka	-	156	120	Huluka	-	113	ETBW6098	56.7	258	292
Isreal	-	156	120	Isreal	-	113	ETBW6130	9.1	258	292
Jafersson	-	157	119	K6290 Bulk	-	113	ETBW6696	14.6	258	289, 292
K6290 Bulk	-	157	120	K6295-4A	-	113	ETBW6861	23.7	258	289, 292
K6295-4A	-	155	120	Kakaba	-	113	ETBW6939	29.3	258	289
Kakaba	-	156	120	Katar	-	113	Galil	20.9	253	292
Katar	-	156	120	KBG-01	-	113	Hawi	58.4	253	289
KBG-01	-	158	121	Kubsa	-	113	Jafersson	34.2	254	292
Kubsa	-	158	120	Kulkulu	-	113	K6295-4A	58.4	249	292
Kulkulu	-	160	127	Mandoyu	-	113	Kubsa	70	264	289
Mada Walabu	-	160	120	Menze	-	113	Mada Walabu	55	258	289
Mandoyu	-	156	120	Meraro	-	113	Mandoyu	14.5	258	292
Meraro	-	159	120	Millennium	-	110	Meraro	22.7	262	289
Millennium	-	158	120	Mitike	-	113	Mitike	50	253	289, 292
Mitike	-	156	120	Ogolcho	-	113	Sanate	4.5	265	289
Ogolcho	-	158	120	Pavon 76	-	113	Shinna	75	NA	289
Pavon 76	-	156	120	Sanate	-	110	Watera	40.8	268	292
PBW343	-	157	120	Senkegna	-	113	Dure	17.1	256	250
Sanate	-	156	120	Shinna	-	113	ETBW5800	15.9	256	250
Senkegna	-	156	120	Shorima	-	112	HAR 719	26	256	NA
Shinna	-	156	120	Sirbo	-	112	Isreal	48	256	299
Simba	-	158	121	Sofumar	-	110	Senkegna	8.4	256	295
Sirbo	-	158	120	Tossa	-	112	Tura	21.3	256	291

Table 2. Contd.

Sofumar	-	156	120	Tura	-	112	Tusie	15.4	256	299	
Sulla	-	157	120	Tusie	-	112	Dinknesh	66.9	254	296	
Tossa	-	153	120	ETBW5879	-	112	ETBW6496	13.1	258	287	
Tusie	-	152	120	Watera	-	112	ETBW6647	49.4	258	282	
ETBW7255	-	164	124	ETBW7255	-	112	HAR 727	3.5	253	296	
Frequency (%)	-	5 (7)		-	-	-	-	-	43 (57)		
Genotype	YR	Yr7		Genotype	YR	Yr8		Genotype	YR	Yr9	
		WMC078				WMC078				Xwms582	
<i>Yr7/6*Avocet S</i>	80	157		<i>Yr8/6*Avocet S</i>	83	120		<i>Yr9/6*Avocet S</i>	83	142	1100
Lee	-	159		Abola	51.7	130		Batu	48.4	142	1100
Abola	51.7	159		Alidoro	32.9	130		Bobicho	66.9	142	1100
Batu	48.4	157		Batu	48.4	130		Dashen	25.7	142	1100
Bobicho	66.9	157		Bobicho	66.9	130		ETBW6496	13.1	142	1100
Bolo	76.7	157		Bolo	76.7	130		ETBW6861	23.7	142	1100
Bonde	55.2	157		Bonde	55.2	130		FH4-2-11	14.4	142	1100
Danda'a	20.4	157		Danda'a	20.4	130		Galil	20.9	142	1100
Dashen	25.7	159		Dashen	25.7	130		HAR 1018	9	142	1100
Dereselign	63.5	157		Dereselign	63.5	130		HAR 1331	10	142	1100
Digalu	81.5	157		Digalu	81.5	130		HAR 723	68.3	142	1100
Dinknesh	0.3	159		Dinknesh	0.3	130		HAR 727	3.5	142	1100
Dodota	25.9	157		Dodota	25.9	130		HAR 820	11.4	142	1100
Dure	17.1	157		Dure	17.1	130		HAR 934	12.1	142	1100
ETBW5800	15.9	159		Enkoy	13.2	130		Hawi	58.4	142	1100
ETBW5890	16.3	159		ET13A2	34.7	130		Katar	60	142	1100
ETBW6094	53.4	157		ETBW5800	15.9	130		Mada Walabu	55	142	1100
ETBW6130	9.1	157		ETBW5890	16.3	130		Meraro	22.7	142	1100
ETBW6496	13.1	157		ETBW6093	50	130		Millennium	42.3	142	1100
ETBW6647	49.4	159		ETBW6094	53.4	130		PBW343	70	142	1100
ETBW6696	14.6	159		ETBW6098	56.7	130		Senkegna	8.4	142	1100
ETBW6861	23.7	159		ETBW6130	9.1	130		Simba	30.9	142	1100
ETBW6939	29.3	157		ETBW6496	13.1	130		Sirbo	30	142	1100
ETBW7698	2.7	159		ETBW6647	49.4	130		Tura	21.3	142	1100
FH4-2-11	14.4	159		ETBW6696	14.6	130		Tusie	15.4	142	1100
Galema	26.7	157		ETBW6861	23.7	130		Watera	40.8	142	1100
Galil	20.9	159		ETBW6939	29.3	130		ETBW7255	32.1	146	1100
HAR 1331	10	157		ETBW7698	2.7	130		Dinknesh	0.3	147	1100
HAR 719	26	159		FH4-2-11	14.4	130		ETBW6696	14.6	147	1100

Table 2. Contd.

HAR 727	3.5	157	Galema	26.7	130	Galema	26.7	147	1100
HAR 820	11.4	157	Galil	20.9	130	Dereselign	63.5	149	NA
HAR 934	12.1	159	Gambo	29.2	130	Digalu	81.5	145	NA
Hawi	58.4	157	Gassay	8.9	130	Abola	51.7	147	NA
Hidassie	32.7	159	HAR 1018	9	130	Alidoro	32.9	147	NA
Hoggana	50	157	HAR 1331	10	130	Bolo	76.7	145	NA
Huluka	26.7	157	HAR 719	26	NA	Bonde	55.2	149	NA
Isreal	48	157	HAR 723	68.3	130	Danda'a	20.4	147	NA
Jafersson	34.2	159	HAR 727	3.5	130	Dodota	25.9	145	NA
K6290 Bulk	55	157	HAR 820	11.4	130	Dure	17.1	149	NA
Kakaba	55.7	159	HAR 934	12.1	130	Enkoy	13.2	149	NA
Kubsa	70	159	Hawi	58.4	130	ET13A2	34.7	149	NA
Kulkulu	58.4	157	Hidassie	32.7	130	ETBW5800	15.9	147	NA
Mandoyu	14.5	157	Hoggana	50	130	ETBW5890	16.3	147	NA
Menze	73.4	157	Huluka	26.7	130	ETBW6093	50	147	NA
Meraro	22.7	159	Isreal	48	130	ETBW6094	53.4	147	NA
Pavon 76	40.7	159	Jafersson	34.2	130	ETBW6098	56.7	147	NA
PBW343	70	159	K6290 Bulk	55	130	ETBW6130	9.1	147	NA
Sanate	4.5	159	K6295-4A	58.4	NA	ETBW6647	49.4	147	NA
Senkegna	8.4	157	Kakaba	55.7	NA	ETBW6939	29.3	147	NA
Shinna	75	159	Katar	60	130	ETBW7698	2.7	147	NA
Shorima	28	157	KBG-01	66.2	130	Gambo	29.2	147	NA
Simba	30.9	157	Kubsa	70	130	Gassay	8.9	147	NA
Sirbo	30	159	Kulkulu	58.4	130	HAR 719	26	147	NA
Sofumar	65	159	Mada Walabu	55	130	Hidassie	32.7	NA	NA
Tossa	65	159	Mandoyu	14.5	130	Hoggana	50	147	NA
Tura	21.3	157	Menze	73.4	130	Huluka	26.7	147	NA
Tusie	15.4	157	Meraro	22.7	130	Isreal	48	149	NA
Alidoro	32.9	141	Millennium	42.3	130	Jafersson	34.2	149	NA
Enkoy	13.2	142	Mitike	50	130	K6290 Bulk	55	149	NA
ET13A2	34.7	155	Ogolcho	28.4	130	K6295-4A	58.4	145	NA
ETBW6093	50	158	Pavon 76	40.7	130	Kakaba	55.7	147	NA
ETBW6098	56.7	156	PBW343	70	130	KBG-01	66.2	147	NA
Gambo	29.2	145	Sanate	4.5	130	Kubsa	70	147	NA
Gassay	8.9	141	Senkegna	8.4	130	Kulkulu	58.4	147	NA
HAR 1018	9	143	Shinna	75	130	Mandoyu	14.5	147	NA
HAR 723	68.3	143	Shorima	28	130	Menze	73.4	145	NA

Table 2. Contd.

K6295-4A	58.4	143	Simba	30.9	130	Mitike	50	147	NA		
Katar	60	158	Sirbo	30	130	Ogolcho	28.4	147	NA		
KBG-01	66.2	158	Sofumar	65	130	Pavon 76	40.7	147	NA		
Madda Walabu	55	141	Sulla	71.7	130	Sanate	4.5	147	NA		
Millennium	42.3	143	Tossa	65	130	Shinna	75	147	NA		
Mitike	50	151	Tura	21.3	130	Shorima	28	147	NA		
Ogolcho	28.4	141	Tusie	15.4	130	Sofumar	65	147	NA		
Sulla	71.7	143	ETBW5879	30.5	130	Sulla	71.7	147	NA		
ETBW5879	30.5	158	Watera	40.8	130	Tossa	65	147	NA		
Watera	40.8	158	ETBW7255	32.1	NA	ETBW5879	30.5	147	NA		
ETBW7255	32.1	158	Frequency (%)	-	0	Frequency (%)	-	25 (33)	-		
Frequency (%)	-	55 (73%)	-	-	-	-	-	-	-		
Genotype	YR	Yr10		Genotype	YR	Yr17		Genotype	YR	Yr18	
		YR10F & YR10R				SC-385	VENTRUP			WMS295	L34DINTF9F L34PLUSR
<i>Yr10/6*Avocet S</i>	60	543	<i>Yr17/6*Avocet S</i>	77	378	258	<i>Yr18/6*Avocet S</i>	78	255	517	
Morro		543	Abola	51.7	378	258	Alidoro	32.9	255	517	
Alidoro	32.9	543	Alidoro	32.9	378	258	Bobicho	66.9	255	517	
Enkoy	13.2	543	Batu	48.4	378	258	Dodota	25.9	255	517	
ET13A2	34.7	543	Bobicho	66.9	378	258	Dure	17.1	255	517	
Galema	26.7	543	Bolo	76.7	378	258	ETBW6093	50	255	517	
Hidassie	32.7	543	Bonde	55.2	378	258	ETBW6130	9.1	255	517	
K6290 Bulk	55	543	Digalu	81.5	378	258	ETBW6496	13.1	255	517	
K6295-4A	58.4	543	Dinknesh	0.3	378	258	ETBW6647	49.4	255	517	
KBG-01	66.2	543	ETBW6093	50	378	258	ETBW6861	23.7	255	517	
Madda Walabu	55	543	ETBW6094	53.4	378	258	FH4-2-11	14.4	255	517	
Meraro	22.7	543	HAR 934	12.1	378	258	HAR 1331	10	255	517	
Mitike	50	543	Hoggana	50	378	258	HAR 719	26	255	517	
Shinna	75	543	Huluka	26.7	378	258	HAR 727	3.5	255	517	
Abola	51.7	NA	KBG-01	66.2	378	258	HAR 820	11.4	255	517	
Batu	48.4	NA	Kulkulu	58.4	378	258	HAR 934	12.1	255	517	
Bobicho	66.9	NA	Madda Walabu	55	378	258	Hidassie	32.7	255	517	
Bolo	76.7	NA	Menze	73.4	378	258	Hoggana	50	255	517	
Bonde	55.2	NA	Meraro	22.7	378	258	K6295-4A	58.4	255	517	
Danda'a	20.4	NA	Sofumar	65	378	258	Kakaba	55.7	255	517	
Dashen	25.7	NA	ETBW5879	30.5	378	258	Katar	60	255	517	
Dereselign	63.5	NA	ETBW7255	32.1	378	258	Madda Walabu	55	255	517	

Table 2. Contd.

Digalu	81.5	NA	Hawi	58.4	378	250	Sanate	4.5	255	517
Dinknesh	0.3	NA	Isreal	48	378	250	Sirbo	30	255	517
Dodota	25.9	NA	Jafersson	34.2	378	NA	Tura	21.3	255	517
Dure	17.1	NA	Danda'a	20.4	393	258	Tusie	15.4	255	517
ETBW5800	15.9	NA	ETBW6098	56.7	393	258	Digalu	81.5	255	517
ETBW5890	16.3	NA	Katar	60	393	258	Watera	40.8	255	517
ETBW6093	50	NA	Kubsa	70	393	258	Bolo	76.7	247	517
ETBW6094	53.4	NA	Millennium	42.3	393	258	Bonde	55.2	255	NA
ETBW6098	56.7	NA	Dashen	25.7	393	250	Dinknesh	0.3	255	NA
ETBW6130	9.1	NA	Dereselign	63.5	379	260	Batu	48.4	255	NA
ETBW6496	13.1	NA	Dodota	25.9	393	273	Millennium	42.3	255	NA
ETBW6647	49.4	NA	Dure	17.1	393	250	PBW343	70	255	NA
ETBW6696	14.6	NA	Enkoy	13.2	393	250	ETBW5879	30.5	255	NA
ETBW6861	23.7	NA	ET13A2	34.7	393	250	ETBW7255	32.1	255	NA
ETBW6939	29.3	NA	ETBW5800	15.9	393	250	Abola	51.7	253	NA
ETBW7698	2.7	NA	ETBW5890	16.3	393	268	Danda'a	20.4	247	NA
FH4-2-11	14.4	NA	ETBW6130	9.1	393	250	Dashen	25.7	248	NA
Galil	20.9	NA	ETBW6496	13.1	393	287	Dereselign	63.5	253	NA
Gambo	29.2	NA	ETBW6647	49.4	393	273	Enkoy	13.2	247	NA
Gassay	8.9	NA	ETBW6696	14.6	393	250	ET13A2	34.7	253	NA
HAR 1018	9	NA	ETBW6861	23.7	393	250	ETBW5800	15.9	253	NA
HAR 1331	10	NA	ETBW6939	29.3	393	250	ETBW5890	16.3	253	NA
HAR 719	26	NA	ETBW7698	2.7	393	250	ETBW6094	53.4	253	NA
HAR 723	68.3	NA	FH4-2-11	14.4	393	250	ETBW6098	56.7	247	NA
HAR 727	3.5	NA	Galema	26.7	393	268	ETBW6696	14.6	248	NA
HAR 820	11.4	NA	Galil	20.9	393	281	ETBW6939	29.3	253	NA
HAR 934	12.1	NA	Gambo	29.2	393	NA	ETBW7698	2.7	253	NA
Hawi	58.4	NA	Gassay	8.9	393	250	Galema	26.7	253	NA
Hoggana	50	NA	HAR 1018	9	393	NA	Galil	20.9	252	NA
Huluka	26.7	NA	HAR 1331	10	393	250	Gambo	29.2	248	NA
Isreal	48	NA	HAR 719	26	393	NA	Gassay	8.9	253	NA
Jafersson	34.2	NA	HAR 723	68.3	393	250	HAR 1018	9	253	NA
Kakaba	55.7	NA	HAR 727	3.5	393	250	HAR 723	68.3	253	NA
Katar	60	NA	HAR 820	11.4	393	250	Hawi	58.4	251	NA
Kubsa	70	NA	Hidassie	32.7	393	250	Huluka	26.7	252	NA
Kulkulu	58.4	NA	K6290 Bulk	55	393	NA	Isreal	48	252	NA
Mandoyu	14.5	NA	K6295-4A	58.4	393	250	Jafersson	34.2	253	NA

Table 2. Contd.

Menze	73.4	NA	Kakaba	55.7	NA	250	K6290 Bulk	55	253	NA
Millennium	42.3	NA	Mandoyu	14.5	393	250	KBG-01	66.2	253	NA
Ogolcho	28.4	NA	Mitike	50	393	NA	Kubsa	70	253	NA
Pavon 76	40.7	NA	Ogolcho	28.4	393	250	Kulkulu	58.4	253	NA
PBW343	70	NA	Pavon 76	40.7	393	252	Mandoyu	14.5	253	NA
Sanate	4.5	543	PBW343	70	393	250	Menze	73.4	253	NA
Senkegna	8.4	NA	Sanate	4.5	393	277	Meraro	22.7	248	NA
Shorima	28	NA	Senkegna	8.4	393	NA	Mitike	50	253	NA
Simba	30.9	NA	Shinna	75	393	280	Ogolcho	28.4	253	NA
Sirbo	30	NA	Shorima	28	393	284	Pavon 76	40.7	247	NA
Sofumar	65	NA	Simba	30.9	393	250	Senkegna	8.4	253	NA
Sulla	71.7	NA	Sirbo	30	393	250	Shinna	75	253	NA
Tossa	65	NA	Sulla	71.7	393	272	Shorima	28	253	NA
Tura	21.3	NA	Tossa	65	393	276	Simba	30.9	253	NA
Tusie	15.4	NA	Tura	21.3	393	250	Sofumar	65	253	NA
ETBW5879	30.5	NA	Tusie	15.4	393	250	Sulla	71.7	252	NA
Watera	40.8	NA	Watera	40.8	393	278	Tossa	65	247	NA
ETBW7255	32.1	NA	Frequency (%)	-	21 (28%)		Frequency (%)	-	28 (37%)	
Frequency (%)	-	12 (16%)	-	-	-		-	-	-	

originally derived from common wheat and it is located on chromosome 3B. An SSR marker, Barc075 was located at 2.4 ± 1.2 cM from *YrRub/Yr4* (Bansal et al., 2010). This marker amplified a single 105 bp band in the reference line (Hybrid46) and the band was absent in wheat lines without *Yr4*. After evaluation by this marker, 10 (14%) cultivars, namely, *Abola*, *HAR1018*, *HAR723*, *HAR820*, *Hawi*, *Hoggana*, *Jafersson*, *Madda Walabu*, *Simba*, and *Sulla* were genotyped to carry *Yr4* (Table 2).

Yellow rust resistance gene, *Yr6*

Li and Niu (2007) reported that the SSR markers

Wmc076 and *Wmc276* were linked to the *Yr6* gene. These two markers amplified 256 and 292 bp fragments, respectively in most lines carrying *Yr6* (Li and Niu, 2007). Marker *Wmc76* amplified 256, 267 and 271 bp bands in the reference line (*Yr6/6*Avocet S*). The expected fragment size (256 bp) that indicates for the presence of this gene was used to genotype for *Yr6* genes in the tested materials. Based on this marker, 70% of the wheat lines possessed this gene. The second marker, *Wmc276* amplified a single band of size 289 bp in the *Yr6/6*Avocet S* and 292 bp in *Pavon 76* (with known *Yr6* gene). In this regard, both fragment sizes were considered to identify this gene from the tested lines. Based on this marker, around 63 (85%) of the wheat lines possess *Yr6* (Table 2). Of the identified genotypes that

possess this gene, 42 (56%) of them showed similar results for both markers.

Yellow rust resistant gene, *Yr7*

Yr7 was first identified in wheat (*Triticum aestivum* L.) cultivar 'Lee'. It is located on the chromosome 2B (Deng et al., 2004). It is one of the *Yr* genes that was used widely by CIMMYT during the 1970s and 1980s, and has been deployed in many commercial cultivars in Ethiopia. An SSR marker, *Xgwm556*, which is linked at 5.3 cM to the target gene was used to genotype for *Yr7*. The marker allowed null allele on nine of the tested genotypes. The fragment sizes produced by other genotypes varied between 131 and 160 bp. The

marker amplified two different fragment sizes on two of the reference lines (Lee and *Yr7/6*Avocet S*). Thirty-two of the genotypes showed a fragment size of 157 bp, similar to *Yr7/6*Avocet S*. A fragment size of 159 bp was observed in 26 genotypes, which was similar to Lee. Six genotypes showed fragment sizes similar to the reference lines. Therefore, 55 (74%) of the genotypes that showed fragment sizes similar to the two reference lines were considered to possess *Yr7* gene.

Yellow rust resistant gene, *Yr8*

Yr8 is located on chromosome 2D. SSR marker Xgwm157 was linked to *Yr8* with a map distance of about 1.1 cM. The marker amplified a fragment size of 120 bp in wheat lines that possessed *Yr8* genes. In this study, this marker amplified the expected fragment size (120 bp) in the reference line, *Yr8/6*Avocet S* only, whereas none of the tested wheat genotypes amplified fragment sizes similar to the reference lines, indicating the absence of this gene in all the tested genotypes (Table 3).

Yellow rust resistant gene, *Yr9*

Yr9 was transferred to wheat through chromosomal translocation of 1B/1R and is linked to the *Lr26*, *Sr31* and *Pm8* resistance genes (Zhou et al. 2004). It is common in CIMMYT-originated bread wheat cultivars. SSR marker Xgwm582 and a resistance gene analogue (RGA) clone marker (iag95) were linked to the *Yr9* gene (Cabuk et al., 2011; Mago et al., 2002). Marker Xgwm582 amplified three bands (142, 149 and 152 bp) on the reference line. The observed fragment size, of 152 bp, was monomorphic, and it was observed in all the tested wheat genotypes. The fragment size of 142 bp which was amplified on chromosome 1B/1R where the target gene (*Yr9*) was located was used to haplotype this gene in the tested materials. Based on this, marker *Yr9* was detected in 34% of the genotypes. On the other hand, based on RGA iag95 marker (gene specific marker), 36% of the wheat genotypes possess *Yr9*. Of the identified wheat genotypes, 25 (32%) of them were amplified by both markers (Figure 1).

Yellow rust resistant gene, *Yr10*

Yr10 was originally found in wheat line PI 178383 and located on the short arm of chromosome 1B. Singh et al. (2009) designed two primer pairs (*Yr10 F/R* and *Yr10 F1/R1*) based on the *Yr10* sequence and produced markers completely linked to *Yr10*. Genotyping with this marker indicated that *Yr10* was identified in 12 (16%) of the commercially released old as well as recently released cultivars such as *Alidoro*, *Enkoy*, *ET13A2*,

Galama, *Hidassie*, *K6290 Bulk*, *K6295-4A*, *KBG-01*, *Mada Walabu*, *Meraro*, *Mitike* and *Shina*.

Yellow rust resistant gene, *Yr17*

Wheat yellow rust resistance gene, *Yr17*, in combination with *Lr37* (leaf rust resistant gene) and *Sr38* (stem rust resistant gene) are located within a segment of *Triticum ventricosum* chromosome 2NS translocated to the short arm of wheat chromosome 2A (Helguera et al., 2003). VENTRIUP-LN2 primers were 2NS specific and the 259 bp PCR amplification product was observed only in plants carrying the 2NS translocation. This marker amplified a fragment size of 259 bp on 26 (35%) entries to have *Yr17* similar to the reference line (*Yr17/6*Avocet S*). Likewise, SC-385 amplified a fragment size of 378 pb in the reference line. Based on this marker, *Yr17* was identified in 24 (32%) of the tested genotypes. These two markers exhibited a different output for nine of the genotypes. Based on these two markers, 21 (28%) of the genotypes carry this gene.

Yellow rust resistant gene, *Yr18*

The adult plant resistance gene *Yr18* was located on the same chromosome segment containing the *Lr34* gene and is tightly linked with it (Singh, 1992b). Additionally, their co-segregation with other traits such as leaf tip necrosis (*Ltn1*), powdery mildew resistance gene (*Pm38*), and tolerance to barley yellow dwarf virus (*Bdv1*) has been reported (Liang et al., 2006; Singh, 1992a; Spielmeyer et al., 2005). *Wms298* (Cabuk et al., 2011) and a gene specific marker L34DINT9F, L34PLUSR (Krattinger et al., 2009) were used to haplotype this gene. Of the 74 entries, 33 (45%) and 28 (36%) were found to carry *Yr18* according to these markers, respectively. All the wheat genotypes which were genotyped by L34DINT9F, L34PLUSR marker showed the presence of this gene by the second marker, *Wms298*.

Contribution of the identified yellow rust resistance (*Yr*) genes

The contribution of each *Yr* gene to yellow rust resistance was evaluated in differential lines that possessed the individual *Yr* gene. As shown in Table 2 and Figure 2, the differential lines carrying *Yr9* and *Yr8* had the highest ACI value (83 each) followed by *Yr6* and *Yr7* with ACI values of 82 and 80, respectively. The lowest (46.4) value was exhibited on a differential line that carried *Yr4*. The ACI values in the other differential lines that carried *Yr1*, *Yr10*, *Yr17* and *Yr18* were 62, 60, 77 and 78, respectively. This indicates that the identified resistance genes do not provide sufficient protection to wheat yellow rust if

Table 3. Total and average number of alleles, polymorphism information content (PIC), allele frequency, and rare alleles for Ethiopian commercial bread wheat and elite lines for *Yr* linked markers.

Marker	Linked to <i>Yr</i> gene	Major allele frequency	Allele No.	Gene diversity	PIC
Bu099658	<i>Yr1</i>	0.4615	6	0.6972	0.6557
Stm673acag	<i>Yr1</i>	0.1923	9	0.9191	0.9143
Barc075	<i>Yr4</i>	0.7308	3	0.4464	0.4253
WMS120	<i>Yr5/18</i>	0.2308	10	0.8830	0.8744
Xgwm76	<i>Yr6</i>	0.2949	7	0.8202	0.8000
Xgwm276	<i>Yr6</i>	0.3462	5	0.7768	0.7489
Wms526	<i>Yr7</i>	0.1026	9	0.9596	0.9581
Xgwm582	<i>Yr9</i>	0.3974	5	0.7163	0.6732
P6M12-P	<i>Yr9</i>	0.0769	15	0.9796	0.9792
iag95	<i>Yr9</i>	0.6154	1	0.4734	0.3613
Yr10F & R	<i>Yr10</i>	0.8333	1	0.2778	0.2392
Wmc273	<i>Yr15</i>	0.3846	5	0.7429	0.7053
SC-385	<i>Yr17</i>	0.7692	5	0.3744	0.3342
URIC-N2	<i>Yr17</i>	0.4359	2	0.7587	0.7373
VENTRUP	<i>Yr17</i>	0.1667	8	0.9313	0.9276
Wms295	<i>Yr18</i>	0.3077	2	0.8024	0.7766
L34DINT9F, L34PLUSR	<i>Yr18</i>	0.6538	1	0.4527	0.3502
Xgwm011	<i>Yr18</i>	0.0641	16	0.9790	0.9785
Xgwm44	<i>Yr18</i>	0.3205	11	0.7959	0.7681
Xgwc198	<i>Yr32</i>	0.8333	2	0.2817	0.2486
Xgwm18	<i>Yr24</i>	0.2436	7	0.8695	0.8576
Wmc410	<i>YrN19</i>	0.1410	11	0.9421	0.9395
Xgwm111	<i>Yr33</i>	0.0641	12	0.9711	0.9703
Xgwm508	<i>Yr35</i>	0.6154	9	0.6042	0.5910
Xgwm192	<i>Yr46</i>	0.0641	9	0.9691	0.9682
Xgwm165	<i>Yr46</i>	0.1282	6	0.9349	0.9312
Barc1182	<i>Yr1</i>	0.4359	4	0.6815	0.6242
Xgwm161	<i>Yr47</i>	0.5128	5	0.6453	0.5929
Xgwm389	<i>Yr57</i>	0.0513	17	0.9832	0.9830
Xgwc251	<i>Yr62</i>	0.1282	8	0.9415	0.9386
Mean	-	0.3530	7.24	0.7540	0.7288




Figure 1. Frequency of wheat genotypes identified with Yr genes using linked markers.




Figure 2. The distribution (%) of each Yr gene in 74 wheat genotypes and its ACI values.

they are used as single genes.

Relationship between the number of pyramided genes and yellow rust resistance

Relationship between the number of pyramided genes and yellow rust resistance is as shown in Figure 3. The number of Yr genes identified from the tested genotypes varied from 0 to 5 genes. Five varieties (*Digalu*, *HAR*

820, *HAR 934*, *Hoggana* and *Mada Walabu*) possessed the maximum number of five resistance genes. Among these varieties, the highest (82) and lowest (11.4) ACI values were recorded on *Digalu* and *HAR 820*, respectively. These five varieties exhibited ACI values of 42. On the other hand no gene was identified from bread wheat elite line, *ETBW6098*. The field study indicated that this variety exhibited an ACI value of 66.2. Twenty-six (35%) genotypes possessed three genes with an ACI value of 42 followed by 23 (31%) genotypes possessing




Figure 3. Number and ACI of *Yr* genes detected in 74 bread wheat genotypes.

two genes with an ACI value of 32.

Genetic diversity

A total of 212 alleles were detected with the 29 markers linked with *Yr* genes reported on A, B and D genomes. The number of alleles per locus ranged from 1 to 17 with an average of 7.24 alleles per locus. The highest number of alleles (17) was detected for marker Xwms389, which was linked to *Yr57* (Table 3). Genetic diversity values ranged from 0.28 to 0.97, with the highest value of 0.98 detected for marker Xwmc389. The lowest genetic diversity value was observed for marker Xwmc198. The mean genetic diversity value was observed to be 0.77. In this study, the polymorphism information (PIC) content value ranged from 0.31 to 0.98 with an average of 0.75. The highest PIC was detected by Xgwm389, which is linked to *Yr57*, whereas the lowest was detected by SC-385, which is linked to *Yr17*.

DISCUSSION

The development of molecular markers for mapping resistance genes to yellow rust and of marker-assisted selection (MAS) has been among the most active areas of research in wheat. The accuracy of MAS is affected by the distance between the target gene and the linked markers. Randhawa et al. (2014) stated that markers

used to map a gene may not be suitable for detecting the gene in diverse genetic backgrounds. In the present study, genotyping was performed only for those yellow rust resistance genes where reference lines were available as positive controls. The presence of *Yr* genes in the wheat genotypes was counted for each yellow rust resistance gene based on the presence of the fragment sizes of two flanking markers, except for a few genes where only one closely linked marker has been reported.

Molecular marker based gene identification showed the presence of *Yr1*, *Yr4*, *Yr6*, *Yr7*, *Yr9*, *Yr10*, *Yr17* and *Yr18* in various frequencies. On the other hand, *Yr8* gene was not detected in any of the tested wheat genotypes. *Yr7* is the most frequent (74%) followed by *Yr6* (56%), *Yr18* (37%) and *Yr9* (32%) whereas *Yr1* was detected at the lowest frequency (7%), followed by *Yr4* (14%). *Yr6* and *Yr7* are located on chromosomes 7BS and 2BL, respectively, and they confer all stage resistance (ASR) to yellow rust. *Yr9* was transferred to wheat through chromosomal translocation of 1B/1R (Zhou et al., 2004). These three genes were used widely by CIMMYT and have been deployed in many commercial cultivars (Van and Rajaram, 1993). Badebo et al. (1990) reported that *Yr9* gene was the most frequently (67%) identified gene from Ethiopian commercial and advanced bread wheat genotypes. But in this study, this gene was only detected in 35% of the genotypes, which may indicate the reduction of the previously widely applied *Yr9* gene in the country. However, yellow rust resistance in wheat is dominated by few *Yr* genes, such as *Yr7* and *Yr6*, which

were present in 74 and 56% of the tested varieties and lines, respectively. A diversity reduction of resistant varieties is unfavorable for breeding varieties with durable resistance. With extensive deployment of numerous commercial cultivars throughout the world, virulence for these genes was wide spread. Chen et al. (2002) reported occurrence of high virulence frequency to *Yr2*, *Yr6*, *Yr7*, and *Yr9* in most wheat producing areas of the world. Virulence frequency as high as 90% was recorded for *Yr6* and *Yr7* to *Pst* isolates collected internationally (Sharma-Poudyal et al., 2013). In Ethiopia, virulence frequencies of 92 to 100% were recorded among 107 isolates collected in 2005 from different parts of the country (Dawit et al., 2012). The average coefficients of infection (ACI) exhibited in the present study were 82 (*Yr6*), 80 (*Yr7*) and 83 (*Yr9*), indicating the ineffectiveness of these genes in the country.

The adult plant resistance gene *Yr18* is located on chromosome 7DL and is tightly linked with leaf rust resistant gene, *Lr34* (Singh, 1992a). Additionally, their co-segregation with other traits such as leaf tip necrosis (*Ltn1*), powdery mildew resistance gene (*Pm38*), and tolerance to barley yellow dwarf virus (*Bdv1*) has been reported (Liang et al., 2006; Singh, 1992a; Spielmeyer et al., 2005). This multi-pathogen resistance locus is a valuable source of resistance in wheat breeding. The use of the slow rusting gene pair *Lr34/Yr18* in combination with other slow rusting genes has been suggested to contribute to near immunity to leaf and yellow rust infections (Singh et al., 2000). Using a sequence specific marker (L34DINT9F, L34PLUSR) that indicated the presence of the cloned *Yr18* in the genomic DNA accurately, this gene was detected in 37% of the tested bread wheat genotypes. Singh (1992a) reported that despite the contribution of this gene in many countries, wheat genotypes with *Yr18* displayed inadequate resistance in some locations in Ecuador and Kenya. In the present study, an ACI value of 78 was recorded on the differential line that possessed this gene, which may indicate the ineffectiveness when used alone in Ethiopia. However, more than 67% of the bread wheat genotypes that possessed *Yr18* in different *Yr* gene combinations exhibited ACI values lower than 78. This study is the first report on the presence of the adult plant resistant gene, *Yr18*, in Ethiopian bread wheat genotypes. Those genotypes that were identified to possess *Yr18* may also possess leaf rust resistance gene (*Lr34*) and other genes linked to *Yr18* such as *Pm38*.

Yellow rust resistance gene, *Yr17* together with other genes such as *Lr37* and *Sr38* are located within a segment of *T. ventricosum* chromosome 2NS translocated to the short arm of wheat chromosome 2A (Helguera et al., 2003). The resistant gene *Yr17* was used in many breeding programs to develop resistant cultivars (Eugene et al., 2015). The yellow rust reaction on wheat seedlings with this gene is influenced by environmental conditions (for example temperature) and genetic background

(Eugene et al., 2015). Thus, it is very difficult to apply gene postulation to identify this gene in wheat genotypes. The VENTRIUP-LN2 marker, which is 2NS specific and amplifies a 259 bp PCR product only in plants carrying the 2NS translocation, was used together with the second marker SC-385 for genotyping of this gene. These markers identified 24 (32%) of the entries to have *Yr17* in the present study. Those wheat genotypes that are genotyped to have *Yr17* may also carry leaf rust (*Lr37*) and stem rust (*Sr38*) resistance genes. Earlier reports indicated that *Yr17* was postulated to be present only in three (7%) of the tested Ethiopian bread wheat genotypes (Badebo et al., 2008). On the contrary, *Yr17* was widely deployed in European wheat cultivars and a virulence frequency close to 100% has been reported in Northern European countries (Villaral et al., 2002). Yellow rust resistance gene *Yr17* was effective with regard to the prevailing races in East Africa (Badebo and Stubbs, 1995). Similarly, Dawit et al. (2012) reported the effectiveness of *Yr17* gene under Ethiopian conditions. In the same report, a virulence frequency of 14% was exhibited on this gene by isolates collected from Ethiopia, but there was no indication for the presence of these genes in the tested Ethiopian wheat cultivars. In the present study, the exhibited ACI on *Yr17* was 77, which may indicate that this gene is no more effective to the prevailing races in the country.

In addition to the aforementioned yellow rust resistant genes, this study identified the seedling resistant genes *Yr1*, *Yr4* and *Yr10*. *Yr1* is located on chromosome 2AL. Virulence to *Yr1* has been found in several countries of the world (Wan et al., 2017). But, the virulence frequency is high in East Africa where 166 Chinese wheat cultivar originated (Zhan et al., 2016). For instance, virulence frequencies of 50 and 74% were recorded in Kenya and Ethiopia, respectively (Sharma-Poudyal et al., 2013). Similarly, high yellow rust disease intensity with ACI value of 62 was exhibited in the present study. Bansal et al. (2009) reported a molecular marker, *stm673acag*, that amplified 120 and 124 bp products in most lines carrying *Yr1* gene, but in the present study the marker amplified a 129 bp fragment size, which may be due to technical reasons. Hasancebi et al. (2014) reported the second marker, *bu099658* that is linked to *Yr1* gene. This marker amplified 206 bp only in those lines that possessed *Yr1* genes, including *Yr1/6*Avocet S*. However, this marker amplified 162 bp product on the positive control (*Yr1*Avocet S*). Both markers were used to haplotype *Yr1* in the present study but showed different results. Thus, only 5 (7%) of the genotypes that exhibited a fragment size similar to both markers were considered to possess this gene. According to Bansal et al. (2009) *stm673acag* is used to haplotype both *Yr1* and *Sr48*, thus those wheat genotypes identified to possess *Yr1* may have also stem rust resistant gene *Sr48*.

Yr4 is originally derived from common wheat and is synonymous with *Yr4a* and *Yr4b*, and it is located on

chromosome 3B A microsatellite marker, Barc075, which is 2.4 ± 1.2 cM distal to *YrRub/Yr4* on chromosome 3BS, was used for genotyping. This marker was initially linked to uncharacterized resistant gene (*YrRub*) in Australian wheat cultivar Rubric. Later, the marker was tested in genotypes known to carry *Yr4* (Hybrid 46 and Avalon). Based on the amplification of the Rubric-specific PCR products at Barc075 loci, it was concluded that *YrRub* could be *Yr4* (Bansal et al., 2010). In the present study, the marker amplified a single 105 bp band in the reference line, *Yr4/6*Avocet S*. After evaluation by this marker, 11 (14%) of the bread wheat cultivars showed a similar band size to the reference line. Two of the cultivars (*HAR723* and *HAR820*), which were previously reported/postulated to have the *Yr4* gene (Badebo, 1990) exhibited a similar fragment size to the reference line. This further confirms the probability of *YrRub* to be *Yr4*. Yellow rust resistant gene *Yr4* was effective in East Africa with regard to the prevailing races (Badebo and Stubbs, 1995). A low virulence frequency of 6% was recorded in Ethiopia (Dawit et al., 2012). In the present study, *Yr4* exhibited an ACI of 50, which may indicate that this gene is no more effective if it is used alone. *Yr10* originated from bread wheat and is located on the short arm of chromosome 1B (It is one of the resistant genes that confers high resistance to *Pst* races in Pakistan (Farrakh et al., 2016) and China (Zheng et al., 2017). In Ethiopia, a yellow rust severity as high as 40% was recorded on the differential lines that possessed this gene. By using a gene specific marker, *Yr10* is confirmed to be present in 16% of the tested wheat genotypes.

Gene pyramiding, combining multiple *Yr* genes in a single genotype is an important strategy to develop durable rust resistant cultivars. The ACI value that was recorded on the genotypes that possessed the maximum number (five) of resistance genes was 42. Cultivars with four and three genes exhibited 36.19 and 42.01 ACI, respectively. This may indicate that pyramiding of those identified genes may not provide sufficient protection against the prevalent races in the country. Thus, there is an urgent need to search for more effective resistance genes to be incorporated in Ethiopian bread wheat cultivars. On the contrary, ACI values ranging from low (8.9) to high (70) were exhibited in cultivars *Gassay* and *Shina*, respectively with one gene. Those cultivars with low average coefficient of infection may indicate the presence of additional *Yr* genes that were not identified in the present study.

The number of alleles per locus ranged from 1 to 17 with an average of 7.24. Genetic diversity values ranged from 0.28 to 0.98, with the highest value of 0.98 detected for markers Xwmc389 and P6M12-P. The mean genetic diversity values were observed to be 0.75. The polymorphic information content (PIC) value ranged from 0.24 to 0.98 with an average of 0.73. The highest PIC value was detected by Xgwm389, which is linked to *Yr57*, whereas the lowest was by Yr10F & R, linked to *Yr10*.

The evidence in this study on the basis of genetic

diversity and the presence of *Yr* genes in the improved wheat genotypes will be helpful for developing appropriate breeding strategies to broadening the genetic base in future wheat breeding programs in Ethiopia.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

ACKNOWLEDGEMENTS

This research was supported by a fellowship to Woubit Dawit by the German Academic Exchange Service (DAAD). The molecular research was conducted at the Leibniz Institute of Plant Genetics and Crop Plant Research (IPK), Gatersleben, Germany. The authors thank Annette Heber for excellent technical support for genotyping.

REFERENCES

- Ahmad I, Ullah U, Inamullah, Ahmad H, Muhammad I (2015). Screening of Pakistani wheat landraces for stripe rust resistance using molecular markers. *Basic Research Journal of Soil and Environmental Science* 3(1):1-3.
- Badebo A, Stubbs RW, Ginkel MV, Gebeyehu G (1990). Identification of resistance genes to *Puccinia striiformis* in seedlings of Ethiopian and CIMMYT bread wheat varieties and lines. *Netherlands Journal of Plant Pathology* 96:199-210.
- Badebo A, Stubbs RW (1995). Valuable sources of resistance of wheat to the East African yellow rust isolates. In: *Breeding for disease resistance with emphasis on durability*. In: Proceedings of a regional workshop for Eastern, Central, and Southern Africa Danial DL (eds.), Njoro, Kenya pp. 206-214.
- Badebo A, Fehrmann H, Yahyaoui A (2008). Status of wheat stripe rust (*Puccinia striiformis*) races and their virulence in major wheat growing areas of Ethiopia. *Pest Management Journal of Ethiopia* 12:1-7.
- Bansal UK, Hayden MJ, Keller B, Wellings CR, Park RF, Bariana HS (2009). Relationship between wheat rust resistance genes *Yr1* and *Sr48* and a microsatellite marker. *Plant Pathology* 58:1039-1043.
- Bansal UK, Hayden MJ, Gill MB, Bariana SH (2010). Chromosomal location of uncharacterized stripe rust resistance gene in wheat. *Euphytica*, 171:121-127.
- Cabuk E, Aydin Y, Uncuoglu AA (2011). Assessing wheat (*Triticum aestivum*) genotypes for *Yr* resistance genes using conserved regions and simple-sequence motifs. *Genetic Molecular Research* 10:3463-3471.
- Chen XM, Moore M, Milus EA, Long DL, Line RF, Marshall D, Jackson L (2002). Wheat stripe rust epidemics and races of *Puccinia striiformis* f. sp. *tritici* in the United States in 2000. *Plant Disease* 86:39-46.
- Central Statistics Authority (CSA) (2017). Report on area and crop production forecast for Major grain crops. Addis Ababa, Ethiopia: statistical bulletin.
- Dawit W, Flath K, Weber WE, Schumann E, Röder MS, Chen X (2012). Postulation and mapping of seedling stripe rust resistance genes in Ethiopian bread wheat cultivars. *Journal of Plant Pathology* 94:403-409.
- Deng ZY, Zhang XQ, Wang XP (2004). Identification and molecular mapping of a stripe rust resistance gene from a common wheat line Qz180. *Acta Botanica* 46:36-241.
- Doyle JJ and Doyle JL (1990) Isolation of plant DNA from fresh tissue. *Focus* 12:13-15.

- Eriksen L, Afsharri F, Christiansen MJ, McIntosh RA, Jahoor A, Wellings CR (2004). *Yr32* for resistance to stripe (yellow) rust present in the wheat cultivar Carstens V. *Theoretical Applied Genetics* 108:567-575.
- Eugene AM, Kevin DL, Gina BG (2015). Characterization of stripe rust resistance in wheat lines with resistance gene *Yr17* and implication for evaluating resistance and virulence. *Phytopathology Society* 105:1123-1130.
- Farrakh S, Khalid S, Rafique A, Riaz N, Mujeeb-Kazi A (2016). Identification of stripe rust resistant genes in resistant synthetic hexaploid wheat accessions using linked markers. *Plant Genetic Resources* 14:219-225.
- Hasancebi S, Mert Z, Ertugrul F, Akan K, Aydin Y, Akfirat FS, Uncuoglu AA (2014). An EST-SSR marker, bu099658, and its potential use in breeding for yellow rust resistance in wheat. *Czech Journal of Genetics and Plant Breeding* 50:11-18
- Helguera M, Khan IA, Kolmer J, Lijavetzky D, Zhong-qi L, Dubcovsky J (2003). PCR assays for the *Lr37–Yr17–Sr38* cluster of rust resistance genes and their use to develop isogenic hard red spring wheat lines. *Crop Science* 43:1839-1847.
- Jia JQ, Li GR, Liu C, Lei MP, Yang ZJ (2011). Characterization of wheat yellow rust resistance gene *Yr17* using EST-SSR and rice syntenic region. *Cereal Research Communications* 39:88-99.
- Kaur N, Street K, Mackay M, Yahiaoui N, Keller B (2008). Molecular approaches for characterization and use of natural disease resistance in wheat. *European Journal of Plant Pathology* 21:387-397.
- Krattinger SG, Lagudah ES, Spielmeier W, Singh RP, Huerta-Espino J, McFadden H, Bossolini E, Selter LL, Keller B (2009). A putative ABC transporter confers durable resistance to multiple fungal pathogens in wheat. *Science* 323:1360-1363.
- Li Y and Niu YC (2007). Identification of molecular markers for wheat stripe rust resistance Gene *Yr6*. *Acta Agriculturae Boreali-Sinica* 22:189-192.
- Liu W, Frick M, Huel R, Nykiforuk CL, Wang X, Gaudet DA, Eudes F, Conner RL, Kuzyk A, Chen Q, Kang Z, Laroche A (2014). Stripe rust resistance gene *Yr10* encodes an evolutionary-conserved and unique CC–NBS–LRR sequence in wheat. *Molecular Plant* 7:1740-1755.
- Liu K and Muse SV (2005). Power Marker: an integrated analysis environment for genetic marker analysis. *Bioinformatics* 21:2128-2129.
- Mago R, Spielmeier W, Lawrence GJ, Lagudah ES, Ellis JG, Pryor A (2002). Identification and mapping of molecular markers linked to rust resistance genes located on chromosome 1RS of rye using wheat-rye translocation lines. *Theoretical Applied Genetics* 104:1317-1324.
- McIntosh RA, Yamazaki Y, Dubcovsky J, Rogers WK, Morris CF (2013). Catalogue of gene symbols for wheat. In: 12th International Wheat Genetics Symposium, 8-13 September 2013 Yokohama, Japan. 31p.
- McIntosh RA, Dubcovsky J, Rogers WJ, Morris C, Appels R, Xia XC (2016). Catalogue of gene symbols for wheat: 2013-14 Supplement. Annual wheat newsletter 58.
- Ma JX, Zhou RH, Dong YS, Wang LF (2001). Molecular mapping and detection of the yellow rust resistance gene *Yr26* in wheat transferred from *Triticum turgidum* using microsatellite markers. *Euphytica* 120:219-226.
- Peterson RF, Campbell AB, Hannah AE (1948). A diagrammatic scale for estimating rust intensity of leaves and stem of cereals. *Canadian Journal of Research* 26:496-500.
- Randhawa M, Bansal U, Valarik M, Klocova B, Dolezel J, Bariana H (2014). Molecular mapping of stripe rust resistance gene *Yr51* in chromosome 4AL of wheat. *Theoretical Applied Genetics* 127:317-324.
- Rohlf FJ (2000). NTSYS-PC: Numerical taxonomy and multivariate analysis system. Exeter Software. New York 44 p
- Sharma-Poudyal D, Chen XM, Wan AM, Zhan GM, Kang ZS, Cao SQ, Jin SL, Morgounov A, Akin B, Mert Z, Shah SJA, Bux H, Ashraf M, Sharma RC, Madariaga R, Puri KD, Wellings C, Xi KQ, Wanyera R, Manninger K, Ganzález MI, Koyda M, Sanin S, Patzek LJ (2013). Virulence characterization of international collections of the wheat rust pathogen, *Puccinia striiformis* f.sp. *tritici*. *Plant Disease* 97:379-386.
- Singh RP (1992a). Genetic association of leaf rust resistance gene *Lr34* with adult plant resistance to stripe rust in bread wheat. *Phytopathology* 82:835-838.
- Singh RP (1992b). Association between gene *Lr34* for leaf rust resistance and leaf tip necrosis in wheat. *Crop Science* 32:874-878.
- Singh RP, Huerta-Espino J, Rajaram S (2000). Achieving near-immunity to leaf and stripe rusts in wheat by combining slow rusting resistance genes. *Acta Phytopathologica et Entomologica Hungarica* 35:133-139.
- Singh R, Datta D, Priyamvada, Singh S, Tiwari R (2009). A diagnostic PCR based assay for stripe rust resistance gene *Yr10* in wheat. *Acta Phytopathologica et Entomologica Hungarica* 44:1-18.
- Spielmeier W, McIntosh R, Kolmer J, Lagudah E (2005). Powdery mildew resistance and *Lr34/Yr18* genes for durable resistance to leaf and stripe rust co-segregate at a locus on the short arm of chromosome 7D of wheat. *Theoretical Applied Genetics* 111:731-735.
- Van Ginkel M and Rajaram S (1993). Breeding for durable resistance to diseases: An international perspective. In: *Durability of Disease Resistance*. Jacobs T, Parlevliet JE (eds.) Wageningen, The Netherlands. The Netherlands 259 p.
- Villalaz LMMA, Lannou C, de Vallavieille-Pope C, Neema C (2002). Genetic variability in *Puccinia striiformis* f.sp. *tritici* populations sampled on a local scale during natural epidemics. *Applied and Environmental Microbiology* 68:6138-6145.
- Vanzetti LS, Campos P, Demichelis M, Lombardo LA, Aurelia PR, Vaschetto LM, Bainotti CT, Helguera M, (2011). Identification of leaf rust resistance genes in Selected Argentinean bread wheat cultivars by gene postulation and molecular markers. *Electronic Journal of Biotechnology* 14(3):9-9.
- Wan A, Kebede T, Habtemariam Z, Bekele H (2017). Virulence characterization of wheat stripe fungus *Puccinia striiformis* f.sp. *tritici* in Ethiopia and Evaluation of Ethiopian wheat germplasm for resistance to races of the pathogen from Ethiopia and the United States. *Plant Diseases* 101:73-80.
- Yaniv E, Raats D, Ronin Y, Korol AB, Grama A, Bariana H, Dubcovsky J, Schulman AH, Fahima T (2015). Evaluation of marker-assisted selection for the stripe rust resistance gene *Yr15*, introgressed from wild emmer wheat. *Molecular Breeding* 35:43.
- Zhan G, Wang F, Wan C, Han Q, Huang Li, Kang Z, Chen X (2016). Virulence and Molecular Diversity of the *Puccinia striiformis* f.sp. *tritici* Population in Xinjiang in Relation to Other Regions of Western China. *Plant Disease* 100:99-107.
- Zheng S, Li Y, Lu L, Liu Z, Zhang C, Ao D, Li L, Zhang C, Liu R, Luo C, Wu Y, Zhang L (2017). Evaluating the contribution of *Yr* genes to stripe rust resistance breeding through Marker-assisted detection in wheat. *Euphytica* 213:50-66.
- Zhou Y, Zhong HH, Zhang GS, Xia LQ, Chen XM, Gao YC, Jing ZB, Yu GJ (2004). Utilization of 1BL/1RS translocation in wheat breeding in China. *Acta Agronomica Sinica* 30:531-535.

Supplementary Table 1. Particulars of Ethiopian-grown bread wheat varieties and lines used in the study.

S/N	Variety/line	Origin/Source	Year released/ Registered	Cross/selection
1	Abolla (HAR 1522)	CIMMYT/ Ethiopia	1997	BOBWHITE/BUCKBUCK
2	Alidoro	Ethiopia/USA	2007	(HK-14-R251)
3	Batu	CIMMYT	1984	GLL/CUC//KVZ/SX (SUNBIRD)
4	Bobicho (HAR2419)	CIMMYT	2002	PEG/PF70354/KAL/BB/ALD/3/MRNG
5	Bolo (HAR 3816)	CIMMYT	2009	VEE/LIRA//BOW/3/BCN/4/KAUZ
6	Bonde	Ethiopia	landrace	landrace
7	Danda'a (DANPHE#1)	CIMMYT	2010	KIRITATI//2*PBW65/2*SERI.1B
8	Dashen	CIMMYT	1984	KAVKAZ/(SIB)BUHO//KALYANSONA/BLUEBIRD
9	Dereselign	CIMMYT	1974	CI8154/2*FR
10	Digalu	CIMMYT	2005	SHA7/KAUZ
11	Dinknesh (HAR3919)	CIMMYT	2007	CARACARA/4/CORYDON/3/PELOTAS 72380/ARTHUR- 71*2/H567.1
12	Dodota (HAR2508)	CIMMYT	2001	BJY/COC//PRL/BOW
13	Dure (HAR1008)	CIMMYT	2001	BOW"S"/YD"s"/ZZ"S" CM62045-1Y-1M-1Y-1M-6Y-1M-OY
14	Enkoy	Kenya	1974	[HEBRAND sel./(WIS 245/ SUP51)]/[FR-FN/Y)2 .A)
15	ET-13A2	Ethiopia	1981	UQ105 Sel. x ENKOY
16	ETBW5800	CIMMYT	Line	WAXWING*2/TUKURU
17	ETBW5890	ICARDA	Line	EALME4SA - 167
18	ETBW6093	CIMMYT	Line	CROC_1/AE.SQUARROSA (205)//KAUZ/3/ENEIDA/4/PSN/BOW//MILAN
19	ETBW6094	CIMMYT	Line	TC870344/GUI//TEMPORALERA M 87/AGR/3/2*WBLL1
20	ETBW6098	CIMMYT	Line	TC870344/GUI//TEMPORALERA M 87/AGR/3/2*WBLL1
21	Wane (ETBW6130)	CIMMYT	2016	SOKOLL/EXCALIBUR
22	ETBW6496	CIMMYT	Line	CROC_1/AE.SQUARROSA (205)//FCT/3/PASTOR
23	ETBW6647	CIMMYT	Line	MARCHOUC*4/SAADA/3/2*FRET2/KUKUNA//FRET2
24	ETBW6696	CIMMYT	Line	PBW343*2/KUKUNA//SRTU/3/PBW343*2/KPASTOR
25	Lemu (ETBW6861)	CIMMYT	2016	WAXWING*2/HEILO
26	ETBW6939	ICARDA	Line	UTIQUE 96/FLAG-1
27	ETBW7698	CIMMYT	Line	FRNCLN/4/WHEAR/KUKUNA/3/C80.1/3*BATAVIA//2*WBLL1
28	FH4-2-11	Ethiopia/Germany	Line	Arb//295//SM/3/149/SM//150/M
29	Galama (HAR 604)	CIMMYT	1995	4777*2//FLN/GB/3/PVN
30	Galil	Israel	2010	HORK/YAMHILL//KALYANSONA/BLUEBIRD/3/BOBWHITE
31	Gambo (QUIAU 2)	CIMMYT	2011	BABAX/LR42//BABAX*2/3/VIVITSI
32	Gassay	CIMMYT	2009	PFAU/SERI//BOBWHITE
33	HAR1018	CIMMYT	Line	2109.36?VEE/4WRM//KAL/BB/3?KAL/BB//ALD
34	HAR1331	CIMMYT	Line	L2266-1406101/BUC'S'//VPM-MOS83.11.4.8/NAC
35	HAR 719	CIMMYT	Line	L1RA 'S'
36	HAR723	CIMMYT	Line	CHIL'S'
37	HAR727	CIMMYT	Line	PEG'S'
38	HAR820	CIMMYT	Line	CHIL'S'
39	HAR934	CIMMYT	Line	TJB788.1039/PVN76
40	Hawi (HAR2501)	CIMMYT/ Ethiopia	2000	CHIL/PRL

Supplementary Table 1 Contd.

41	Hidasse (ETBW5795)	CIMMYT/ Ethiopia	2012	YANAC/3/PRL/SARA//TSI/VEE#5/4CROC-1/AE.SQUAROSA(224)//OPATTA
42	Hoggana (ETBW5780)	No inf.	2011	PYN/BAU//MILAN
43	Hulukka (ETBW5496)	ICARDA/ Ethiopia	2010	UTQUE96/3/PYN/BAU//MILAN
44	Israel	Unknown	Pre-1949	NA
45	Jefferson	USA	2012	Imported by Morell PLC
46	K 6290-Bulk	Kenya	1977	AF.M2*ROMANY
47	K 6295-4A	Kenya	1980	ROMANZ x GB -GAMENYA
48	Kakaba (Picaflor#1)	CIMMYT	2010	KIRITATI/SERI/RAYO
49	Katar (HAR 1899)	CIMMYT	1999	COOK/VEE//DOVE/SERI/3/BIY/COC
50	KBG-01	Ethiopia/Germany	2001	300-SM-501-M/HAR-1709
51	Kubsa (HAR 1685)	CIMMYT	1994	ND/VG9144//KAL/BB/3/YACO /4/VEERY #5 (ATTILA)
52	Kulkulu	ICARDA	2009	PYN/BAU//MILAN
53	Mada-Welabu (HAR1480)	ICARDA	2000	TL/3/FR/Th/Nar59*2/4/BOL'S'C M56569-/AP-1AP-5AP-2AP-OAP
54	Mandoyu	CIMMYT	2014	WORRAKATTA/PASTOR
55	Menze (HAR 3008)	CIMMYT	2007	MILAN/SHANGHAI#7
56	Meraro (11-6-24)	CIMMYT	2005	M/4/HAR1709/3/M//24/E
57	Millennium (ETBW4921)	CIMMYT	2007	ALONDRA/CEP 75630//CEP 75234/PAT 7219/3/BUCKBUCK/BIY/4/
58	Mitike (HAR1709)	Ethiopia	1994	BOW 28 / RBC
59	Ogolcho (ETBW5520)	CIMMYT	2012	WORRAKATTA/2*PASTOR
60	Pavon 76	CIMMYT	1982	VCM//CNO/7C/3/KAL/BB
61	PBW343	CIMMYT/India	1995	NORD-DESPREZ/VG-1944//KALYANSONA//BLUEBIRD/3/YACO(SIB)/4/VEERY-5
62	Sanate	CIMMYT parent	2014	14F/HAR 1685
63	Senkegna (HAR3646)	CIMMYT	2005	CHUAN-MAI-18/BAGULA
64	Shina (HAR 1868)	CIMMYT	1999	GOV9/AZ//MUS/3/R37/GHL21//KAL/BB/4/ANI
65	Shorima (ETBW5483)	ICARDA	2011	UTIQUE-96/3/PAYNE/BAGULA//MILAN
66	Simba (HAR 2536)	CIMMYT	2000	PARULA/VEERY-6//MYNA/VULTUREPRINIA
67	Sirbo (HAR 2192)	CIMMYT	2001	VS-73-600/MIRLO/3/BOBWHITE/YECORA-70//TRIFON
68	Sofumer (HAR1889)	CIMMYT	200	LIRA'S'/TAN'S'
69	Saulla	Ethiopia	2007	HAR710/RBC
70	Tossa (HAR 3123)	CIMMYT	2004	ATTILA
71	Tura (HAR 1775)	Ethiopia	1999	AROYR Sel.60/1989
72	Tusie (HAR 1407)	CIMMYT	1997	COOK/VEE//DOVE/SERI
73	Honqolo (EBW5879)	CIMMYT	2014	NJORO SD-7
74	Wetera (HAR 1920)	CIMMYT	2000	MON/VEE//SARA
75	ETBW7255	ICARDA	Line	AGUILAL/FLAG-3