

*Full Length Research Paper*

# The reactivity of CysF9[93] $\beta$ sulphhydryl group of des-HisHC3[146] $\beta$ human haemoglobin A

J. Oyebamiji Babalola<sup>1\*</sup>, N. Adesola Babarinde<sup>2</sup> and A. Idowu Adeogun<sup>3</sup>

<sup>1</sup>Department of Chemistry, University of Ibadan, Ibadan, Nigeria.

<sup>2</sup>Department of Chemical Sciences, Olabisi Onabanjo University, Ago Iwoye, Ogun, Nigeria.

<sup>3</sup>Department of Chemistry, University of Agriculture, Abeokuta, Nigeria.

Accepted 5 September, 2011

The pH dependence of the second order rate constant of the reaction of 5,5'-dithiobis(2-nitrobenzoate) (DTNB) with CysF9[93] $\beta$  sulphhydryl group of human haemoglobin A at 50 mmol dm<sup>-3</sup> is complex. The removal of the terminal HisHC3[146] $\beta$  of haemoglobin A by enzymatic cleavage with carboxypeptidase A breaks the salt bridge between HisHC3[146] $\beta$  and AspFG1[94] $\beta$  and reduces the strain on CysF9[93] $\beta$  sulphhydryl group. The pH dependence profiles of the second order rate constant for the reaction of DTNB with CysF9[93] $\beta$  sulphhydryl group of the modified haemoglobin A derivatives at 50 mmol dm<sup>-3</sup> became simple with significant reduction in the reaction rates contrary to expectations. The implication is that CysF9[93] $\beta$  becomes occluded and hence less reactive. The mean pK<sub>a</sub>s of the ionizable groups linked to the reactivity of CysF9[93] $\beta$  sulphhydryl group of des-HisHC3[146] $\beta$  human haemoglobin A were 5.52  $\pm$  0.01 and 8.29  $\pm$  0.1. These values are assigned to HisH21[143] $\beta$  and CysF9[93] $\beta$  amino acid residues, respectively.

**Key words:** Haemoglobin, CysF9[93] $\beta$ , carboxypeptidase A, 5,5'-dithiobis(2-nitrobenzoate), salt bridge.

## INTRODUCTION

Haemoglobin is the most studied protein, yet it remains evergreen (Bettati et al., 2009) because of the lack of agreement on the mechanisms underlying the fine regulations of its structural and functional properties. The reactivity of CysF9[93] $\beta$  sulphhydryl group has been of used as indicator of tertiary and quaternary structure changes in haemoglobin in the past decades (Guidotti, 1965; Antonini and Brunori, 1969; Gibson, 1973; Perutz, et al., 1974; Hensley et al., 1975; Heidner et al., 1976; Baldwin, 1980; Shanaan, 1983; Okonjo et al., 2008, 2009; 2010). The reactivity of CysF9[93] $\beta$  sulphhydryl group has been shown to be affected by the amino sequence of the haemoglobin, increased ionic strength and the presence of allosteric effectors such as inositol hexakisphosphate (Okonjo et al., 1995, 1996, 2008, 2010).

The pH dependence of the apparent second order rate constant of the reaction of 5,5'-dithiobis (2-nitrobenzoate) (DTNB) with CysF9[93] $\beta$  of human haemoglobin A at

ionic strength of 50 mmol dm<sup>-3</sup> gives a complex profile (Okonjo, et al., 1995; 1996). However, at an ionic strength of 200 mmol dm<sup>-3</sup>, the complex profile becomes simple, resembling the titration curve of a diprotic acid.

This is because the electrostatic environment of the sulphhydryl group has been screened off. A similar observation was obtained when organic phosphate, inositol hexakisphosphate, was added. This organic phosphate is known to bind to haemoglobin at the amino acid groups, ValNA1[1] $\beta$ , HisNA2[2] $\beta$ , LysEF6[82] $\beta$  and HisH21[143] $\beta$ , at the dyad axis. The change in profile observed in the presence of organic phosphate is believed to have been caused by the reduced exposure of CysF9[93] $\beta$  to sulphhydryl reagent induced by the organic phosphate. However, Bonaventura et al. (1972) had earlier reported that addition of inositol hexakisphosphate reverses the characteristics of normal haemoglobin, which were lost upon the removal of the HisHC3[146] $\beta$  residues. HisHC3[146] $\beta$  is known to contribute about 50% to the Bohr effect observed in human haemoglobin A (Perutz, et al., 1980). The cleavage of HisHC3[146] $\beta$  is therefore expected to remove the major part of the ion pair effect and to ease the strain on AspFG1[94] $\beta$  and subsequently

\*Corresponding author. E-mail: [bamijibabalola@yahoo.co.uk](mailto:bamijibabalola@yahoo.co.uk).  
Tel: +234 803 454 0881.

its neighbor CysF9[93] $\beta$  sulphhydryl group.

The influence of the ion pair formation between HisHC3[146] $\beta$  and AspFG1[94] $\beta$  on the reactivity of CysF9[93] $\beta$  sulphhydryl group should be prominent since AspFG1[94] $\beta$  is next to the CysF9[93] $\beta$  in the amino acid sequence. Therefore, this study was aimed at investigating the effect of this ion pair on the reactivity of CysF9[93] $\beta$ , since it is possible to remove the ion pair by enzymatic cleavage of the terminal HisHC3[146] $\beta$  by carboxypeptidase-A (CPA) hydrolysis of haemoglobin A. It is expected that when the strain of the salt bridge is removed from CysF9[93] $\beta$ , the amino acid will assume a relaxed condition, lower energy configuration and react faster with DTNB.

## MATERIALS AND METHODS

### Sample preparation

Blood sample of non-smoking healthy adult with homozygous haemoglobin A was obtained from the blood Bank. Haemoglobin was prepared from the blood as previously described by Okonjo et al. (1995). Enzymatic cleavage was carried out on the oxy haemoglobin A by sequential digestion according to the method of Bettati et al. (1997) using carboxypeptidase A (CPA). The samples purity was confirmed using the methods adopted by Bettati et al. (1997). Des-HisHC3[146] $\beta$  haemoglobin samples were stored in the freezer and thawed when required. The various derivatives of haemoglobin were prepared as earlier described by Okonjo et al. (1995). Prior to use for experiments, each haemoglobin sample was passed through a Dintzis ion exchange column to remove endogenous compounds and undesired ions.

### Kinetic experiment

The kinetics of the reaction of DTNB with des-HisHC3[146] $\beta$  haemoglobin samples were monitored at 412 nm on a Shimadzu 1600PC double beam UV-Visible spectrophotometer interfaced to a computer and thermostated with a Lauda 30D table Cryostat. The reaction was carried out under a second order reaction conditions as described by Okonjo et al. (1995).

## RESULTS AND DISCUSSION

Cysteine is the most reactive amino acid in haemoglobin. Human haemoglobin A has only one reactive sulphhydryl group located at the position nine on the F helix of the  $\beta$ -chain, CysF9[93] $\beta$ . The other sulphhydryl group on position 104 of the  $\alpha$ -chain is at the sub-unit interface, hence it is occluded and does not react. Cysteine only reacts in the thiolate form of DTNB. Its reactivity has been an indicator for both the tertiary and quaternary changes in haemoglobin.

The pH dependence profile of the apparent second order rate constant,  $k_{app}$ , for the reaction of des-HisHC3[146] $\beta$  oxyhaemoglobin A with DTNB is shown in Figure 1. The profile is simple and resembles the titration curve of a diprotic acid. This means that there are two


ionizable amino acids that are linked to the reactivity of CysF9[93] $\beta$ . A comparison of the reaction of des-HisHC3[146] $\beta$  HbA to that of normal haemoglobin A (inset of Figure 1) shows that the complex pH dependence profile obtained for normal haemoglobin A has been made simple for des-HisHC3[146] $\beta$ . The rate of reaction was also drastically reduced. This implies that the salt-bridge between HisHC3[146] $\beta$  and AspFG1[94] $\beta$ , which has been removed by the enzymatic hydrolysis HbA to form des-HisHC3[146] $\beta$  HbA has great implication on haemoglobin structure. This is not surprising because HisHC3[146] $\beta$  has been reported to contribute about half of the Bohr effect in haemoglobin (Perutz et al., 1980).

In order to check the effect of ligand binding to the haem on the reactivity of des-HisHC3[146] $\beta$  HbA, the pH dependence profile on  $k_{app}$  for the reaction of DTNB with des-HisHC3[146] $\beta$  carbonmonoxyhaemoglobin A is reported in Figure 2. The profile is similar to that obtained for des-HisHC3[146] $\beta$  oxyhaemoglobin A except that the rate was slightly higher. The increase in the rate compared with that of des-HisHC3[146] $\beta$  oxyhaemoglobin is not surprising because the reaction of normal carbonmonoxyhaemoglobin with DTNB is also higher than that of oxyhaemoglobin. The inset in Figure 2 shows a comparison of the reaction of normal carbonmonoxyhaemoglobin A with DTNB to that of des-HisHC3[146] $\beta$  carbonmonoxyhaemoglobin A. Apart from the change in the shape of the pH dependence profile, carbonmonoxyhaemoglobin A reacted between six and 20 times faster than des-HisHC3[146] $\beta$  carbonmono-xyhaemoglobin A.


The pH dependence profile on  $k_{app}$  for the reaction DTNB with des-HisHC3[146] $\beta$  aquomethaemoglobin A is reported in Figure 3. A similar profile to those of des-HisHC3[146] $\beta$  oxy- and carbonmonoxyhaemoglobin A was obtained. This implies that the  $Fe^{3+}$  on the haem do not have any effect on the profile. The rate was only slightly higher than that of carbonmonoxy derivative. That is, the rate of the reaction of DTNB with the derivatives of des-HisHC3[146] $\beta$  HbA was in the order; aquomet- > carbonmonoxy- > oxy-. The inset in Figure 3 shows the comparison with normal aquomethaemoglobin. The same explanations made for the oxy-derivative is also applicable. The effect of the removal HisHC3[146] $\beta$  brought about a simplification of the reaction profile thereby removing the effect of the electrostatic environment linked to the reactivity of the CysF9[93] $\beta$  sulphhydryl group. The effect also led to reduction in the accessibility of the sulphhydryl reagent to CysF9[93] $\beta$  thereby leading to the drastic reduction in the reaction rate.

### Analysis of pH dependence profiles


It has been suggested that the reactivity of sulphhydryl group depends on the conformation of the sulphhydryl group and the electrostatic effects of the charged



**Figure 1.** Dependence of  $k_{app}$  on pH for the reaction of DTNB with CysF9[93] $\beta$  sulphhydryl group of des-HisHC3[146] $\beta$  oxyhaemoglobin A at 50 mmol dm<sup>-3</sup> ionic strength. Inset is the comparison of the theoretical best fit line for normal human oxyhaemoglobin A to that of des-HisHC3[146] $\beta$  oxyhaemoglobin A. Conditions: 10  $\mu$ mol dm<sup>-3</sup> haemoglobin concentration, (haem) (5  $\mu$ mol dm<sup>-3</sup> in reactive sulphhydryl groups); 100  $\mu$ mol dm<sup>-3</sup> DTNB concentration; 5.6  $\geq$  pH  $\geq$  8.0 phosphate buffers; pH $\geq$ 8 borate buffers; ionic strength made up with NaCl; temperature of 25°C.



**Figure 2.** Dependence of  $K_{app}$  on pH for the reaction of DTNB with CysF9[93] $\beta$  sulphhydryl group of des-HisHC3[146] $\beta$  carbonmonoxyhaemoglobin A at 50 mmol dm<sup>-3</sup> ionic strength. Inset is the comparison of the theoretical best fit line for normal human carbonmonoxyhaemoglobin A to that of des-HisHC3[146] $\beta$  carbonmonoxyhaemoglobin A. Conditions: 10  $\mu$ mol dm<sup>-3</sup> haemoglobin concentration, (haem) (5  $\mu$ mol dm<sup>-3</sup> in reactive sulphhydryl groups); 100  $\mu$ mol dm<sup>-3</sup> DTNB concentration; 5.6  $\geq$  pH  $\geq$  8.0 phosphate buffers; pH $\geq$ 8 borate buffers; ionic strength made up with NaCl; temperature of 25°C.



**Figure 3.** Dependence of  $k_{app}$  on pH for the reaction of DTNB with CysF9[93]β sulphhydryl group of des-HisHC3[146]β aquomethaemoglobin A at 50 mmol dm<sup>-3</sup> ionic strength. Inset is the comparison of the theoretical best fit line for normal human aquomethaemoglobin A to that of des-HisHC3[146]β aquomethaemoglobin A. Conditions: 10 μmol dm<sup>-3</sup> haemoglobin concentration, (haem) (5 μmol dm<sup>-3</sup> in reactive sulphhydryl groups); 100 μmol dm<sup>-3</sup> DTNB concentration; 5.6 ≤ pH ≤ 8.0 phosphate buffers; pH ≥ 8 borate buffers; ionic strength made up with NaCl; temperature of 25 °C.

ionizable groups on the protein, as well as the pK of the sulphhydryl group. On the basis of these facts, Equation 1 can be tentatively considered for the analysis of the second order kinetic profile. Similarly, simple profiles like those in the Figures had been previously analysed using Equation 1.

$$k_{app} = k_1 \frac{Q_1}{Q_1 + [H^+]} + k_2 \frac{Q_2}{Q_2 + [H^+]} \quad (1)$$

In this Equation,  $k_1$  is the limiting apparent second-order rate constant at high pH for the DTNB reaction when the reactivity of the CysF9(93)β sulphhydryl group is linked to the ionization of HisH21(143)β, with ionization constant  $Q_1$ ;  $k_2$  is the limiting apparent second order rate constant at high pH when the sulphhydryl reactivity is linked to the ionization of CysF9(93)β, with ionization constant  $Q_2$ . The analyses of the simple profiles in the Figure 1 with Equation (1) gave the best-fit parameters reported in Table 1. The mean  $pQ_1$  and  $pQ_2$  values are  $5.52 \pm 0.01$  and  $8.29 \pm 0.1$  respectively, which are lower than the mean values of 6.6 and 8.8 obtained for  $pQ_1$  and  $pQ_2$  for

intact haemoglobin A (Okonjo et al., 1995). However the values can still be attributed to the same ionizable residues electrostatically linked to the reactivity of CysF9[93]β sulphhydryl group.

#### The effect of removing C-terminal of the β-chains on pH dependence profile

Salt bridges play important roles in protein structure; these include oligomerization, molecular recognition and allosteric regulation (Perutz, 1970; Fersht, 1972). Contrary to the expectation that removal of salt bridge will increase the accessibility to sulphhydryl reagent and subsequently an increased rate of reaction, it was observed that native HbA reacted faster than the des-HisHC3(146)β haemoglobin. The expectation is based on the fact that the salt bridge between HisHC3[146]β and AspFG1[94]β imposes strain on CysF9[93]β which is just next to AspFG1[94]β. The removal of the strain was expected to give CysF9[93]β freedom to react faster. However, at the removal of the ion pair between HisHC3[146]β and AspFG1[94]β, the pH dependence of

**Table 1.** Reaction of DTNB with CysF9[93] $\beta$  sulphhydryl group of des-HisHC3[146] $\beta$  human haemoglobin A.

Parameter	$k_1$ ( $\text{dm}^3\text{mol}^{-1}\text{s}^{-1}$ )	$k_2$ ( $\text{dm}^3\text{mol}^{-1}\text{s}^{-1}$ )	$pQ_1$	$pQ_2$
Oxy	5.56	6.76	5.55	8.06
Carbon monoxy	5.95	11.71	5.57	8.51
Aquomet	5.61	12.61	5.53	8.31
Mean			$5.52 \pm 0.01$	$8.29 \pm 0.1$

$K_{app}$  of the reaction of haemoglobin A with DTNB became simplified and the rate of reaction decreased in all the pH range by more than half in all the haemoglobin derivatives. The implication is that the strain on CysF9[93] $\beta$  was actually responsible for its exposure to sulphhydryl reagents and the breakage of the salt bridge occluded it. The change in conformation of CysF9[93] $\beta$  made it to lose the electrostatic environment that gave it a complex pH dependence profile. By implication, half of the Bohr effect of haemoglobin is related to complex profile and the high rate of reaction of CysF9[93] $\beta$ . This has further confirmed the fact that CysF9[93] $\beta$  is an indicator of tertiary and quaternary changes in haemoglobin

## Conclusion

The removal of HisHC3[146] $\beta$  - AspFG1[94] ion pairs of haemoglobin A is of structural consequence. CysF9[93] $\beta$  sulphhydryl group, rather than been exposed to the sulphhydryl reagent, recoiled inward thereby reducing the accessibility of the reagent. The resultant effect is obvious in the reduced rate of reaction and simple profile obtained for the pH dependence of the apparent second order rate constant.

## ACKNOWLEDGMENTS

The authors are very grateful to Prof. K.O. Okonjo for his immense assistance. The work also benefitted from the support of the Academy of Sciences for the Developing Countries, Trieste, Italy; Alexander von Humboldt-Stiftung, Bonn, Germany; the International Centre for Theoretical Physics, Trieste and the Senate Research Grant (SRG/FSC/2006/16A) of the University of Ibadan.

## REFERENCES

- Antonini E, Brunori M (1969). On the rate of conformation change associated with ligand binding in hemoglobin. *J. Biol. Chem.* 244: 3909-3912.
- Baldwin JM (1980). The structure of human carbonmonoxyhemoglobin at 2.7 Å resolution. *J. Mol. Biol.* 136: 103-128.
- Bettati S, Viappiani C, Mozzarelli A (2009). Hemoglobin, an "evergreen" red protein. *Biochim. Biophys. Acta.* 1794: 1317-1324.
- Bonaventura J, Bonaventura C, Girdiana B, Antonini E, Brunori M, Wyman J (1972). Partial Restoration of Normal Functional Properties in Carboxypeptidase A – Digested Hemoglobin. *Proc. Natl. Acad. Sci.* 8: 2174-2178.
- Gibson QH (1973). p-Mercuribenzoate as an indicator of conformation change in hemoglobin. *J. Biol. Chem.* 248: 1281-1284.
- Guidotti G (1965). The rates of reaction of the sulphhydryl group of human hemoglobin. *J. Biol. Chem.* 240: 3924-3927.
- Heidner EJ, Lardner RC, Perutz MF (1976). Structure of horse carbonmonoxyhaemoglobin. *J. Mol. Biol.* 104: 707-722.
- Hensley P, Edelstein SJ, Wharton DC, Gibson QH (1975). Conformation and spin state in methemoglobin. *J. Biol. Chem.* 250: 952-960.
- Okonjo KO, Aboluwoye CO, Babalola JO, Usanga IA (1995). Organic phosphate binding groups electrostatically linked to the reactivity of the CysF9[93] $\beta$  sulphhydryl group of hemoglobin. *J. Chem. Soc. (Faraday Trans.)* 91: 2095-2100.
- Okonjo KO, Aken'ova YA, Aboluwoye CO, Nwozo S, Akhigbe FU, Babalola JO Babarinde NA (1996). Effect of A3[6] $\beta$ Glu→Val mutation on reactivity of CysF9[93] $\beta$  sulphhydryl group of human hemoglobin S. *J. Chem. Soc. (Faraday Trans.)* 92: 1739-1746.
- Okonjo KO, Bello OS, Babalola JO (2008). Transition of hemoglobin between two tertiary conformations: the transition constant differs significantly for the major and minor hemoglobins of the Japanese quail (*Cortunix cortunix japonica*). *Biochim. Biophys. Acta* 1784: 464-471.
- Okonjo KO, Adeogun I, Babalola JO (2009). Transition of haemoglobin between two tertiary conformations: Inositol hexakisphosphate increases the transition constant and the affinity of sheep haemoglobin for 5,5'-dithiobis(2-nitrobenzoate). *Biochim. Biophys. Acta.* 1794: 398-409.
- Okonjo KO, Adeogun I, Babalola JO (2010). Tertiary conformational transition in sheep haemoglobins induced by reaction with 5,5 - dithiobis(2-nitrobenzoate) and by binding inositol hexakisphosphate. *Biophys. Chem.* 145: 65-75.
- Perutz MF (1970). Stereochemistry of cooperative effects in hemoglobin. *Nature.* 228: 726-739.
- Perutz MF, Fersht AR, Simon SR, Roberts GCK (1974). Influence of globin structure on the state of the heme. II. Allosteric transitions in methemoglobin. *Biochemistry.* 13: 2174-2286.
- Perutz MF, Kilmartin JV, Nishikura K, Fogg JH, Butler PJG, Rollema HS (1980). Identification of residues contributing to the Bohr effect of human haemoglobin. *J. Mol. Biol.* 138: 649-668.
- Shaanan B (1983). Structure of human oxyhemoglobin at 2.1 Å resolution. *J. Mol. Biol.* 171: 31-59.