

*Full Length Research Paper*

# Effects of gibberellic acid on sugarcane plants exposed to salinity under a hydroponic system

Mahmoud Shomeili<sup>1,2\*</sup>, Majid Nabipour<sup>1</sup>, Mosa Meskarbashee<sup>1</sup> and Hamid Rajabi Memari<sup>1</sup>

<sup>1</sup>Agronomy Department, College of Agriculture, Shahid Chamran University, Ahvaz, Iran.

<sup>2</sup>Agronomy Department, Sugarcane Research Center, Iran.

Accepted 1 June, 2011

A hydroponical experiment was conducted to evaluate the possible interaction of salinity (EC 0 and 9 dSm<sup>-1</sup>) and occurrence of exogenous gibberellic acid (GA<sub>3</sub>) (100 ppm) on imbalance and partitioning of nutrients in a popular sugarcane genotype (var. CP48-103) and its somaclonal tolerant variant. The results revealed that the uptake and partitioning of N, K<sup>+</sup>, Na<sup>+</sup> and Cl<sup>-</sup> content were affected as a result of salinity, so salt stress induced the accumulation of toxic elements, namely: Na<sup>+</sup> and Cl<sup>-</sup>, and particularly in CP48-103. The tolerant variant performed better by maintaining higher N and K<sup>+</sup>. Some important parameters, namely shoot/root dry matter ratio and chlorophyll content decreased, while soluble sugars and protein content increased due to salt stress. Irrespective of the genotypes, supplementing GA<sub>3</sub> (100 ppm) as foliar application play important role on imparting salt tolerance in terms of enhancing nutrient uptake, as well as the morphological and physiological aspects. The result of this study showed that inhibition of the growth of sugarcane plantlets by salt stress was removed by GA<sub>3</sub>.

**Key words:** Sugarcane, salt stress, somaclonal variant, gibberellic acid.

## INTRODUCTION

Salinization is one of the most devastating forms of land degradation threatening food production worldwide, especially in arid and semi-arid countries such as Iran. In Iran, due to a number of factors such as climate change and irrigation practices, an increasing number of landscapes became saline, causing a decreased yield of crops. However, climate change predictions indicated less rainfall and higher temperatures in the near future in most of the agricultural regions in Iran, especially in Khuzistan province, and so, experts worry that the changes will lead to even more saline lands and predict that salinity will increase from 4 to 9 dSm<sup>-1</sup> in the future. Progress in developing salt tolerant varieties has been very slow because of less knowledge on the mechanism of salt damage and complex nature of salt tolerance. Thus, understanding the adaptive mechanisms of each crop becomes necessary to improve or produce the salt resistant genotypes. Salinity may cause damage to the plants through osmotic stress, nutrient imbalance and

specific ion toxicity (Munns et al., 1986). As such, ionic imbalance is increasingly important among the others. The availability and uptake of essential elements are affected under salinity conditions, and they lead to accumulation of some elements in toxic concentrations (Munns, 1985; Marchaner, 1986). Palaniswamy and Moshi (1973) observed in sugarcane that high proportion of Na<sup>+</sup> in the exchangeable complex of soil increased the Na<sup>+</sup> content of the plant and decreased the yield. Syed and El-Swaif (1972) reported that as the concentrations of Na<sup>+</sup> and Cl<sup>-</sup> increased, N and K<sup>+</sup> decreased and P did not changed in leaf tissue of salt tolerant sugarcane genotype under saline water irrigation. According to Kwon et al. (1999), salinity tolerance was associated with Na<sup>+</sup> exclusion, while the selectivity uptake of K<sup>+</sup> over Na<sup>+</sup> was maintained at higher K<sup>+</sup>/Na<sup>+</sup> ratio in the leaf and stem of canola.

Reports are available that the external application of GA<sub>3</sub> can alleviate deleterious effects of salinity. Ashraf et al. (2002) showed that GA<sub>3</sub> application increased the nutrient uptake, dry weights, plant height, leaf area and yield of wheat under saline conditions. There is also evidence that GA<sub>3</sub> can significantly relieve NaCl-induced growth inhibition in rice (Wen et al., 2010). Starck and

\*Corresponding author. E-mail: [shomeli@gmail.com](mailto:shomeli@gmail.com). Tel: 00986 113432303. Fax:00986113432321.

Kozinska (1980) reported that the GA<sub>3</sub> caused more absorption of P and Ca<sup>2+</sup> and less absorption of Na<sup>+</sup>, while it partially adjusted the ion ratios in bean. Bejaoui (1985) concluded that the effects of exogenously applied GA<sub>3</sub> in alleviation of salt stress may be caused by activation of special enzymes which participate in RNA and protein synthesis. Aloni and Pressman (1980) discussed possible interaction between salinity and the GA<sub>3</sub> effect on petiole elongation, cellular breakdown and bolting in celery. The response to the exogenous GA<sub>3</sub> treatments depends on the elapsed time from the beginning of salinization (Amzallag and Lerner, 1992). Many studies have shown positive effects of GA<sub>3</sub> upon photosynthesis, sucrose biosynthesis and utilization of stored sucrose in stress conditions particularly when used in a range of 100 ppm in sugarcane (Alexander, 1973; Buren et al., 1979; Moore and Ginoza, 1980). However, it can increase the sugarcane stalk length from 200 to 400% in normal conditions (Bull, 1964).

The present study focuses on the following objectives:

(a) To understand the effect of salt on nutrient imbalance and variability in partitioning of nutrients in different organs, and (b) to test the efficiency of GA<sub>3</sub> on the impart of salt tolerance in terms of better nutrient uptake under salt stress situation.

## MATERIALS AND METHODS

A hydroponic culture with the use of ¼ strength Hoagland's solution (Hoagland and Arnon, 1950) was conducted by using two salinity treatments (NaCl), which include EC = 0 and 9 dSm<sup>-1</sup> mixed with and without 100 ppm GA<sub>3</sub> as foliar spray, for the 60 days old plantlet. The experiment was carried out in the Department of Agronomy and Plant Breeding, Ahvaz University, Iran, in 2010 using a popular sugarcane (*Saccharum officinarum* L.) genotype, namely CP48-103, and its salt-tolerant variant was derived from a tissue culture process in this department. The experimental trays were arranged in a factorial complete randomized design with 4 replicates.

The observations on physiological parameters and nutrient analysis were recorded when the plants were 150 days old, after which the plants were harvested, dried in an oven at 70°C, weighed and grounded to constant mass for further analysis. Dry matter partitioning to shoots and roots was estimated by calculating shoot:root ratios in dry weight basis. Nitrogen was estimated following micro Kjeldal's method described by Humphries (1956) and then the total nitrogen concentration was determined according to the method of Lindner (1944). The potassium and sodium content was estimated by the flame photometer (Jenway PFP 7, ELE Instrument Co. Ud.) method. Calcium was estimated by versene titration method as described by Jackson (1973). Chloride content was estimated by volumetric method with AgNO<sub>3</sub> solution using potassium chromate as an indicator after extraction of leaf powder with 0.05 N potassium sulphate solutions as described by Saffiqua et al. (1977). Chlorophyll content was determined using the spectrophotometric method of Hille et al. (1986). Soluble sugars were determined by the anthrone-sulphuric acid method (Irigoyen et al., 1992), while soluble proteins were determined according to the method of Bradford (1976). All measurements were conducted on three plants per each treatment. The data were subjected to analysis of variance (ANOVA), and comparisons between the mean

(LSD) tests using the statistical software SAS (SAS Institute, 1992).

## RESULTS

### Uptake and partitioning of nutrients

Results indicated that irrespective of genotypes and plant parts, reduction in N content was observed under salinity condition; however, the reduction was higher in shoot as compared to root. An overall reduction in N content in different plant parts was also associated with more accumulation of toxic ions in different plants (Table 1). Although, Nasir et al. (2010) mentioned that nitrate reductase is the enzyme that catalyses the N assimilation, which appears to suppress its activities by salinity, GA<sub>3</sub> can overcome its effect. In this study, GA<sub>3</sub> improved non-significantly the N content in two types of sugarcane plants even under NaCl-stress.

The plants supplemented with NaCl exhibited a significant increase in Na<sup>+</sup> and Cl<sup>-</sup> content of shoot and root as compared to the plants grown under non-saline condition. GA<sub>3</sub> diminished this increase clearly in the tolerant variant grown with or without NaCl. In the tolerant variant, the accumulation of these harmful ions (that is, Cl<sup>-</sup> and Na<sup>+</sup>) was lower than the parent as there was limited transfer. Tolerant variant seems to have a genetically ability to absorb Cl<sup>-</sup> in low rate and a mechanism to transfer small amounts of that toward shoot. However, GA<sub>3</sub> decreased markedly the accumulation of Cl<sup>-</sup> in both shoot and root, particularly in tolerant variant. While concentration of K<sup>+</sup> decreased markedly in shoots and roots of source variety due to salt stress, no significant change was observed in shoots of tolerant variant, except in its roots. After application of hormone, K<sup>+</sup> accumulation in the shoots of all sugarcane plants increased under non-saline and saline stress (Table 1). In contrast, Ca<sup>2+</sup> contents of shoot and root showed a little increase in both sugarcane type plants under NaCl stress, but the effect of hormone application was observed adversely even without the presence of NaCl.

The results obtained here are consistent with those of Ashraf et al. (2002) in wheat (*Triticum aestivum* L.), Kaya et al. (2006) in maize (*Zea mays* L.), Rodriguez et al. (2006) in rice (*Oriza sativa* L.) and Nasir et al. (2010) in linseed (*Linum usitatissimum* L.); however, they were in contrast with Maggio et al. (2010) findings in tomato (*Lycopersicon esculentum* L.). As a result, application of GA<sub>3</sub> to the NaCl-exposed plants was found to be more effective in the reversal of the adverse effect of NaCl stress, which caused significant responses.

### Dry matter accumulations and physiological responses to salt stress

The results for dry matter accumulation of shoots and roots

**Table 1.** Changes in sugarcane nutrient content in response to salinity stress and exogenous GA<sub>3</sub>.

Salinity (dSm <sup>-1</sup> )	GA <sub>3</sub> (ppm)		Tolerant variant					Source variety				
			N	Na <sup>+</sup>	K <sup>+</sup>	Ca <sup>2+</sup>	Cl <sup>-</sup>	N	Na <sup>+</sup>	K <sup>+</sup>	Ca <sup>2+</sup>	Cl <sup>-</sup>
0	0	Shoot	1.45	0.20	1.43	0.14	0.19	1.32	0.31	1.46	0.10	0.26
		Root	1.04	0.31	1.48	0.09	0.16	1.03	0.62	1.39	0.04	0.24
	100	Shoot	1.50	0.20	1.78	0.12	0.18	1.48	0.24	2.00	0.09	0.20
		Root	1.07	0.24	1.20	0.08	0.12	1.07	0.32	1.15	0.04	0.15
9	0	Shoot	1.35	0.27	1.44	0.15	0.39	1.12	0.49	1.17	0.11	0.43
		Root	0.96	0.55	1.19	0.11	0.48	0.94	0.95	1.14	0.06	0.87
	100	Shoot	1.39	0.24	1.69	0.15	0.26	1.20	0.39	1.35	0.11	0.35
		Root	0.99	0.41	0.95	0.11	0.29	0.98	0.95	1.31	0.05	0.49
L.S.D. (5%)			0.08	0.06	0.12	0.01	0.05	0.08	0.12	0.12	0.02	0.11


**Table 2.** Changes in sugarcane dry matter accumulation in response to salinity stress and exogenous GA<sub>3</sub>.

Salinity (dSm <sup>-1</sup> )	GA <sub>3</sub> (ppm)	Tolerant variant			Source variety		
		Dry matter (gplant <sup>-1</sup> )	Shoot mass (gplant <sup>-1</sup> )	Root mass (gplant <sup>-1</sup> )	Dry matter (gplant <sup>-1</sup> )	Shoot mass (gplant <sup>-1</sup> )	Root mass (gplant <sup>-1</sup> )
0	0	1.41	0.94	0.47	1.53	1.02	0.51
	100	1.43	0.96	0.47	1.61	1.15	0.46
9	0	1.29	0.81	0.48	0.98	0.59	0.39
	100	1.37	0.95	0.42	1.21	0.76	0.45
L.S.D. (5%)		0.05	0.06	0.04	0.21	0.18	0.04

showed that NaCl had a significant adverse effect on them (Table 2). Both the two plant types mirrored this response with approximately 36 and 8.5% reduction for parent variety and its tolerant variant, respectively. Application of GA<sub>3</sub> also had a significant positive effect on the biomass

production of both sugarcane types. From the mean data of dry weights of shoots and roots, it was evident that although dry matter accumulation declined consistently with increase in NaCl level, application of GA<sub>3</sub> was found to have alleviated the effect of salt stress on both sugarcane types.

In the absence of salinity, the parent variety was affected by GA<sub>3</sub> as compared to its tolerant variant. So, it had a superior distribution of dry matter benefit to shoot; although, this condition was completely reversed when NaCl was added. Table 2 clearly depicts that the tolerant variant had an



**Figure 1.** Effects of salinity and GA<sub>3</sub> on protein content of sugarcane plant tissue (mg.g<sup>-1</sup> fresh matter).

aimed response to GA<sub>3</sub> application under salinity.

Salinity and GA<sub>3</sub> treatments significantly affected protein content both in the source variety and in its tolerant variant (Figure 1). The significant interaction between GA<sub>3</sub> and salinity treatments revealed a positive effect of GA<sub>3</sub> on the protein content of plants, as the highest level of protein content was found in salt stressed plants, which were subjected to GA<sub>3</sub>.


The presence of NaCl caused a significant decrease in chlorophyll content as compared to the control (NaCl-free stress). The highest pigment content was found in the tolerant variant which was subjected to GA<sub>3</sub>. However, under salt stress, the lowest chlorophyll content was recorded in the source variety plants, supplied with or without GA<sub>3</sub> treatments (Figure 2). Application of GA<sub>3</sub> could not bring about a significant change in chlorophyll content especially in tolerant variant.

Soluble sugar content had an increasing trend in shoot and root tolerant variant and no significant change in source variety with the increase in salt concentration,

whereas with application GA<sub>3</sub>, the effects on this variable were better only in tolerant variant as was markedly reported in its shoots (Figure 3a and b). Overall, the effect of growth regulator application on the source variety was not significant with respect to this attribute.

## DISCUSSION

The results reported in the present study showed that GA<sub>3</sub> application helped the salinity-challenged plants to a different degree in the reversal of altered growth and physio-chemical processes in sugarcane. It has been suggested that salt stress modifies the biochemical changes taking place in the cell wall during growth, thereby preventing extension (VanVolkenburgh and Boyer, 1985; Kaya et al., 2006). GA<sub>3</sub> has been reported to promote cell division and cell elongation in sugarcane (Alexander, 1973). Thus, the observed increase in seedling growth of salt stressed sugarcane with GA<sub>3</sub>




**Figure 2.** Effects of salinity and GA<sub>3</sub> on protein contents of sugarcane plant tissue (mg mg<sup>-1</sup> fresh matter).

treatment could be due to the inherent attribute of GA<sub>3</sub> in increasing impaired cell division and cell elongation under stressed conditions. As such, salt stress was strong enough to inhibit plant growth due to reduction in gibberellin production (Kaya et al., 2006). Addition of exogenous gibberellic acid might increase seedling growth by enhancing the content of endogenous gibberellic acid as that mentioned by Rodriguez et al. (2006).


Salt stress is known to enhance the uptake and accumulation of toxic ions such as Na<sup>+</sup> and Cl<sup>-</sup> in crop species, including sugarcane (Muniaswamy, 1998; Ashraf et al., 2002; Kaya et al., 2006; Nasir et al., 2010). However, GA<sub>3</sub>-treated plants accumulated less Na<sup>+</sup> and Cl<sup>-</sup> in both shoots and roots than the untreated plants but in significant lower range in tolerant variant. These results agree with Aldesuquy (1995) who reported that GA<sub>3</sub> reduced the accumulation of toxic ions in leaves of wheat under saline conditions. Although the shoot and root N content decreased consistently in both types of sugarcane plants under salt stress, GA<sub>3</sub> application promoted the uptake of N in both shoots and roots. It is

pertinent to mention that GA<sub>3</sub> application might have mitigated the adverse effect of salt stress on nitrate reductase and enhanced its activity as mentioned by Nasir et al. (2010). In order to maintain K<sup>+</sup> transport, GA<sub>3</sub> was more effective on tolerant variant than on source variety in the presence of salinity; but without salt stress, the effects were higher in source variety.

This study showed that inhibition of the growth of sugarcane plantlets by salt stress was removed by GA<sub>3</sub>. In contrast, increasing salinization (EC) of the nutrient solution caused a reduction of the leaf area, shoot dry matter percentage and total dry matter. The observed chlorophyll depletion may be considered to be a result of the inhibition of chlorophyll biosynthesis following an increase in ethylene production brought about by the elevated NaCl content (Shah, 2007). Moreover the GA<sub>3</sub> can promote the synthesis of ethylene (Alexander, 1973), so this study did not have a positive effect on chlorophyll content. Consequently, in this study, GA<sub>3</sub> did not increase the chlorophyll in sugarcane; so, these results do not agree with those of Ashraf et al. (2002) study. During salt stress, plants adapt to osmotic stress by accumulating



(a)



(b)

**Figure 3.** Effects of salinity and GA<sub>3</sub> on (a) shoot and (b) root soluble sugar contents of sugarcane plant tissue (mg mg<sup>-1</sup> fresh matter).

some compatible solutes such as proteins, soluble sugars, proline, glycine betaine, polyols and trehalose (Ryoun et al., 1999; Sakamoto and Murata, 2002). These compounds play a predominant role in protecting plants from osmotic stress. Although many reports showed that the activities of some antioxidant enzymes were increased by salt stress, it was observed that it may be due to the application of GA<sub>3</sub> to salt-stressed plants which enhanced the activities of these enzymes and so diminished the disadvantage of salinity effects (Nasir et al., 2010). Thus, antioxidants and compatible solutes may provide a strategy to enhance salt tolerance in plants and the later studies need to determine the appearance of these agents especially in tolerant plants as quality and quantity amounts. The cumulative response of these parameters might have provided an environment to the plants to perform photosynthesis normally, which is further confirmed by the enhanced dry matter production.

On a whole-plant basis and under field conditions, there are evidences which indicate that vigorous plants may better cope with salinity (Munns et al., 2006), possibly by delaying the onset of the salinity tolerance threshold (Dalton et al., 2000). In contrast, GA<sub>3</sub>-dependent growth reductions were reported to be critical in stress adaptation and/or survival (Achard et al., 2006). Although the independent roles of GA<sub>3</sub> have been well documented (Maggio et al., 2010), it remains uncertain how this hormone coordinately regulate plant growth and stress adaptation.

## ACKNOWLEDGMENTS

The authors wish to thank M. Meskarbashee and H. Rajabi Memari who guided them in all academic necessities. They acknowledge E. Emam and S. Shahmansori, who researchers of Biotechnology Department of Karun Agro-Industry, Iran, for their assistance and comments. Also, they acknowledge S. Rahdarian for her contributions to the lab analysis.

## REFERENCES

- Achard P, Cheng H, De Grauwe L, Decat J, Schoutteten H, Moritz T, Van Der Straeten D, Peng J, Harberd NP (2006). Integration of plant responses to environmentally activated phytohormonal signals. *Sci.*, 311: 91-94.
- Aldesuquy HS (1995). Hormones induced modifications in the response of wheat flag leaf area to NaCl. *Biol. Plant*, 387: 605-611.
- Alexander AG (1973). Sugarcane physiology. Elsevier Sci. Pub. Co. Amsterdam, Netherlands. p. 752.
- Aloni B, Pressman E (1980). Interaction with salinity of GA<sub>3</sub>-induced leaf elongation, petiole pithiness and bolting in celery. *Sci. Hortic.*, 13(2): 135-142.
- Amzallag GN, Lerner HR, Poljakoff-Mayber A (1992). Interaction between mineral nutrients, Cytokinin and Gibberellic acid during growth of Sorghum at high NaCl salinity. *J. Exp. Bot.*, 43(1): 81-87.
- Ashraf M, Karim F, Rasul E (2002). Interactive effects of gibberellic acid (GA<sub>3</sub>) and salt stress on growth, ion accumulation and photosynthetic capacity of two spring wheat (*Triticum aestivum* L.) cultivars differing in salt tolerance. *Plant Growth Regul.*, 36(1): 49-59.
- BeJaoui M (1985). Interaction between NaCl and some phytohormones on soybean growth. *J. Plant Physiol.*, 120: 95-110.
- Bradford MM (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye-binding. *Anal. Biochem.*, 72: 248-254.
- Bull TA (1964). The effects of temperature, variety and age on the response of *Saccharum* spp. to applied gibberellic acid. *Aust. J. Agric. Res.*, 15: 77-84.
- Buren LL, Moore PH, Yamasaki Y (1979). Gibberellin studies with sugarcane. II. Hand sampled field trails. *Crop Sci.*, 19: 425-428.
- Dalton FN, Maggio A, Piccinni G (2000). Simulation of shoot chloride accumulation: Separation of physical and biochemical processes governing plant salt tolerance. *Plant Soil*, 219: 1-11.
- Hille J, Verheggen F, Roelvink P, Franssen H, Vankammen A, Zabel P (1986). Bleomycin resistance: A new dominant selectable marker for plant cell transformation. *Plant Mol. Biol.*, 7: 171-176.
- Hoagland DR, Arnon DI (1950). The water-culture method of growing plants without soil. *Calif. Agr. Exp. Sta. Circ.*, p. 347.
- Humphries EC (1956). Modern methods of plant analysis. Springer Verlag, Berlin, pp. 468-502.
- Irigoyen JJ, Emerich DW, Sanchez-Diaz M (1992). Water stress induced changes in concentrations of proline and total soluble sugars in nodulated alfalfa (*Medicago sativa*) plants. *Physiologia Plantarum*, 84: 55-60.
- Jackson ML (1973). Soil chemical analysis. Prentice Hall of India Pvt. Ltd. New Delhi, p. 372.
- Kaya CA, Tuna L, Alves AAC (2006). Gibberellic acid improves water deficit tolerance in maize plants. *Acta Physiol. Plant*, 28 (4): 331-337.
- Kwon TR, Harris PJC, Bourne WF (1999). Partitioning of Na<sup>+</sup>, K<sup>+</sup>, proline and total soluble sugar in relation to the salinity tolerance of *Brassica juncea* and *Brassica rapa*. *J. Kor. Soc. Hortic. Sci.*, 40(4): 425-430.
- Lindner RC (1944). Rapid analytical methods for some of the more common inorganic constituents of plant tissues. *Plant Physiol.*, 19: 76-89.
- Maggio A, Barbieri G, Raimondi G, De Pascale S (2010). Contrasting Effects of GA<sub>3</sub> Treatments on Tomato Plants Exposed to Increasing Salinity. *J. Plant Growth Regul.*, 29: 63-72.
- Marchaner H (1986). Adaptation of plants to adverse chemical soil conditions. In: Mineral Nutrition of Higher Plants. Academic Press, London, pp. 477-542.
- Moore PH, Ginoza H (1980). Gibberellin studies with sugarcane. III. Effects of rate and frequency of gibberellic acid application on stalk length and fresh weight. *Crop Sci.*, 20: 78-82.
- Muniaswamy P (1998). Growth and development of sugarcane varieties in saline environment MSc (Agri) Thesis submitted in Tamil Nadu Agricultural University Coimbatore and in collaboration with Sugarcane Breeding Institute, Coimbatore.
- Munns R (1985). Na<sup>+</sup>, K<sup>+</sup> and Cl<sup>-</sup> in xylem sap following to shoots of NaCl treated barley. *J. Exp. Bot.*, 36: 1032-1042.
- Munns R, James RA, Lauchli A (2006). Approaches to increasing the salt tolerance of wheat and other cereals. *J. Exp. Bot.*, 57: 1025-1043.
- Munns R, Termaat A (1986). Whole-plant responses to salinity. *Aust. J. Plant Physiol.*, 13: 143-160.
- Nasir KM, Siddiqui MH, Mohammad F, Naeem M, Masroor M, Khan A (2010). Calcium chloride and gibberellic acid protect linseed (*Linum usitatissimum* L.) from NaCl stress by inducing antioxidative defense system and osmoprotectant accumulation. *Acta Physiol. Plant*, 32: 121-132.
- Palaniswamy N, Moshi MJ (1973). Studies on saline irrigation waters. II. Influence on soil properties. *Madras Agric. J.*, 60: 805-809.
- Rodriguez AA, Stella AM, Storni MM, Zulpa G, Zaccaro MC (2006). Effects of cyanobacterial extracellular products and gibberellic acid on salinity tolerance in *Oryza sativa* L. *Saline Syst.*, 2: 7.
- Ryoun KT, Harris PJC, Bourne WF (1999). Partitioning of Na<sup>+</sup>, K<sup>+</sup>, proline and total soluble sugar in relation to the salinity tolerance of *Brassica juncea* and *Brassica rapa*. *J. Korean Soc. For. Hortic. Sci.*, 40(4): 425-430.
- Saffiqua PG, Keeney DR, Tanner CB (1977). Nitrogen chloride and water balance with irrigated Reser Burbank potatoes in sandy soil. *Agron. J.*, 69: 251-257.
- Sakamoto A, Murata N (2002). The role of glycine betaine in the

- protection of plants from stress: Clues from transgenic plants. *Plant Cell Environ.*, 25: 163-171.
- SAS Institute (1992). *Fundamentals of the SAS system (version 6)* SAS Ins. Inc. NY, U.S.A., p. 555.
- Shah SH (2007). Effects of salt stress on Mustard as affected by gibberellic acid application. *Gen. Appl. Plant Physiol.*, 33(1-2): 97-106.
- Starck Z, Kozinska M (1980). Effect of phytohormones on absorption and distribution of ions in the salt stressed bean plants. *Acta Soc. Bot. Poloniae*, 49: 111-125.
- Syed MM, El-Swaify SA (1972). Effect of saline water irrigation on NCo 310 and H 50-7209 cultivars of sugarcane II. Chemical composition of plants. *Trop. Agric.*, 50(1): 45-51.
- VanVolkenburgh E, Boyer JS (1985). Inhibitory effects of water deficit on maize leaf elongation. *Plant Physiol.*, 77: 190-194.
- Wen F, Zhang Z, Bai T, Xu Q, Pan Y (2010). Proteomics reveals the effects of gibberellic acid (GA<sub>3</sub>) on salt-stressed rice (*Oryza sativa* L.) shoots. *Plant Sci.*, 178: 170-175.