

Full Length Research Paper

Floristic assessment of semi evergreen forests of a peripheral site in Hadagarh Sanctuary, Odisha, India

Ashis K. Dash, V. P. Upadhyay* and H. K. Patra

Department of Botany (Environmental Sciences), Faculty of Science, Utkal University, Bhubaneswar-751015, India.

Received 12 February, 2020; Accepted 16 March, 2020

Phytosociological study was carried out in the outer periphery region of Hadagarh Wildlife Sanctuary, Odisha, India. An attempt has been made to find out the characteristics of the vegetation community of the forests of the sanctuary. Analysis of the vegetation was done following the standard ecological methods widely adopted for such studies (quadrat method). A total of 68 plant species from 30 families were documented comprising of 14 species of tree, 42 species of shrubs and saplings and 12 species of herbs and seedlings. This study provides a baseline data for all such future studies as no previous ecological assessment about the forest vegetation of the sanctuary is available. In terms of Importance Value Index (IVI), *Azadirachta indica* with IVI value of 61.45 among trees was found most dominant followed by *Shroea robusta* (46) and *Schleichera oleosa* (37). Similarly most common plant species amongst shrubs and herbs were *Ageratum conyzoides* L. (IVI-40.08) and *Mimosa pudica* L. (IVI-45.67). The sanctuary has been under various anthropogenic pressures resulting in depletion of the vegetation. Near absence of saplings and seedlings of dominant forest trees is a matter of concern in the peripheral region of the sanctuary. The study revealed that, a suitable long term management intervention to step up regeneration of population in this area will go a long way in improving overall ecological and aesthetic value of the forests of the sanctuary area.

Key words: Importance Value Index (IVI), Santhal tribe, natural vegetation, traditional knowledge, strict enforcement.

INTRODUCTION

The concept of Sustainable development has received worldwide acceptance. Sustainable planning of ecologically important regions could not be taken up due to lack of sufficient data on structure and functioning of ecosystem. Hadagarh Wildlife Sanctuary is one such area containing diverse floral and wildlife composition but does not have much information about its phytosociological composition. The floristic composition and phytosociological attributes are useful for comparing

one community with the other from season to season and from year to year under certain environmental stress factors (Singh and Weigand, 1994). Large number of animals and plants inhabit the earth in a variety of habitats and ecosystems (Wilson, 1992). Odisha is one of the richest biodiversity regions in Southeast Asia. Saxena and Brahmam (1996) reported 2,727 species of plants under 228 families and 1062 genera of which 2561 species are indigenous and 166 species are cultivated.

*Corresponding author. E-mail: vpupadhyay@gmail.com.




Figure 1. Odisha Wildlife Organisation (2020).

There is such a vast repositories of animals and plants found across the planet with a distinct biological co-relationship maintained and sustained by various adaptations vis a vis changing environment, inter dependence among the different life forms, sustained productivity and food chain/web relationships, nutrient cycling and energy flow etc. Phytosociology reveals the organization and structure of plant diversity which helps in determination of the distribution pattern of individuals among the species in a particular habitat therefore such analysis is important for understanding the functioning of any community (Warger and Morrel, 1976). Phytosociological status of Kuldiha Wildlife Sanctuary Odisha, India reveals that there is a big gap between the values of various structural parameters and tree species having very low values of IVI deserve more attention (Rout et al, 2018). Another study on Similipal Biosphere Reserve (SBR) listed 266 species of angiospermic plants represented under 72 families. Out of 266 plant species 115, 37, 103 and 11 are trees, shrubs, herbs and climbers distributed under 42, 22, 34 and 8 families, respectively (Mishra et al, 2008). Presence of high species richness and diversity, mean stand density and species rarity indicates the uniqueness and potentiality of SBR for conservation of ecosystem in its totality (Reddy et al., 2007). Forests in the peninsular India are decreasing at a fast rate, though the data on structure and functional dynamics of these forests (Parthasarathy and Sethi, 1997) are scarce. A few studies in Eastern Ghats of Tamil Nadu (Kadavul and Parthasarathy, 1999a, b; Chittiababu and Parthasarathy, 2000; Natarajan et al., 2004) have been conducted reflecting the floristic composition of forests in the area. More studies in tropical forests are (Tripathi and Singh 2009; Hegde et al., 2011, Bajpai et al., 2012; Dangwal et al., 2012; Sahu et al., 2012; Verma et al., 2013; Sangma and Lyngdoh, 2014; Ahmed and Sharma 2014; Pradhan and Rahman, 2015; Knight, 2015; Sundarapandian and Subbiah 2015;

Shahid and Joshi, 2016; Bajpai et al., 2017; Masens et al., 2017; Shiferaw et al., 2018). Sahu et al. (2010a) and Bhadra and Pattanayak (2016, 2017, 2019) have also studied tree Species Population Dynamics in the Tropical Dry Deciduous Forest of Gandhamardan Hills, Eastern Ghats, Western Odisha. Behura et al. (2016) highlighted the ecological significance of native species which are effective ecological tools towards conserving the biodiversity and enhancing the ecological services in a restored habitat. From various studies, it is seen that phytosociological characteristics of Western Ghats have been carried out over a period of time on the forest vegetation of the area whereas a few data are available about the forests of Eastern Ghats, particularly about Odisha. The phytosociological studies conducted in other parts of Odisha are by Sahu et al. (2007), Ekka and Behera, (2011), Behura et al. (2015) and Nayak et al. (2016) etc. Odisha accounts for about 45% of the forest area in the Eastern Ghats. Therefore, the present study was aimed at documenting the status of the present vegetation structure of Hadagarh Wildlife Sanctuary through a detailed phytosociological analysis of the forest ecosystem involving parameters like relative density, abundance, relative frequency, relative basal area and Important Value Index, which will be helpful to plan for a better management intervention to enhance the quality of forest ecosystem.

MATERIALS AND METHODS

Study area

The Hadagarh Wildlife Sanctuary spreads across the districts of Keonjhar and Mayurbhanj in the state of Odisha, covering an area of 191.06 Sq Kms (Longitude 86°10' to 86°22' E Latitude 21°23' to 21° 12' N) which was notified vide Government notification dated 06.12.1978 (Figure 1) (Odisha Wildlife Organization, 2020. www.wildlife.odisha.gov.in). This area, therefore came under the status of a 'protected area' under the provisions of the Government from



Figure 2. A view of study area.

the date of its notification.

It lies in the catchment of Salandi River, one of the major tributaries of river Baitarani and is close to the Hadgarh Reservoir of Salandi dam. This region is rich in mixed deciduous forests and provides ideal habitat to variety of wildlife. The Baula area of sanctuary consists of two hills on either sides of the river and spreads across Satkosia RF of Mayurbhanj and Baula RF of Keonjhar district. The valley comprises of Hadagarh reservoir and its catchments. There are 16 villages, largely dependent on the ecological resources of the sanctuary. The demography of the area reveals dominance of backward classes and tribes like Santhals and Hos. Their regular day to day needs like firewood, grazing of domestic animals, timbers, medicinal plants etc are fulfilled from the resources available within the forests. Villagers' residing near the boundary allow cattle into the sanctuary, which is a common feature in the area. The relative humidity generally remains above 60%, maximum temperature 35°C during April-May and 6 degree Celsius during November –January of the year. Average rainfall ranges from 1718 mm to 2369 mm with 86 rainy days in a year. Cyclone with wind is frequently observed during September to December of the year (Management Plan of Hadagarh Sanctuary, 2011-2012 to 2020-2021). The study area comes under the district Kendujhar, Odisha. The District of Keonjhar is very often visited by calamities like droughts, floods, cyclone and other natural calamities. (District Disaster Management and Response Plan, Keonjhar, 2016). The forest type of this area has been categorised as Dry Peninsular Sal Forests and Northern dry mixed deciduous forests (Champion and Seth, 1968) (Figure 2). The sanctuary comes under Biogeographic classification of Deccan peninsula sub-division of Chotanagpur plateau. Sandy loam is the prominent soil type and Quartzite, Quartz, Schists, Laterite are the main rocks found in the area. (Management Plan of Hadagarh Sanctuary, 2011-2012 to 2020-2021).

Vegetation study

The sampling of vegetation was done for trees and shrubs only

once for one year during summer (March –May) and for herbaceous vegetation and seedlings three times in a year i.e. for summer (March-May), winter (December-January) and rainy season (July-September). Similar observation continued for the 2nd year also. The data thus obtained were used for analysis of all structural vegetation parameters for phytosociological analysis.

Random sampling plots were identified for the study. The quadrat size was taken 10 x 10 m for trees, 5 x 5 m for shrubs and 1 x 1 m for herbs and seedlings. All the plants were classified into following three categories taking into account their girth class i.e. circumference at breast height (cbh) (i) Herbs and Seedlings (0-10 cm cbh) (ii) Shrubs and Saplings (10-30 cm cbh) and (iii) Trees - more than 30 cm cbh and a total of 20 quadrats were sampled for each strata of vegetation. The size and number of quadrats, collection of data from the quadrats were based on standard ecological methods of Misra (1968) and Kershaw (1973). Analysis was done for arriving at phyto-sociological parameters following the Philips (1959) and Curtis (1959) to compute density, basal area (BA), frequency, relative density (RD), abundance, relative frequency (RF), relative basal area (RBA) and Important Value Index (IVI). IVI was determined as the sum of the relative frequency, relative density and relative dominance for tree layer. The following formulae were used for calculation:

$$\text{Frequency} = \frac{\text{Number of quadrats in which species occur}}{\text{Total number quadrats under study}} \times 100$$

$$\text{Relative Frequency} = \frac{\text{Number of occurrences of a species}}{\text{Total number occurrences of all species}} \times 100$$

$$\text{Density} = \frac{\text{Total number of individuals}}{\text{Total number of quadrats under study}}$$

Table 1. Species, density, basal area, frequency, distribution pattern and Importance Value Index (IVI) of Tree species of a peripheral site of Hadagarh Sanctuary.

Name of Species	Local name	Density/ha	BA m ² /ha	Freq (%)	Abundance	RD	RF	RBA	IVI
<i>Azadirachta indica</i> A Juss., Mem	Neemba	105	0.348	50	2.1	30.435	26.316	4.708	61.459
<i>Shorea robusta</i> Gaertn.f. Fruct	Sal	75	0.261	40	1.875	21.739	21.053	3.537	46.329
<i>Schleichera oleosa</i> (Lour.) Oken	Kusum	5	2.407	5	1	1.449	2.632	32.595	36.675
<i>Tamarindus indica</i> L. Sp. Pi	Tentuli	20	1.932	5	4	5.797	2.632	26.161	34.590
<i>Naringi crenulata</i> (Roxb.) M. Roem.	Benta	10	0.115	10	1	2.899	5.263	23.919	32.081
<i>Lannea corromandelica</i> (Houtt)	Jia	40	0.188	20	2	11.594	10.526	2.541	24.661
<i>Strychnos nux-vomica</i> (L.)	Kuchila	20	0.072	15	1.333	5.797	7.895	0.980	14.672
<i>Eucalyptus tereticornis</i> Sm. Spec	Nilgiri	20	0.304	5	4	5.797	2.632	4.118	12.547
<i>Protium serratum</i> (Wall. ex Colebr)	Rimili	10	0.013	10	1	2.899	5.263	0.182	8.344
<i>Cassia fistula</i> L. Sp. Pi	Sunari	10	0.001	10	1	2.899	5.263	0.007	8.168
<i>Diospyros malabarica</i> (Desr.) Kostel	Kali Kendu	10	0.017	5	2	2.899	2.632	0.235	5.765
<i>Cycas circinalis</i> L. var. <i>orixensis</i>	Katha Bheru	10	0.001	5	2	2.899	2.632	0.007	5.537
<i>Acacia nilotica</i> L.	Babul	5	0.054	5	1	1.449	2.632	0.737	4.818
<i>Catunaregam spinosa</i> (Thunb.) Tirveng.	Kalei Kanta	5	0.020	5	1	1.449	2.632	0.273	4.354

Total Indi.-Total number of individuals, BA-Basal Area, Freq-Frequency, RD-Relative Density, RF-Relative Frequency, RBA-Relative Basal Area, IVI-Important Value Index.

$$\text{Relative Density (RD)} = \frac{\text{Number of Individuals of the species}}{\text{Number of individuals of all species}} \times 100$$

$$\text{Abundance} = \frac{\text{Total number of individuals of the species}}{\text{Total number of quadrats of occurrence}} \times 100$$

$$\text{Relative Abundance (RA)} = \text{Relative Frequency} + \text{Relative Density}$$

$$\text{Relative Basal Area (RBA)} = \frac{\text{Total basal area of the species}}{\text{Total basal area of all the species}} \times 100$$

Finally, the Importance Value Index was calculated on the basis of the following formula Importance Value Index (IVI) = Relative Frequency + Relative Density + Relative Dominance. Specimen were identified with the help of Working Plan (WP, 2008, 2017) of Keonjhar forest division, Flora of Orissa (Saxena and Brahmam, 1994-1996), Botany

of Bihar and Orissa (Haines, 1921-1925) and herbarium of Botany department, Utkal University, Bhubaneswar, India.

RESULTS AND DISCUSSION

Analysis of vegetation reveals that 68 plant species from 30 families were available in the area with 14 species of tree, 42 species of shrubs and saplings and 12 species of herbs and seedlings (Tables 1 to 3). The species of trees, shrubs and saplings and herbs and seedlings belonged to 13, 29 and 9 families respectively. Importance value Index (IVI) is the measurement of ecological amplitude of species (Ludwig and Reynolds, 1988) which indicates one of the ability of a species to establish over an array of habitats. It gives composite information by taking into account the relative density, relative frequency and

relative basal area. The frequency, abundance and density values are suitable for herbs and shrubs (Airi et al., 2000). IVI is an important information for all species. The vegetation characteristics of the area should be looked into through upper-story, middle-story and under-story vegetation for the purpose of a holistic assessment. The high importance value index of the species states its dominance and ecological success, its better power of regeneration and ecological amplitude in the area (Bhandari et al., 1999).

The dominant tree species having maximum RD and RF values were *Azadirachta indica* followed by *Shorea robusta* and *Lannea corromandelica* (Table 1). The IVI of *A. indica* was highest followed by *S. robusta*, *Naringi crenulata*, *Tamarindus indica* and *Schleichera oleosa*. Mishra et al., (2012) *Shorea robusta* was found as the dominant species in the SBR having IVI of 77.67 followed



Figure 3. View of a peripheral site of Hadagarh Sanctuary.

by *Terminalia alata* (16.13) and *Anogeissus latifolia* (13.43). *Wendlandia* sp. had IVI of 0.25 and was considered as the rare species of the reserve. All other tree species showed intermediate range of IVI. Among the shrubs and saplings which occupy the middle storey vegetation *Ageratum conyzoides* exhibited highest IVI followed by *Combretum roxburghii*, *Holarrhena antidysenterica* and *Combretum decandrum* (Table 2).

The middle storey vegetation comprised of a good numbers of woody climbers and liana like *Combretum roxburghii*, *Smilax macrophylla*, *Cissampelos pareira*, *Ichnocarpus frutescens*. Similarly, analysis of herbs and seedlings (understorey vegetation) (revealed *Cynodon dactylon* having the highest IVI followed by *Arundo donax*, *Mimosa pudica* and *Sida rhombifolia*. The plant species observed and encountered in the study belonged to 30 different families with most of the species belonging to Mimosaceae followed by Ebenaceae, Acanthaceae, Asteraceae, Combretaceae etc. It is important to note that majority of the families were represented by only two or less number of species. Though vegetation can be described in terms of a number of parameters including frequency, density and cover the use of any one of these quantitative parameters could lead to over-simplification or under-estimation of the status of the species (Kigomo et al., 1990; Oyun et al., 2009). Low ecological status of most of the tree species in the present investigation as evidenced by the IVIs, may be attributed to lack of dominance by any one of these species, suggesting positive interactions among the tree species (Mishra et al., 2012). In addition to the above findings, several cutting stumps were also observed indicating impact of anthropogenic activities in one of the forest vegetation in the area. Plant species like *Shorea robusta*, *Cassia fistula* have local economic importance. Similarly, plant species like *Diospyros ebenum*, *Acacia nilotica*, *Azadirachta*

indica, *Strychnos nux-vomica*, *Andrographis paniculata*, *Alangium salvifolium*, *Holarrhena antidysenterica*, *Smilax macrophylla*, *Asoaragus racemosus*., *Xantolis tomentosa* etc have been used by the local population for years for preparation of medicines. Similar use of such plant species by people have been reported by various studies in India and abroad (Brahmam and Saxena, 1990; Girach, 1992; Sahoo and Mudgal, 1995; Mohanty et al., 1996; Brahmam et al., 1996; Aminuddin and n, 1996; Mehra et al., 2014; Bajpai et al., 2016; Rout et al., 2018). These plant species need urgent attention from the conservation point of view as their numbers are decreasing owing to factors like unsystematic and uncontrolled harvesting, grazing pressure from domestic animals and many other anthropogenic pressures (WP, 2008, 2017).

The population residing inside and adjoining area of the sanctuary have been dependent on the resources of this forest ecosystem which will have long term implications. This may be fairly attributed towards their lack of awareness on the issues concerning ecosystem services. Extensive use of natural vegetation in the sanctuary in the past has led to decrease in the provisioning services and increasing human demands has been the major cause of deterioration in the condition of the natural habitats and increasing rarity of plant biodiversity (Giam et al 2010). The middle and lower storey vegetation showed presence of many saplings and seedlings of tree species like *Morinda tinctoria*, *Cassia siamea*, *Holarrhena antidysenterica*, *Azadirachta indica*, *Cesearia elliptica*, *Acacia nilotica*, *Diospyros ebenum*. These plant species need proper protection to grow as big trees. These plants are facing fierce competition in the ecosystem for space and sunlight due to threat from demands, weed species and other biotic impacts. As a result, there are a low percentage of old grown trees in the peripheral site.

Table 2. Species, density, basal area, frequency, distribution pattern and Importance Value Index (IVI) of Shrubs and saplings of a peripheral site of Hadagarh Sanctuary.

Scientific name	Local name	Frequency	R F	Density	R D	Abundance	RA	IVI
<i>Ageratum conyzoides</i> L.	<i>Poka Sungha</i>	62.5	14.164	1290	20.925	5.160	4.994	40.083
<i>Combretum roxburghii</i> Spreng.Syst	<i>Dhala Atundi</i>	55	12.465	1065	17.275	4.841	4.685	34.425
<i>Holarrhena antidysenterica</i> Wall.	<i>Kurei</i>	15	3.399	430	6.975	7.167	6.937	17.311
<i>Albizia marginata</i> (Lam)	<i>Beranga</i>	23.75	5.382	430	6.975	4.526	4.381	16.738
<i>Combretum decandrum</i> Roxb.PI.	<i>Atundi (kala)</i>	12.5	2.833	355	5.758	7.100	6.872	15.463
<i>Diospyros malabarica</i> (Desr.)	<i>Kali kendu</i>	28.75	6.516	300	4.866	2.609	2.525	13.907
<i>Strychnos nux-vomica</i> (L.)	<i>Kochila</i>	18.75	4.249	290	4.704	3.867	3.743	12.696
<i>Andrographis paniculata</i> (Burm.f.)	<i>Bhuini Nimba</i>	10	2.266	235	3.812	5.875	5.686	11.764
<i>Streblus asper</i> Lour.	<i>Sahada</i>	26.25	5.949	165	2.676	1.571	1.521	10.146
<i>Cesearia elliptica</i> Willd.	<i>Khakada</i>	18.75	4.249	185	3.001	2.467	2.387	9.638
<i>Morinda tinctoria</i> Roxb.	<i>Achu</i>	7.5	1.700	150	2.433	5.000	4.839	8.972
<i>Cissus quadrangula</i> L	<i>Hada shankha</i>	18.75	4.249	155	2.514	2.067	2.000	8.764
<i>Pergularia daemia</i> (Forssk.)	<i>Utu rali</i>	7.5	1.700	145	2.352	4.833	4.678	8.730
<i>Cissampelos pareira</i> L.var.hirsuta	<i>Akandabindu</i>	10	2.266	150	2.433	3.750	3.630	8.329
<i>Indigofera cassioides</i> Rottl.	<i>Gilri</i>	18.75	4.249	110	1.784	1.467	1.420	7.453
<i>Manilkara hexandra</i> (Roxb.)	<i>Lal banduri</i>	17.5	3.966	100	1.622	1.429	1.383	6.971
<i>Jasminum pubescens</i> Wild.Sp.	<i>Bana malli</i>	12.5	2.833	115	1.865	2.300	2.226	6.924
<i>Symphorema polyandrum</i> Wight	<i>Badichang/Mahasindu</i>	10	2.266	115	1.865	2.875	2.783	6.914
<i>Clerodendrum viscosum</i> Vent.	<i>Gobra</i>	13.75	3.116	65	1.054	1.182	1.144	5.314
<i>Woodfordia fruticosa</i> (L)	<i>Dhataki</i>	2.5	0.567	35	0.568	3.500	3.388	4.522
<i>Paederia foetida</i> L.Mant Pl.	<i>Gandhli</i>	5	1.133	40	0.649	2.000	1.936	3.718
<i>Acacia nilotica</i> L.	<i>Babul</i>	1.25	0.283	15	0.243	3.000	2.904	3.430
<i>Diospyros ebenum</i> Koenig	<i>Kendu</i>	1.25	0.283	15	0.243	3.000	2.904	3.430
<i>Asoaragus racemosus</i> Willd.	<i>Isware jata</i>	6.25	1.416	35	0.568	1.400	1.355	3.339
<i>Cassia siamea</i> Lam.	<i>Chakunda(Desi)</i>	3.75	0.850	30	0.487	2.000	1.936	3.272
<i>Diospyros montana</i> Roxb.	<i>Halada</i>	6.25	1.416	25	0.406	1.000	0.968	2.790
<i>Atalantia monophylla</i> L.	<i>Narguni</i>	3.75	0.850	20	0.324	1.333	1.291	2.465
<i>Glycosmis pentaphylla</i> (Retz.)	<i>Dubuduba/Haumircha</i>	1.25	0.283	10	0.162	2.000	1.936	2.381
<i>Ichnocarpus frutescens</i> (L.)	<i>Suama Noi</i>	2.5	0.567	10	0.162	1.000	0.968	1.697
<i>Imperata cylindrica</i> (L)	<i>Dara gadi</i>	2.5	0.567	10	0.162	1.000	0.968	1.697
<i>Alangium salvifolium</i> (L..)Wang	<i>Ankula</i>	2.5	0.567	10	0.162	1.000	0.968	1.697
<i>Thunbergia fragrans</i> Roxb.var.hispida Gamble.	<i>Natka koli</i>	2.5	0.567	10	0.162	1.000	0.968	1.697
<i>Protium serratum</i> (Wall.exColebr)	<i>Rimili</i>	1.25	0.283	5	0.081	1.000	0.968	1.332
<i>Xantolis tomentosa</i> (Roxb.)	<i>Jyestha madhu</i>	1.25	0.283	5	0.081	1.000	0.968	1.332
<i>Combretum decandrum</i> Roxb.	<i>Kala Atundi</i>	1.25	0.283	5	0.081	1.000	0.968	1.332
<i>Achyranthes aspera</i> L.	<i>Apa maranga</i>	1.25	0.283	5	0.081	1.000	0.968	1.332

Table 2. Contd.

<i>Smilax macrophylla</i> Roxb.FI	Muturi	1.25	0.283	5	0.081	1.000	0.968	1.332
<i>Mimosa himalayana</i> Gamble,Kew	kirki koli kanta	1.25	0.283	5	0.081	1.000	0.968	1.332
<i>Grewia tillifolia</i> Vahl,Symb	Kulutha	1.25	0.283	5	0.081	1.000	0.968	1.332
<i>Flacourtia jangomas</i> (Lour.Raeusch.Nom)	Mambuli Kuli	1.25	0.283	5	0.081	1.000	0.968	1.332
<i>Shorea robusta</i> Gaertn.f.Fruct	Sal	1.25	0.283	5	0.081	1.000	0.968	1.332
<i>Sida rhombifolia</i> L..Sp.PI	Bajramuli	1.25	0.283	5	0.081	1.000	0.968	1.332

Total Indi.-Total number of individuals, A-Basal Area, Freq-Frequency, RD-Relative Density,RF-Relative Frequency,RBA-Relative Basal Area, IVI-Important Value Index.

Table 3. Species, density, basal area, frequency, distribution pattern and Importance Value Index (IVI) of Herbs and seedling species of a peripheral site of Hadagarh Sanctuary.

Name of Species	Local Name	Frequency	RF	density/m2	RD	Abundance	RA	IVI
<i>Cynodon dactylon</i> L.Sp.Pi	Duba Ghasa	23.75	13.768	2.175	30.000	9.158	21.936	65.704
<i>Arundo donax</i> L.	baunsapatri grass	31.25	18.116	1.575	21.724	5.040	12.072	51.912
<i>Mimosa pudica</i> L.	Lajkuli lata	33.75	19.565	1.250	17.241	3.704	8.871	45.678
<i>Sida rhombifolia</i> L..Sp.PI	Bajramuli	16.25	9.420	0.438	6.034	2.692	6.449	21.904
<i>Blumea lacera</i> (Burm.f)	Pokasungha	13.75	7.971	0.438	6.034	3.182	7.621	21.627
<i>Pachyrrhizus</i> sp DC Rich.ex	Sankha Saga	13.75	7.971	0.388	5.345	2.818	6.750	20.066
<i>Andrographis paniculata</i> (Burm.f)Wall.ex	Bhuin neemba	15	8.696	0.363	5.000	2.417	5.789	19.484
<i>Lygodium flexuosum</i> (L.) Sw.J	Mahajala	8.75	5.072	0.225	3.103	2.571	6.159	14.335
<i>Ageratum conyzoides</i> L.Sp.Pi	Bok sunga/ Deksingi	3.75	2.174	0.100	1.379	2.667	6.387	9.941
<i>Solanum insanum</i> L.Mant.PI	Kanta Baigana	5	2.899	0.113	1.552	2.250	5.389	9.840
<i>Sida acuta</i> Burm.f	Sunakhadika	5	2.899	0.113	1.552	2.250	5.389	9.840
<i>Cassia occidentalis</i> L.Sp.Pi	Kala Chakunda	2.5	1.449	0.075	1.034	3.000	7.186	9.670

Total Indi.-Total number of individuals, BA-Basal Area, Freq-Frequency, RD-Relative Density,RF-Relative Frequency,RBA-Relative Basal Area, IVI-Important Value Index.

These species have shown good regeneration pattern in some of the areas of the study. Therefore, it is important that anthropogenic activities in this area like overgrazing, encroachments, illegal felling etc need to be curtailed suitable intervention (WP, 2008, 2017).

Despite having the status of a protected area of the country; the vegetation needs more attention

from the habitat conservation point of view. The area needs management intervention for speedy regeneration of plant species. The phytosociological analysis reveals good generation potential of the habitat and the tree species having low IVIs, tree saplings and seedlings deserve due attention to ensure strict enforcement of the rules and better monitoring for enhancement

of ecosystem stability of this protected area. For addressing forest management issues of the sanctuary, collection and analysis of long term ecological data by scientific baseline studies covering all structural parameters will be helpful to know the present state of ecological health of the ecosystem. Various forms of anthropogenic pressures have affected the habitat which include

logging, illegal hunting, and other development challenges like mining in peripheral areas. The conservation efforts as per working (management) plan prescriptions have not so far yielded desired result. Therefore, for a sustainable ecosystem and community management and for reducing biotic pressure, the management methodology also needs to be modified for developing a Long Term Research Network to ensure improvement in structure and function of ecosystem.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

ACKNOWLEDGEMENTS

The authors express their gratitude to the State Forest Department of Odisha and all those involved in the study.

REFERENCES

- Ahmed J, Sharma S (2014). Spatial pattern, diversity and phytosociological analysis of woody plant species in Ponda watershed, Rajouri, J&K, India. *International Journal of Current Research* 6(6):7022-7027.
- Airi S, Rawal, RS, Dhar U, Purohit AN (2000). Assessment of Availability and Habitat Preference of Jatamasi a Critically Endangered Medicinal Plant of West Himalayas. *Current Science* 79:1467-1470.
- Aminuddin GRD (1996). Native phytotherapy among the Paudi bhuinya of Bonai hills. *Ethnobotany* 8:66-70.
- Bajpai O, Kumar A, Mishra AK, Sahu N, Pandey J, Behera SK, Chaudhary LB (2012). Recongregation of tree species of Katarniaghat Wildlife Sanctuary, Uttar Pradesh, India. *Journal of Biodiversity and Environmental Sciences* 2(12):24-40.
- Bajpai O, Pandey J, Chaudhary LB (2016). Ethnomedicinal uses of tree species by Tharu tribes in the Himalayan Terai region of India. *Research Journal of Medicinal Plant* 10(1):19-41.
- Bajpai O, Suman S, Upadhyay N (2017). Ecological exploration of Kuwana forest A tropical moist deciduous forest of eastern Terai, India. *Annals of Plant Sciences* 6(12):1811-1816.
- Behura S, Kar M, Upadhyay VP (2015). Phytosociological Study of herbs species at two reclaimed sites of Sukinda Chromite mining region of Odisha, India. *Geo-Eco-Tropical* 39(2):329-342.
- Behura S, Kar M, Upadhyay VP (2016). Ecological analysis of tree species at two reclaimed sites of Sukinda chromite mining region of Odisha, India. *International Journal of Advanced Research in Biological Sciences* 3:51-59.
- Bhadra AK, Pattanayak SK (2016). Abundance or Dominance: Which is more justified to calculate Importance Value Index (IVI) of plant species? *Asian Journal of Science and Technology* 7(9):3577-3601.
- Bhadra AK, Pattanayak SK (2017). Dominance is more justified than Abundance to calculate Importance Value Index (IVI) of plant species. *Asian Journal of Science and Technology* 8(2):4304-4326.
- Bhadra AK, Pattanayak SK (2019). Tree Species Population dynamics in the Tropical Dry Deciduous Forest of Gandhamardan Hills, Eastern Ghats, Western Odisha. *Indian Forester* 145(2):101-113.
- Bhandari BS, Nautiyal DC, Gaur RD (1999). Structural attributes and Some noteworthy folk-medicinal uses. *Ethnobotany* 2:71-79.
- Brahmam M, Dhal NK, Saxena HO (1996). Ethnobotanical studies among the Tania of Malaygiri hills in Dhenkanal district, Orissa, India. In: Jain, S.K. (Ed.). *Ethnobiology in Human Welfare*, Deep Publications, New Delhi pp. 393-396.
- Champion HG, Seth SK (1968). A revised survey of the forest types of India. *Manager Publications, New Delhi*, pp. 16-17.
- Chittiababu CV, Parthasarathy N (2000). Attenuated tree species diversity in human-impacted tropical evergreen forest sites at Koll hills, Eastern Ghats, India. *Biodiversity and Conservation* 9:1493-1519.
- Curtis JT (1959). *The vegetation of Wisconsin. An ordination of plant Communities*. University of Wisconsin Press, Madison Wisconsin.
- Dangwal LR, Singh T, Singh A, Sharma A (2012). Species composition of woody plants in forest of block Nowshera, district Rajouri (J&K), India. *Journal of Current Research* 4(5):5-10.
- District Disaster Management and Response Plan, Keonjhar (2016). Available at <https://kendujhar.nic.in/document/district-management-and-response-plan-2016-17/>.
- Ekka NJ, Behera N (2011). Species composition and diversity of vegetation developing on an age series of coal mine spoil in an opencast coal field in Odisha, India. *Tropical Ecology* 52:337-343.
- Giam X, Bradshaw CJA, Tan HTW, Sodhi NS (2010). Future habitat loss and the conservation of plant biodiversity. *Biological Conservation* 143:1594-1602.
- Girach RD (1992). Medicinal plants used by Kondh tribe of district Phulbani (Orissa) in Eastern India. *Ethnobotany* 4:53-66.
- Haines HH (1925). *The Botany of Bihar and Orissa*. Vol. I-III. London, Botanical Survey of India, Calcutta (Repr. Edn. 1961).
- Hegde GT, Murthy IK, Bhat PR, Swarnim S, Alipuria AK, Ravindranath NH (2011). Vegetation status in degraded forest, community and private lands of Himanchal Pradesh. *Indian Forester* 137(5):544-553.
- Kadavul K, Parthasarathy N (1999a). Plant biodiversity and conservation of tropical semi-evergreen forest in the Sherwarayan hills of Eastern Ghats, India. *Biodiversity and Conservation* 8:421-439.
- Kadavul K, Parthasarathy N (1999b). Structure and composition of woody species in tropical semi-evergreen forest of Kalrayan hills, Eastern Ghats, India. *Tropical Ecology* 40:247-260.
- Kershaw KR (1973). *Quantitative and Dynamic Plant Ecology*. Edward Arnold Ltd., ISBN0713120991, London.
- Kigomo BN, Savill PS, Woodell SR (1990). Forest Composition and Its Regeneration Dynamics: A Case Study of Semi-Deciduous Tropical Forest in Kenya. *African Journal of Ecology* 28(3):174-187.
- Knight DH (2015). From Phytosociology and Plant Evolution to Ecosystem Analysis. *Ecological Society of America Bulletin* 96(2):209-210.
- Ludwig JA, Reynolds JF (1988). *Statistical Ecology: A Primer on Methods and Computing*. Wiley-Interscience, ISBN 10-0471832359, New York.
- Lyngdoh N, Sangma DM (2014). Diversity Assessment of Nokrek Biosphere Reserve, Meghalaya.
- Management Plan of Hadagarh Sanctuary (2011). Keonjhar Wildlife Division, (2011-12 to 2020-21) Keonjhar, Orissa pp. 1-64.
- Masens da-Musa YB, Ngbolua KTN, Masens M, Tambu TM, Gédéon NB (2017). Phytocological study of Nzundu massif forest of Imbongo city, Kwilu Province, Democratic Republic of the Congo. *Tropical Plant Research* 4(3):363-375.
- Mehra A, Bajpai O, Joshi H (2014). Diversity, utilization and sacred values of Ethno-medicinal plants of Kumaun Himalaya. *Tropical Plant Research* 1(3):80-86.
- Mishra RK, Upadhyay VP, Nayak PK, Pattanaik S, Mohanty RC (2012). Composition and Stand Structure of Tropical Moist Deciduous Forest of Similipal Biosphere Reserve, Orissa, India. *Forest Ecosystems – More than Just Trees*. Edited by Juan AB, Yueh-Hsin L. InTech Publishers Janeza Trdine 9, 51000 Rijeka, Croatia.
- Mishra RK, Upadhyay VP, Mohanty RC (2008). Vegetation Ecology of the Similipal Biosphere Reserve, Orissa, India. *Journal of Applied Ecology and Environmental Research* 6(2):89-99.
- Misra R (1968). *Ecology Work-Book*. Oxford and IBH Publishing Co., New Delhi.
- Mohanty RB, Padhy SN, Dash SK (1996). Traditional phytotherapy for diarrhoeal diseases in Ganjam and Phulbani districts of Orissa, India. *Ethnobotany* 8:60-65.
- Natarajan D, Britto SJ, Balaguru B, Nagamurugan N, Soosairaj S, Arockiasamy DI (2004). Identification of conservation priority sites using remote sensing and GIS – A case study from Chitteri hills, Eastern Ghats, Tamil Nadu. *Current Science* 86:1316-1323.
- Nayak S, Bhuyan BK, Satapathy GK, Das S, Kumar V (2016). Phytosociological studies on biodiversity of Bonaigarh forest division,

- Sundergarh district, Odisha. Journal of Plant Development Sciences 8(6):261-273.
- Odisha Wildlife Organization, Official Website of PCCF (WL) & Chief Wildlife Warden, Odisha (2020). www.wildlife.odisha.gov.in
- Oyun MB, Bada SO, Anjah GM (2009). Comparative Analysis of the Floral Composition at the Edge and Interior of Agulii Forest Reserve, Cameroon. Journal of Biological Science 9(5):431-437.
- Parthasarathy N, Sethi P (1997). Tree and liana species diversity and population structure in a tropical dry evergreen forest in south India. Tropical Ecology 38:19-30.
- Philips EA (1959). Methods of Vegetation Study. Henry Holt Co. Inc New York, USA P 107.
- Pradhan B, Rahman CH (2015). Phytosociological study of plant species in three tropical dry deciduous forests of Birbhum District, West Bengal, India. Journal of Biodiversity and Environmental Sciences 7(2):22-31.
- Reddy S, Pattanaik C, Mohapatra A, Biswal AK (2007). Phytosociological Observations on Tree Diversity of Tropical Forest of Similipal Biosphere Reserve, Orissa, India C. Taiwan 52(4): 352-359.
- Rout SD, Panda SK, Panda T (2018). Phytosociological and floristic evaluation of Kuldiha Wildlife Sanctuary, Odisha, India. Tropical Plant Research 5(3):419-430.
- Sahoo AK, Mudgal V (1995). Less known ethnobotanical uses of plants of Phulbani district, Orissa, India. Ethnobotany 7:63-68.
- Sahu SC, Dhal NK, Bhadra AK (2010a). Arboreal Taxa Diversity of tropical forests of Gandhamardan Hill range, Eastern Ghats, India: An approach to sustainable biodiversity conservation. Taiwan 55(3):208-215.
- Sahu SC, Dhal NK, Mohanty RC (2012). Tree species diversity, distribution and population structure in a tropical dry deciduous forest of Malygiri hill ranges, Eastern India. Tropical Ecology 53(2):163-168.
- Sahu SC, Dhal NK, Reddy CS, Pattanaik C, Brahman M (2007) Phytosociological study of tropical dry deciduous forest of Boudh district, Orissa, India. Research Journal of Forestry 1(2):66-72.
- Saxena HO, Brahman M (1989). The Flora of Similipahar, Orissa. Regional Research Laboratory, Bhubaneswar.
- Saxena HO, Brahman M (1996). The Flora of Orissa, Vol I-IV. Orissa Forest Development Corporation Limited, Bhubaneswar. India pp. 89-1735.
- Shahid M, Joshi SP (2016). Phytosociological assessment and distribution patterns of tree species in the forests of Doon Valley, Shivalik hills of lower Himalaya. Tropical Plant Research 3(2):263-271.
- Shiferaw W, Lemenih M, Gole TWM (2018). Analysis of plant species diversity and forest structure in Arero dry Afromontane forest of Borena zone, South Ethiopia. Tropical Plant Research 5(2):129-140.
- Singh KB, Weigand S (1994). Identification of resistant sources in Cicer species to *Liriomyza ciceris*. Genetic Resources and Crop Evolution 41:75-79.
- Sundarapandian SM, Subbiah S (2015). Diversity and tree population structure of tropical dry evergreen forests in Sivagangai district of Tamil Nadu, India. Tropical Plant Research 2(1):36-46.
- Tripathi KP, Singh B (2009). Species diversity and vegetation structure across various strata in natural and plantation forests in Katernia Ghat Wildlife Sanctuary, North India. Tropical Ecology 50(1):191-200.
- Verma MK, Niranjana RK, Pal A (2013). Phytosociological attributes of a tropical dry deciduous forest of Bundelkhand region of Uttar Pradesh, India. Journal of Biodiversity and Environmental Sciences 3(10):86-99.
- Warger MJA, Morrel VE (1976). Plant species and plant communities: Some conclusion. In: Proceedings of the International Symposium, Nijmegen. The Netherlands pp. 167-175.
- Wilson EO (1992). The Diversity of Life, Edward O. Harvard University Press, Cambridge, MA. 464 p. ISBN: 0-674-21298-3.
- WP (2008, 2017). Working Plan for Keonjhar (T) Division and Keonjhar (WL) Division, (2007-08 to 2016-17). Government of Odisha pp. 9-104.