

Full Length Research Paper

Genetic variability of Maure zebu in Mali using SSR markers

**Drissa Konaté^{1*}, Diakaridia Traoré^{1,2}, Sognan Dao¹, Oumar Ouattara¹, Ramata Diop¹,
Aboubacrine Mahamane Toure¹, Adama Konaté¹, Rokiatou Fané¹, Amadou Hamadoun
Babana¹, Youssouf Sanogo¹, Ousmane Nialibouly³ and Mamadou Dougakoro Coulibaly³**

¹Research Laboratory in Microbiology and Microbial Biotechnology (LaboREM-Biotech), Department of Biology, Faculty of Sciences and Techniques, University of Sciences, Techniques and Technologies of Bamako, Bamako, Mali.

²National Center for Artificial Animal Insemination, Bamako, Mali.

³Institute of Rural Economy, Niono, Mali.

Received 16 June, 2021; Accepted 3 December, 2021

In Mali, Maure's zebu breed has good dairy production abilities. Females are often used as a matrix in crosses in peri-urban areas with exotic breeds. Despite the enthusiasm of its productive abilities, the genetic diversity of this breed is not yet sufficiently assessed. In order to contribute to a better understanding of the genetic diversity of this breed, 82 animals were chosen from the farms of four districts in two regions in Mali for genetic evaluation, using Simple Sequence Repeats (SSR) markers. Results from the investigations identified a total of 41 alleles with an average of 3.15 alleles per locus. The Polymorphism Information Content (PIC) ranged from 0.293 (ILSTS011) to 0.786 (HEL5) with a mean of 0.528. By locus, the means of observed heterozygosity (Ho) and expected heterozygosity (He) were 0.400 ± 0.032 and 0.416 ± 0.026 , respectively. Likewise, by subpopulation, their values varied respectively from 0.379 ± 0.057 (Kayes) to 0.417 ± 0.070 (Niono) for the observed heterozygosity and from 0.360 ± 0.068 (Baraouéli) to 0.454 ± 0.049 (Diéma) for the expected heterozygosity. The overall mean of genetic differentiation (Fst) was moderate, indicating 9% of the genetic deviation corresponded to differences between subpopulations. This study will strengthen knowledge on the genetic diversity of Maure zebu for the better strategy adoption for its conservation and promotion.

Key words: Maure zebu, genetic, variability, SSR markers, Mali.

INTRODUCTION

Mali has a large herd of cattle in West Africa with the majority concentrated in areas above 600 mm isohyet. Malian cattle breeds are characterized by low production potential (Lhoste, 1983; Nialibouly, 1999), but have on the other hand a great adaptability to the harsh climatic conditions of the Sahel and Sudanian zones. The cattle

are comprise zebus (*Bos indicus*) concentrated in the north and center of the country, taurines (*Bos taurus*) and the products of zebu-taurine from crosses found in the south of the country (Planchenault, 1988). Among the zebus, the Maure zebu occupies a place of choice among the population who rear it. It is a rather rambling animal,

*Corresponding author. E-mail:konatedrissa80@gmail.com.

with strong bones (FAO, 1957; Planchenault, 1988). Its limbs are quite strong and well adapted to walking (Planchenault, 1988). The horns are thin, short in the male but longer in the female, they are round in section and greyish or brownish in color (Ministry of Livestock and Fisheries, 2016). The coat is generally dark red, (Planchenault, 1988) but sometimes it could be black and pie-black coats, white on the muzzle on the belly (Ministry of Livestock and Fishing, 2016). The height of the withers varies from 1.40 to 1.50 m for the male and from 1.25 to 1.30 m for the female. The thoracic perimeter varies from 1.60 to 1.85 m with a scapulo-ischial length varying from 1.24 to 1.50 m (Ministry of Livestock and Fisheries, 2016). The weight of the breed is between 250 and 400 kg. Daily milk production varies from 2 to 3 kg reaching 8 kg during good seasons (Nialibouly et al., 2003). Its natural habitat straddles the border between Mali and Mauritania (Ministry of Livestock and Fisheries, 2016). This breed is also known under the following names: Arab zebu, Gabarouyé zebu and Mauritanian zebu (FAO, 1957). In peri-urban rearing, Maure females are increasingly used as a matrix in crosses with exotic breeds for milk production. Good knowledge of the genetic diversity of this breed will enable Malian technicians and breeders to better design genetic improvement and management programs for the Maure zebu. The objective of this study is to assess the genetic diversity of Maure zebu from four districts in two different regions of Mali.

MATERIAL AND METHODS

Ethical considerations

This study titled “Genetic variability study of Maure zebu in Mali using Simple Sequence Repeats (SSR) markers” complies with the Agricultural Orientation Law, the National Policy for Development of Livestock, the National Priority Investment Program for the Agricultural Sector, the Pastoral Charter and the National Strategy for Conservation, Selection and Dissemination of Native Breeds. The study was approved by the ethics committee of the National Institute for Public Health Research under N° 17/2019 / CE-INRSP.

Sampling site

Sampling was carried out in the farms of the Circles of Kayes, Diéma (region of Kayes), Baraouéli and the station of the Regional Agronomic Research Center of Niono (region of Ségou). These sites were chosen according to their accessibility, availability of cow breeders and their herds. The animals were chosen according to the morphological criteria of the breed. A maximum of 5 individuals were sampled per farm. The sites constitute breeding and transit areas for transhumant to the country's wetlands. A sample of 82 heads including 32 from Kayes district, 30 from Diéma district, 8 from Baraouéli district and 12 from Niono district were selected for genetic diversity investigations.

Herd management

The herds were led in an extensive breeding system through the

exploitation of natural ranges during the twelve months of the year for the peasant farms of Kayes, Diéma and Baraouéli and in semi-intensive breeding for the animals of the Regional Agronomic Research Center of Niono (CRRRA-Niono). Some peasant herds received crop residues and concentrates (cereal bran) during dry periods in addition to grazing. The animals at the Niono CRRRA received supplementation (food concentrates), crop residues from rice during the dry season. The health interventions were carried out by the technical breeding agents, if necessary, in the peasant farms. At the Niono CRRRA, sanitary prophylaxis was applied to the herd. Mating was natural in all herds. A sire was used for reproduction.

Collection of blood samples

Four milliliters (4 ml) of blood were taken from the jugular vein in EDTA tubes, from males and females randomly selected from the herds. The blood samples were labeled, kept under ice, and transported to the Research Laboratory in Microbiology and Microbial Biotechnology (LaboREM-Biotech) of the Faculty of Sciences and Techniques (FST) of the University of Sciences, Techniques and Technologies of Bamako (USTTB) for molecular analyses.

Extraction of genomic DNA

Zebu genomic DNA samples were extracted from whole blood using the Promega ReliaPrep™ Blood gDNA Miniprep System Extraction Kit. The concentration of the DNAs extracted was determined using the EPPENDORF BioPhotometer® spectrophotometer 6131. DNAs were diluted to 20ng / µl and stored at 4°C for analyzes.

DNA amplification

DNA samples were amplified according to Konaté et al. (2018) with 13 microsatellite primer pairs (Table 1). The markers were chosen from the panel recommended by the International Society of Animal Genetics (ISAG) and the Food and Agriculture Organization (FAO, 2011) for genetic diversity analyzes. These markers have been used in cattle, goat and sheep characterizations (Gororo et al., 2018; Konaté et al., 2018; Al-Atiyat, 2016; Hussain et al., 2016). They are co-dominant, reproducible and polymorphic (Lirón et al., 2002; Portetelle et al., 2000) to generate vital information on the genetic variability of cattle.

Electrophoresis of amplified products

The gels were prepared with 0.5X TBE (Tris, Borate Acid, EDTA) and 30µl of 10% ethidium bromide (1mg/ml) (Konaté et al., 2018). The MS4 CONDA 3% agarose gel was loaded with 10 µl of each PCR product and migrated for 2 h 30 min at 80V. The gels were photographed with the Gel Documentation System E-BOX VX2 version 15.06.

Statistical analysis

The band (alleles) sizes of each microsatellite primer were determined in base pair using E-Capt software version 15.06. The genetic diversity of each locus was assessed based on the following statistical parameters: number of alleles, frequency of alleles, number of genotypes, genetic diversity, and PIC (Polymorphism Information Content) using Power Marker software

Table 1. Names, sequences and repeating patterns of the microsatellite primers used

SSR name	Chrs		Primers	References
BM1824	1	Forward	GAGCAAGGTGTTTTTCCAATC	Bishop et al., 1994
		Reverse	CATTCTCCAAGTCTTCCTTG	
INRA023	3	Forward	GAGTAGAGCTACAAGATAAACTTC	Vaiman et al., 1994
		Reverse	TAACACAGGGTGTAGATGAACTCA	
MNB42	4	Forward	CTTTAGCTGTTCAGCAGATGG	Hassane, 2011
		Reverse	TGATTCTCAGCTCTTTCTAGCC	
ILSTS034	5	Forward	TACTCGTAGGGCAGGCTGCCTG	Sodhi et al., 2011
		Reverse	GAGACCTCAGGGTTGGTGATCAG	
BM143	6	Forward	ACCTGGGAAGCCTCCATATC	Vaiman et al., 1994
		Reverse	CTGCAGGCAGATTCTTTATCG	
ILSTS006	7	Forward	TGTCTGATTTCTGCTGTGG	Armstrong et al., 2006
		Reverse	ACACGGAAGCGATCTAAACG	
MM12	9	Forward	CAAGACAGGTGTTTCAATCT	Mommens et al., 1994
		Reverse	ATCGACTCTGGGGATGATGT	
ILSTS005	10	Forward	GGAAGCAATGAAATCTATAGCC	Brzeziński and Majid, 1993.
		Reverse	TGTTCTGTGAGTTTGTAAAGC	
INRA37	11	Forward	GATCCTGCTTATATTTAACCAC	Vaiman et al., 1994
		Reverse	AAAATTCATGGAGAGAGAAAC	
ILSTS011	14	Forward	GCTTGCTACATGGAAAGTGC	Brzeziński and Majid, 1993
		Reverse	CTAAAATGCAGAGCCCTACC	
TGLA 53	16	Forward	GCTTTCAGAAATAGTTTGCATTCA	Georges and Massey, 1992
		Reverse	ATCTTCACATGATATTACAGCAGA	
TGLA 122	21	Forward	CCCTCCTCCAGGTAAATC AGC	Georges and Massey, 1992
		Reverse	AATCACATGGCAAATAAGTACATAC	
HEL5	21	Forward	GCAGGATCACTTGTAGGGA	Kaukinen and Varvio, 1993
		Reverse	AGACGTTAGTGATACATTAAC	

Chrs: Chromosome; **SSR:** Simple sequence repeat.

version 3.25. The observed (H_o) and theoretical or expected (H_e) heterozygosity, the overall inbreeding coefficient (F_{it}), the inbreeding coefficient (F_{is}), the standardized variance (F_{st}) were calculated using the GenAEx software version 6.5. Principal component analysis based on genetic distance was performed with the same software.

RESULTS

Molecular analysis of 82 Maure zebus blood, from four circles (Kayes, Diéma, Baraouéli and Niono) in Mali with thirteen (13) microsatellite markers, revealed a total of 41 alleles forming 73 different genotypes (Figure 1). The number of alleles ranged from 2 (ILSTS006, ILSTS011 and INRA023) to 5 (TGLA122) with an average of 3.15 alleles per locus (Table 2). The PIC ranged from 0.293 (ILSTS011) to 0.786 (HEL5) with a mean of 0.528. Majority of SSR markers exhibited a mean PIC greater than or equal to 0.50. The observed genetic diversity varied from 0.356 to 0.812 respectively for the ILSTS011 and HEL5 loci with an average of 0.576. Values for H_o and H_e were calculated for each locus and subpopulation

under Hardy Weinberg equilibrium. Thus, H_o varied from 0.241 (INRA37) to 0.889 (BM143) with a mean of 0.400 ± 0.032 . By subpopulation, it varied from 0.379 ± 0.057 (Kayes) to 0.417 ± 0.07 (Niono) (Table 4). Likewise, H_e varied from 0.229 (ILSTS005) to 0.626 (MM12) with a mean of 0.416 ± 0.026 (Table 2). Its values per subpopulation from Baraouéli and Diéma varied respectively from 0.360 ± 0.068 to 0.454 ± 0.049 (Table 4).

The markers BM143 and INRA37 loci showed a significant deviation from the Hardy-Weinberg equilibrium in Kayes, Diéma and Niono subpopulations. The MNB42 locus showed a significant deviation from the Hardy-Weinberg equilibrium in Kayes, Diéma and Baraouéli subpopulations. The ILSTS034 locus was significant in Kayes and Diéma subpopulations. Significant deviation from the Hardy-Weinberg equilibrium was noticed with TGLA122 and HEL5 loci in the respective subpopulations of Diéma and Niono (Table 5). The subpopulations of Kayes, Diéma, Baraouéli and Niono presented respective fixation indices $F = 0.033 \pm 0.082$, $F = 0.049 \pm 0.075$, $F = -0.097 \pm 0.071$ and $F = 0.074 \pm 0.148$ with an average of




Figure 1. Profiles of the TGLA122 (A) and MNB42 (B) SSR markers after migration on a 3% metaphor gel. **M:** Size marker (in base pair), **N:** Negative control; **Z-22 to Z-49:** Maure zebu samples.

Table 2. Number of alleles, allele frequency, genetic diversity and PIC of Maures zebus from Mali.

Marker	Nbre d'allèle	Taille en pb		Gene Diversity	PIC	Ho	He
		Mini	Max				
BM143	4	95	203	0.358	0.340	0.889	0.539
BM1824	3	179	191	0.450	0.385	0.262	0.277
HEL5	3	150	168	0.812	0.786	0.341	0.585
ILSTS005	3	183	204	0.438	0.376	0.283	0.229
ILSTS006	2	196	324	0.485	0.379	0.376	0.296
ILSTS011	2	264	293	0.356	0.293	0.313	0.247
ILSTS034	3	161	234	0.771	0.742	0.447	0.553
INRA023	2	203	205	0.442	0.384	0.242	0.240
INRA37	3	112	137	0.541	0.506	0.241	0.319
MM12	4	112	141	0.762	0.743	0.465	0.626
MNB42	4	68	126	0.706	0.654	0.483	0.592
TGLA53	3	184	250	0.682	0.625	0.562	0.503
TGLA122	5	143	174	0.682	0.657	0.295	0.399
Mean	3,15			0.576	0.528	0.400±0.032	0.416±0.026

PIC: Polymorphism Information Content, **Ho** = Heterozygosity observed, **He** = Heterozygosity expected.

0.017 ± 0.049. These positive indices reflect a deficit of heterozygosity in the subpopulations, while the negative index indicates an excess of heterozygosity (Table 3).

Wright's F-statistics F_{is} , F_{it} and F_{st} calculated for all Maure zebu populations in the study area were 0.005 ± 0.083, 0.087 ± 0.088 and 0.090 ± 0.022, respectively (Table 5). The overall mean of genetic differentiation (F_{st}) was moderated indicating that 9% of the genetic deviation corresponded to the differences between populations. All of the amplified microsatellite markers contributed to genetic differentiation (F_{st}) in sub-

populations. However, two microsatellite markers INRA023 (31.3%) and TGLA122 (15.3%) showed significant differentiations. The mean F_{is} revealed a deficit of heterozygosity in overall population. However, loci BM143, ILSTS005, ILSTS006, ILSTS011, INRA023 and TGLA53 showed excess heterozygosity with negative F_{is} values (Table 5).

Analysis of Molecular Variance (AMOVA) revealed high genetic variation within individuals (97.33%). The analysis of the principal coordinates revealed a percentage of genetic variation of 12.76 and 24.34% respectively for

Table 3. Observed, expected, non-biased heterozygosity and fixation index in the population of Kayes, Diéma and Baraouéli.

Population	Ho	He	F
KAYES	0.379±0.057	0.411±0.054	0.033±0.082
DIEMA	0.406±0.048	0.454±0.049	0.049±0.075
BARAOUELI	0.397±0.086	0.360±0.068	-0.097±0.071
NIONO	0.417±0.07	0.438±0.035	0.074±0.148
Mean	0.400±0.032	0.416±0.026	0.017±0.049

Ho = Heterozygosity observed, He = Heterozygosity expected, F = Fixation index.

Table 4. Summary of chi-square tests for Hardy-Weinberg equilibrium by Maure zebu population of Mali.

Locus	Populations			
	Kayes	Diéma	Baraouéli	Niono
BM143	*	***	ns	***
BM1824	ns	ns	ns	ns
HEL5	**	ns	ns	*
ILSTS005	ns	ns	monomorphic	ns
ILSTS006	ns	ns	ns	ns
ILSTS011	ns	ns	ns	ns
ILSTS034	***	*	ns	ns
INRA023	ns	ns	ns	ns
INRA37	**	*	ns	***
MM12	**	***	ns	ns
MNB42	***	***	*	ns
TGLA53	ns	ns	ns	ns
TGLA122	ns	**	ns	ns

*: P <0.05, **: P <0.01, ***: P <0.001, ns: Négatif.

Table 5. F-statistics of whole populations for each locus.

Locus	Fis	Fit	Fst
BM143	-0,650	-0,631	0,011
BM1824	0,054	0,065	0,012
HEL5	0.417	0.483	0.113
ILSTS005	-0.233	-0.139	0.076
ILSTS006	-0.267	-0.191	0.060
ILSTS011	-0.268	-0.186	0.065
ILSTS034	0.192	0.268	0.094
INRA023	-0.009	0.307	0.313
INRA37	0.244	0.290	0.061
MM12	0.256	0.320	0.086
MNB42	0.184	0.278	0.114
TGLA53	-0.116	-0.110	0.006
TGLA122	0.260	0.373	0.153
Mean	0.005±0.083	0.087±0.088	0.090±0.022

Fis: fixation index, Fit: global differentiation, Fst: differentiation.




Figure 2. Principale component analysis of SSR genotypes of 82 Maure zebu.

the first two principal axes with a total of 37.10% (Figure 2). The graph generated by the first two axes, showed individual's dispersion and do not form specific homogeneous groups. Nevertheless, a mixture of three subpopulations (Kayes, Diéma and Baraouéli) was observed. On the other hand, the subpopulation of Niono was different to that of Kayes, Diéma and Baraouéli.

DISCUSSION

Genetic diversity study of 82 Maure zebus revealed a total of 41 alleles with an average of 3.15 alleles per locus. The mean PIC and the mean genetic diversity were 0.528 and 0.576, respectively. These values show a strong diversity within the Maure zebus coming from the various localities of Mali. Study conducted by N'diaye et al. (2015) in Senegal on zebus from Saint-Louis and Kaolack revealed 81 alleles with an average of 7.364 ± 0.544 alleles per locus and an average PIC of 0.719. These values are clearly higher than those obtained in the present study and also showed a very high genetic variability of Maure zebus. All the amplified loci were polymorphic based on the mean of the observed PICs. However, 53.85% of the amplified markers showed a PIC greater than or equal to 0.5. On the other hand, this rate is lower than those of N'diaye et al. (2015) on Maure zebus in Senegal, with an average PIC of 100%, greater than 0.5.

The observed mean H_o and expected H_e recorded at the target loci were 0.400 ± 0.032 and 0.416 ± 0.026 , respectively. By subpopulation, H_e varied from 0.360 ± 0.068 (Baraouéli) to 0.454 ± 0.049 (Diéma) with an average of 0.416 ± 0.026 . N'diaye et al. (2015) obtained an average value of 0.769 ± 0.099 on Maure zebus originating from two localities in Senegal. However, in Kayes, Diéma and Niono subpopulations, the observed heterozygosity was less than the expected heterozygosity suggesting a deficit in heterozygosity. Population from Baraouéli showed a great H_o than the H_e indicating an excess of heterozygosity. The high rate of heterozygosity

in this locality could be linked to the contact with other varieties of zebu, the Fulani zebu in particular.

The mean value of the fixation index (F_{is}) of all the loci was 0.005 ± 0.083 , determining a lower heterozygosity deficit compared to the overall population fixation index ($F_{it} = 0.087 \pm 0.088$). The BM1824, HEL5, ILSTS034, INRA37, MM12, MNB42 and TGLA122 SSR markers contributed to this deficit, through positive F_{is} values. As for the F_{st} genetic differentiation, it is relatively very important for the INRA023 locus. In contrast, F_{st} displays low values at the BM143, BM1824, TGLA53 locus levels. The mean differentiation is moderate between subpopulations with a value of 0.090 ± 0.022 indicating the origin of the total genetic variation in the breed.

Principal component analysis (PCA) revealed a mixture of genotypes of the Maure zebu populations of Kayes, Diéma and Baraouéli. Genotypes from Niono were separated from those of other localities. This could be explained by the fact that these animals were bred and selected for several decades for the production of milk. These results reflect the geographic distance between the different districts where the samples were collected. Nearby districts have animals with high genetic similarity. The districts of Kayes and Diéma are contiguous. On the other hand, the district of Niono is more than 700 km away from that of Diéma and Kayes.

Conclusion

The molecular characterization of Maure zebu in Kayes, Diéma, Baraouéli and at the station of Niono showed a genetic variability at the loci level. The results from zebu in Kayes and Diéma presented lower observed heterozygosity than the expected one. This supposes that the flock is not subject to genetic introgression from other breeds. In Baraouéli, the observed heterozygosity was higher than the expected one. This can be explained by a possible contact of Maure zebu with Fulani zebu. Baraouéli constitutes the breeding area of Fulani zebu. The obtained results can contribute to build

conservations and breeding program of Maure zebu in Diéma, Kayes and Niono.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

ACKNOWLEDGMENTS

The authors appreciate the Competitive Fund for Research and Technological Innovation for its financial support. They are also grateful to the technical livestock services: Local Service for Animal Productions and Industries, Support Units for Animal Productions and Industries, Veterinary Services, as well as the breeders for their support and collaboration.

REFERENCES

- Al-Atiyat RM (2016). Association of allele diversity and polymorphism of microsatellites markers in the tropical goat. *Research Journal of Biotechnology* 11(9):29-36.
- Armstrong E, Postiglioni A, Martínez A, Rincón G, Vega-Pla JL (2006). Microsatellite analysis of a sample of Uruguayan Creole bulls (*Bos taurus*). *Genetics and Molecular Biology* 29(2):267-272.
- Bishop MD, Kappes SM, Keele JW, Stone RT, Sunden SL, Hawkins GA, Yoo J (1994). A genetic linkage map for cattle. *Genetics* 136(2):619-639.
- Brzeziński T, Majid S (1993). Quantum group gauge theory on quantum spaces. *Communications in Mathematical Physics* 157(3):591-638.
- Food and Agriculture Organization (FAO) (1957). *African Cattle, Types and Breeds*. FAO Agricultural Studie, Rome (37):56-60.
- Food and Agriculture Organization (FAO) (2011). *Molecular genetic characterization of animal genetic resources*. FAO Animal Production and Health Guidelines 9 Rome.
- Georges M, Massey JM (1992). Polymorphic DNA markers in Bovidae. Patent WO 92/13102.
- Gororo E, Makuza SM, Chatiza FP, Chidzondo F, Sanyika TW (2018). Genetic diversity in Zimbabwean Sanga cattle breeds using microsatellite markers. *South African Journal of Animal Science* 48(1):128-141.
- Hassane YA (2011). Application of microsatellite markers polymorphism for genetic density study of zebus in Niger. Master thesis, University of Polytechnic, Bobo Dioulassao 87, Burkina Faso.
- Hussain T, Babar ME, Peters SO, Wajid A, Ali A, Azam A, De Donato M (2016). Microsatellite Markers Based Genetic Evaluation of Pakistani Cattle Breeds. *Pakistan Journal of Zoology* 48(6):1633-1641.
- Kaukinen J, Varvio SL (1993). Eight polymorphic bovine microsatellites. *Animal Genetics* 24(2):148.
- Konaté D, Traoré D, Dao S, Fané R, Ouattara O, Diop R, Babana A (2018). Genetic Diversity Study of N'Dama Breed in Mali Using SSR Markers. *International Journal of Animal Science and Technology* 2(3):30-35.
- Lhoste P (1983). Development of animal traction and evolution of pastoral systems in Siné-Saloum, Senegal (1970-1981). *Tropical Animal Husbandry and Veterinary Medicine Review* 36(3):291-300.
- Lirón JP, Ripoli MV, De Luca JC, Peral-García P, Giovambattista G (2002). Analysis of genetic diversity and population structure in Argentine and Bolivian Creole cattle using five loci related to milk production. *Genetics and Molecular Biology* 25(4):413-419.
- Ministry of Livestock and Fisheries (2016). *Directory of cattle, sheep and goat breeds in Mali* pp. 31-33.
- Mommens G, Coppieterst W, Weghe A, Zeveren AV, Bouquet Y (1994). Dinucleotide repeat polymorphism at the bovine MM12E6 and MM8D3 loci. *Animal Genetics* 25(5):368-368.
- N'diaye NP, Sow A, Dayo GK, Ndiaye S, Sawadogo GJ, Sembène M (2015). Genetic diversity and phylogenetic relationships in local cattle breeds of Senegal based on autosomal microsatellite markers. *Veterinary World* 8(8):994-1005.
- Nialibouly O (1999). Production and reproduction performance of Maure and Fulani zebus (1986–1995) improving at the Niono Agronomic Research Station in Mali: Master thesis, Cheikh Anta Diop University of Dakar, Faculty of Sciences and technologies P 128.
- Nialibouly O, Coulibaly MD, Dolo D (2003). Improving Dairy Production Performance of Maure and Peul zebu by Selecting at the Niono Station, Mali. *Rural Economy Notebooks*, Issue (0):3
- Planchenault D (1988). L'élevage In : *Animal husbandry and sahelian pastoral potentialities*. Cartographic synthesis. Mali. CIRAD-IEMVT - FRA. Wageningen: CTA-CIRAD-IEMVT, 26-27.
- Portetelle D, Haezebroeck V, Mortiaux F, Renaville R (2000). Traçabilité dans la filière animale. *Biotechnologie Agronomie Société et Environnement* 4(4):233-240.
- Sodhi M, Mukesh M, Mishra BP, Ahlawat SPS, Prakash B, Sobti RC (2011). Microsatellite analysis of genetic population structure of zebu cattle (*Bos indicus*) breeds from North-Western region of India. *Animal Biotechnology* 22(1):16-29.
- Vaiman D, Mercier D, Moazami-Goudarzi K, Eggen A, Ciampolini R, Lépingle A, Lévézuel H (1994). A set of 99 cattle microsatellites: characterization syteny mapping and polymorphism. *Mammalian Genome* 5(5):288-297.