

Full Length Research Paper

Tillage and manure effect on soil physical and chemical properties and on carbon and nitrogen mineralization potentials

H. Kheyrodin^{1*} and H. Antoun²

¹Department of Natural Resources Faculty of desertification of Semnan University, Semnan-Iran.

²Département des sols et de génie agroalimentaire Pavillon Paul-Comtois, local 2211 Université Laval, Canada.

Accepted 28 September, 2009

The objective of this work was to study the effects of tillage and liquid manure applications on some physical and chemical properties and also on the carbon and nitrogen mineralization potential from a meadow soil. Our results indicated that tillage and manure applications had no effect on the concentration of total N and organic C in the 0 - 15 cm layer of soil after 15 years of treatment. However soil P, Ca, Mg and Zn contents increased significantly with manure applications. Soil organic matter and total N significantly decreased in the 15 – 30 cm depth. No significant change could be detected in soil structural stability in both layers. Moreover, tillage affected significantly soil soluble C and the C/N ratio. Application of 100 t ha⁻¹ manure significantly increased soil soluble C. The results of this study suggest that tillage increased significantly the soil N mineralization rate. The potentially mineralizable nitrogen (N₀) was higher in tilled than in no-tilled soil and was at its maximum in the 0 – 15 cm layer of the soil. Furthermore, a significant positive interaction was observed between tillage and manure application on N mineralized after 1.4 wk (N_e). No significant change was detected in both C mineralization rate (C_m) and potentially mineralizable C (C₀). The total amounts of mineralizable carbon (C_m) and nitrogen (N_m) significantly decreased in 15 – 30 cm depth and were very closely correlated with the total amounts of C or N and mineralization rate constants (K).

Key words: Carbon and nitrogen mineralization potentials, k constants, physical and chemical properties.

INTRODUCTION

Soil biological properties play an important role in the transformation and the cycling of plant nutrients, especially carbon and nitrogen. In general, soil organic matter is the main source of nitrogen by means of microbial mineralization. Tracy et al. (1990) found for winter wheat that no-tilled soils accumulated greater nitrate ion (NO₃-N), Ammonium ion (NH₄-N), Phosphate ion (PO₄-P) at 0 - 2.5 cm soil depth, than did recently plowed soil. Below the 5 cm depth, tillage did not influence net nitrogen, phosphorus and phosphorus mineralization. There were higher CO₂-C evolution rates early of the incubation in soil from no-till, compared with those measured after plowing

treatments, but at the end of the incubation period all treatments were releasing similar amounts of CO₂-C (Tracy et al., 1990). El-Haris et al. (1983) determined nitrogen mineralization potential (N₀) in a soil from a long-term crop rotation tillage experiment and found that the average N₀ was unaffected by tillage or crop rotation in the 0 – 15 cm depth when sampling was performed in the fall. In the spring sampling, average N₀ for either chisel plowing or no-till was significantly higher than for moldboard plowing. Moldboard tillage also resulted in significantly lower organic C and a narrower C/N ratio in the top 15 cm relative to the no-tilled treatment (El-Haris et al., 1983). Bennett et al. (1975) also found that N₀ in the surface 15 cm layer was higher for the no-tilled than for conventionally tilled corn (*Zea mays*) plots. Carter and Rennie (1982) reported that potential net mineralizable C and N were significantly greater in surface soil under

*Corresponding author. E-mail: hkheyrodin@yahoo.com. Tel: 00989121029221. Fax: 00982188716603.

zero tillage in comparison to conventional tillage.

Animal manure supplies additional mineralizable C and N that directly stimulate microbial activity and growth (McGill et al., 1986). Manure improved soil fertility by increasing the labile organic N constituent such as the amino compounds. This was reflected by a higher N-supplying power in manure soils (Campbell et al., 1986). The rate of net mineralization of N depends on manure composition and on soil physical, chemical and biological properties. If animal manure is applied regularly in large quantities over a long period of time, it increases the soil organic matter content, soil porosity, mineralizable N, Na-HCO₃-extractable-P, microbial activity and reduce the bulk density (Sommerfeldt and Chang, 1985, Spratt and McCurdy, 1966, Sweeten and Mathers, 1985).

Potentials of N mineralization values can be calculated based on the hypothesis that the rate of N mineralization is proportional to the quantity of N comprising the mineralizable substrate (Stanford and Smith, 1972). However, N₀ is affected by many factors such as moisture, aeration, temperature, nature and quantity of organic matter, nature and quantity of the previous crop residues as well as other soil physical, chemical, and biotic properties (Ellert and Bettany, 1992; Simard and N'dayegamiye, 1993).

Tillage and manure application influence soil moisture, soil temperature, pH, soil organic matter (OM) distribution and soil physical, chemical and biological properties and thus their rates (K_n, K_c) and potentials of C and N mineralization (Flowers and Arnold, 1983; Doran, 1980a, b; N'dayegamiye and Coté, 1989).

The objective of this study was to determine the influence of tillage and pig liquid manure application on soil physical and chemical properties and on C and N mineralization potential in a meadow soil in Quebec, Canada.

MATERIALS AND METHODS

Soil sampling

This study was conducted in 1994 for three years on a silt-loam at the MAPAQ (Ministère de l'Agriculture, des Pêcheries et de l'Alimentation du Québec) experimental farm in St. Lambert, Quebec, Canada. Treatments were in a split-plot design consisting of no-till and tillage as main treatments and three liquid manure rates (0, 50, 100 t ha⁻¹) as secondary treatments. Since 1978, some plots have never been plowed down (no-tilled treatments) whereas in the tilled treatment, soil was plowed in the fall every 5 years. Liquid manure was applied in the spring of each year. Soil sampling for this study was performed in June 1994 and in July 1995. Soil samples were taken from 0 – 15 cm and 15 – 30 cm depths. The moist soil samples were sieved (> 6 mm) to remove roots and stubble in the field and were kept at 4°C until performing both physical and microbial activity analyses. For chemical analysis, sub-samples were air-dried and screened at 2 mm and 250 µm.

Physical analyses

Distribution of aggregate size was carried on samples sieved at 6 mm. Structural stability measurements were determined by wet sieving (> 2 mm; > 1 mm; > 0.5 and 0.25 mm) on moist soil as described by Angers and Mehuys (1993).

Chemical analyses

Soil pH was measured in a soil : water ratio of 1:2. The organic C content was determined by wet oxidation procedure (Walkley and Black, 1934). Total N was estimated by Kjeldahl digestion (Nelson and Sommers, 1982). Macro- and micronutrients were extracted using the Mehlich-III solution (Mehlich, 1984). Soil water-soluble C was assessed as previously described by Dormaar et al. (1984).

Nitrogen and carbon mineralization measurements (Incubation studies)

Field-moist samples (400 g) were incubated for 270 days in 4-L cylinders to ensure optimum aeration and microbial activity (Stanford and Smith, 1972). Carbon dioxide evolution was trapped in 1N NaOH (5 mL) solution and the excess of NaOH was titrated with HCL 1M (5 mL) (Anderson, 1982). For N mineralization analysis, the soil samples were leached as described by Stanford and Smith (1972), with 100 mL of 0.01 M CaCl₂ followed by 25 mL of N-free nutrient solution (0.002 M CaSO₄·2H₂O; 0.002M MgSO₄; 0.005M Ca(H₂PO₄)₂·H₂O; and 0.0025M K₂SO₄). Nitrogen-N was determined colorimetrically on a Technicon Autoanalyzer.

Data analyses

Analysis of variance for determining the effects of both tillage and manure upon soil biological, physical and chemical characteristics was determined by the general linear model procedures (GLM) of the Statistical Analyses System (SAS, 2004). The multiple comparison test was carried out with protected Fisher's least significant difference (LSD). Linear correlations were calculated between physical and chemical characteristics and between C and N mineralization parameters. The soil carbon and nitrogen mineralization potentials (C₀ and N₀) were determined by the cumulative model of Ellert and Bettany (1988), $I_{ct} = M_0 (1 - e^{-kt})$ where I_{ct} is the cumulative amount of C and N mineralized (mg kg⁻¹) after time t (in wk) and M_0 is the potentially mineralizable C or N and k is the first order rate constant (week⁻¹).

RESULTS

Soil Physical Properties

Tillage and manure application did not have any significant effect on soil aggregates distribution. The observed soil structural stability determined by Mean Weight Diameter (MWD) was generally low for this type of soil and was not influenced by tillage or manure application.

Soil chemical properties

Some chemical properties of the soil samples from the

Table 1. Analysis of variance the effect of tillage and liquid manure on some chemical properties of a meadow soil.

Treatment	pH (H ₂ O)	P	K	Ca	Mg	Cu	Mn	Zn
Tillage (T)	0.62	4.45**	9.7**	2.64	2.98	4.75	3.65	39.2**
Manure (M)	6.65**	27.3**	0.75	5.66*	5.61*	1.64	1.25	31.6**
Depth (D)	0.08	6.46*	1.59	24.4**	1.08	9.43	0.11	87.9**
T X M	0.41	4.18*	0.98	2.5	1.62	1.96	1.64	23.1**
T X D	0.06	6.13*	0.09	0.0	0.06	4.33	0.13	21.7**
D X M	0.32	5.81**	1.89	1.72	1	1.81	0.95	26.3**
T X M X D	0.58	5.42*	0.14	0.39	1.01	0.68	0.95	16.4*

** Significant at $P < 0.01$.

*: Significant at $P < 0.05$.

T: Tillage, M: Manure, D: Depth, T X M: Tillage * Manure, T X D: Tillage * Depth, D X M: Depth * Manure.

T X M X D: Tillage * Manure * Depth.

different tillage and liquid manure treatments are presented in Table 1. Liquid manure application significantly increased soil P ($P \leq 0.01$), Ca, Mg and Zn levels. In the absence of manure application tillage significantly reduced surface soil (0 - 15) P content. However, surface manured plowed soils had P levels that were higher than those observed in no-tilled soils. At the 15 – 30 cm depth, tillage did not affect soil P content. In both tilled and notilled soils the addition of 50 t ha⁻¹ liquid manure slightly increased soil pH.

Soil organic C was not affected by the addition of manure and was generally present at higher concentration in the surface soil (0 – 15 cm). In the absence of manure treatment the surface soil under no-tillage contained significantly more carbon than under tillage

Soil C and N Levels and C and N Mineralization

Results on the influence of tillage and liquid manure upon soil C and N levels and mineralizable C and N data are presented in Table 2. Contents of organic carbon are maximum in the upper layer (0 – 15 cm) of the no-tilled treatment that did not receive manure application and the upper layer of the tillage treatment with 100 t ha⁻¹ manure application. The soil N contents significantly decreased in 15 – 30 cm depth. No effect of tillage and manure application was observed for soil C and N contents ($P > 0.05$). Without manure application, tillage slightly increased soil soluble carbon (Cs). However, soils receiving 100 t ha⁻¹ liquid manure had the highest content of Cs which was significantly reduced by tillage. The C mineralized over 10 days (Ce) significantly decreased in 15 – 30 cm depth. Tillage increased significantly the net mineralized nitrogen (Nm) and the N mineralized over 1.4 wk. Cumulative net mineralized N and C showed a curvilinear relationship with time. Figures 1 and 2 illustrate the effect of tillage

and liquid manure application on the cumulative C and N mineralized in 270 days incubation.

Soil C and N mineralization potentials

Results from on the effect of tillage and liquid manure on the potentially mineralizable C and N (C₀ and N₀) are shown in Table 2. Tillage and manure treatments did not have any significant effect on C₀ and N₀ or on their mineralization rate constants K_c and K_n.

We did not observe any effect of manure application on the initial N mineralization (N mineralization over 10 days, N_e) or on the total amount of N mineralized (after 270 days incubation, N_m). The rate constant (K_n) values calculated with this model ranged from 0.0045 to 0.0458 day⁻¹. Nitrogen mineralization potential (N₀) was affected by many factors as soil moisture, aeration, temperature, nature and quantity of organic matter, nature and quantity of the previous crop residues.

Relations between some chemical properties and C and N mineralization parameters

The pH was negatively related to the rate constant of N mineralization ($P < 0.01$). Soil C content is significantly related to net mineralized N (N_m; $P < 0.05$) and initially mineralized N (N_e; $P < 0.01$). Also the soil N content was highly correlated to N_m ($P < 0.01$) and N_e ($P < 0.01$). The soil structural stability (MWD) was not related to nitrogen mineralization parameters. The soil N content was highly correlated to the total amount of C mineralized (C_m; $P < 0.01$). The soluble C is significantly related to potentially mineralizable C (C₀; $P < 0.01$). The soil structural stability (MWD) was not related to carbon mineralization parameters.

Table 2. The effect of tillage and manure on carbon and nitrogen mineralization parameters and on some chemical characteristics of a meadow soil.

Treatment	C	N	C _s	C/N	OM/C _s	N _m	C _m	C _e	N _e
Tillage (T)	3.39	1.02	14.68**	13.4 **	0.00	13.9 **	0.93	2.8	0.11
Manure (M)	2.84	0.19	34.99**	1.36	3.54	1.47	1.64	3.18	1.8
Depth (D)	44.23**	86**	0.65	47.7**	6.43*	28.2**	8.35**	7.38*	18.92**
T X M	0.92	0.38	17.16**	3.5	3.82*	0.99	1.27	1.71	5.56*
T X D	0.01	0.37	5.69*	0.00	0.62	2.68	0.44	2.09	0.09
M X D	3.77*	2.22	0.61	1.56	0.06	1.59	0.66	0.24	0.98
T X M X D	0.26	1.13	6.68**	3.29	0.18	0.75	0.58	1.13	0.97

** Significant at P < 0.01.

*: Significant at P < 0.05.

N_m, total amount of N mineralized and N_e, N mineralized over 10 days. C_m, total amount of C mineralized and C_e, C mineralized over 10 days. OM, organic matter C_s, soluble carbon.




Figure 1. Effect of tillage and liquid manure on cumulative N mineralized over 270 d incubation.




Figure 2. Effect of tillage and liquid manure on cumulative C mineralized over 270 d incubation.

Table 3. Linear correlation coefficient of carbon and nitrogen mineralization parameters for meadow soil over 270 days incubation.

Soil characteristics								
	N_m	N_0	K_n	N_e	C_m	C_0	C_e	K_c
N_m	-	0.058	-0.399*	0.494**	0.360	-0.250	-0.25	-0.294
N_0		-	-0.337	-0.180	-0.332	-0.204	-0.284	-0.063
K_n			-	0.321	-0.245	-0.205	0.459**	0.712**
N_e				-	0.382	0.006	0.409*	0.156
C_m					-	0.398*	0.417*	-0.473*
C_0						-	-0.233	-0.57**
C_e							-	0.716**
K_c								-

N_0 , N mineralization potential; K_n , rate of N mineralization; N_m , total amount of N mineralized and N_e , N mineralized over 10 days.

C_0 , C mineralization potential; K_c , rate of C mineralization; C_m , total amount of C mineralized and C_e , C mineralized over 10 days.

*, *: Significant at $p \leq 0.05$, and $p \leq 0.01$, respectively.

Relationships between calculated and measured C and N mineralization parameters

The linear correlation between measured and calculated mineralization parameters are given in Table 3. The total amount of N mineralized (N_m) was correlated with rate constant K_n ($P < 0.05$) and with the N_e fraction ($P < 0.01$). The total amount of C mineralized (C_m) was correlated with C_0 ($P < 0.05$), and with the initial mineralizable C (C_e) ($P < 0.05$). The C_m was inversely related to the rate constant K_c of the Ellert equation ($P < 0.01$). The potential C mineralization (C_0) was also inversely related to K_c ($P < 0.05$). The initial mineralizable C (C_e) was closely correlated to K_c of the Ellert equation ($P < 0.01$).

DISCUSSION

In this work no difference was observed between soil physical properties measured for a meadow soil following different tillage and manure treatments. Sommerfeldt and Chang (1985) observed that increasing rates of manure on irrigated land tended to decrease the amount of soil aggregates < 1 mm and increase those > 1 mm of soil. They reported that tillage treatments affected the aggregate size distribution at a depth of 0 – 15 cm. The Mean Weight Diameter (MWD), it was 24% and 15% higher in the no-till soil than in the tilled soil at 0 – 15 cm and 15 – 30 cm depth respectively. Our results

corroborate the results of Karlen et al. (1994) who obtained average stability values (MWD) of 46 and 96% for long-term no-till and plow treatments respectively.

Phosphorus is relatively immobile in the soil and so tends to remain near the site of application unless mixed or incorporated to soil (ref). Under zero-tillage (ZT) and conventional tillage (CT) available P was concentrated in the surface 15 cm of soil. However, as previously reported (Campbell et al., 1986) the available P level increased significantly following manure application.

Grant and Bailey (1994) found that the concentration of K was generally higher in the surface soil under zero-tillage, presumably because of the lack of mixing normally attributed to plowing. Tillage did not have any effect on soil Ca and Mg levels. However, as previously observed by N'dayegamiye (1990) the addition of manure significantly increased soil Ca and Mg contents.

Tillage practices did not influence the soil pH as observed by Doran (1987). Our results indicated, however, that there was significant effect of manure application on soil pH ($p \leq 0.01$). The application of 50 t ha⁻¹ of manure increased the soil pH values as observed by Antoun et al. (1985).

This is in agreement with the results of Blevins et al. (1983) who found in the 0 – 5 cm surface layer that organic C and N were approximately twice as high with no-tillage as with conventional tillage. Grant and Lafond (1994) also observed that total carbon content was higher in the surface soil of the zero tillage treatment compared with the conventional tillage system. These results

indicate that tillage influences the organic C pools. In no-tilled soil, C accumulation can be attributed to a decrease in the rate of organic matter oxidation resulting from higher soil moisture and lower temperature (Tracy et al., 1990). On the other hand, C loss rate from soil is greatly accelerated by conventional tillage as indicated by Doran (1980a).

Richter et al. (1990) observed that annual tillage changed dominant plant species from grasses to annual herbs and consequently altered the distribution of carbon in root biomass. In annually tilled plots, herb-dominated root biomass averaged about 232 g C/m² compared with about 753 g C/m² in no-tilled plots. As previously observed by N'dayegamiye and Coté (1989), manure application did not affect soil total N content because addition of liquid manure increased significantly the total amount of nitrogen mineralization quality.

Cumulative N mineralization after 270 days (Nm) was higher in plowed treatments than in no-tilled treatments. Our data confirm those of Carter and Rennie (1984) and El-Haris et al. (1983) who found that cumulative N mineralization was higher under conventional than zero tillage system.

We observed that the initial N mineralization (Ne) was higher in the 15 - 30 cm plowed soil (51%) than in no-tilled soil. The observed rates of Ne are in the same range as those obtained from comparable meadow soils from Québec (Simard and N'dayegamiye, 1993).

Other soil physical and biotic properties can influence Kn (Cassman and Munns, 1980; Macdonald et al., 1995; Simard and N'dayegamiye, 1993). Higher Kn values and lower N₀ values in the no-till treatments compared with those calculated for the plowed soils suggest that the mineralizable substrate concentration was lower but decomposed more rapidly under no-till condition.

Under no-till, the ratios of organic matter/soluble C (OM/Cs) are in general higher than in plowed soils. This indicated that N₀ and Kn were quite different in the two tillage systems. An inverse relationship between N₀, Nm and Kn values was evident, suggesting that a major part of the microbial biomass is easily mineralizable (Torben and Rosswal, 1987).

Our results with liquid manure are not comparable to those of Campbell et al. (1986) who observed that mineralizable N increased significantly by the addition of barnyard manure. As observed by Lindemann and Cardenas (1984) the addition of 50 and 100 t ha⁻¹ liquid manure increased the total amount of N mineralized respectively by 34 and 41% in the 0 - 15 cm soil. We observed that total amount N mineralization decreased with increasing manure application in 15 - 30 cm depth in plowed treatments.

In this study, Kn values decreased with increasing rates of manure application. Our results are different from the observations of Boyle and Paul (1989), who found that N and C mineralization rates (Kn, Kc) increased with sludge

application rate. The average kn value was slightly greater for subsoil than surface soil in plowed treatments as previously calculated by Campbell et al. (1981).

As compared with the value for untreated soil, N₀ values were higher with 50 t ha⁻¹ (32.45%) and 100 t ha⁻¹ (42.78%) manure application in the surface plowed soil. Also N₀ values increased with 50 and 100 t ha⁻¹ manure application in no-till soil at all depths. Our results agree with those of N'dayegamiye and Coté (1989) who also observed that high liquid manure application rates increased N₀.

Tillage also influenced strongly the amount of soluble C. The soluble C (Cs) has been suggested as an index of organic matter decomposition and humification (Hu et al., 1972). In this study, Cs was higher in plowed treatments than in no-till treatments. Cumulative net mineralized C showed a curvilinear relationship with time. These results suggested that CO₂-C production in all samples decreased during incubation.

It is interesting to note that in the absence of manure application, the Cs values obtained from plowed soils are negatively correlated with the Cm values ($r = -1$; for Cs, 10.22 and 14.7 mg kg⁻¹ and Cm, 314.54 and 249.62 mg kg⁻¹). Under no-till condition Cs values are positively correlated with the Cm values ($r = 1$ for Cs, 8.73 and 7.64 mg kg⁻¹ and Cm, 265.55 and 241.91 mg kg⁻¹). This indicated that the index developed by Hu et al. (1972) would apply mainly for forest soils and for soil maintained under no-till conditions. In manured soils calculated r values are not statistically significant but in tilled soils Cs values seem to be positively correlated to the Cm values and the contrary is observed under no-till conditions. More work is required to establish the exact relationship between Cs and Cm values in soils submitted to different tillage and manure applications.

The cumulative model of Ellert and Bettany (1988) was used to describe the effect of tillage on the level of the potential mineralizable C (C₀), and to calculate the first order rate constant for mineralizable C (kc). Tillage showed no significant effect on C mineralization potential (C₀). The Kc values were higher in no-till soil than plowed soil in 0 - 15 cm depth but were higher in plowed soil than in no-till soil in 15 - 30 cm depth. Higher Kc values were obtained by Franzluebbers et al. (1995) in wheat and soybean crops under conventional tillage (0.33 day⁻¹) and no-tillage (0.34 day⁻¹).

Results suggested that there was no effect of tillage on total mineralized C (C_m), however depth significantly affected C_m values. In fact under both tillage and zero-tillage systems, greater C_m were observed in surface (0 - 15 cm) soils compared with deeper soil (15 - 30 cm). Carter and Rennie (1982) also observed that potential net mineralizable C and N were significantly greater in surface soil under zero-tillage in comparison to conventional tillage. But the reverse situation was observed at the lower depth.

Easily mineralizable C (mineralizable C after 10 days of incubation, C_e) was 51.61% higher in no-till soil than in plowed treatments in 0 - 15 cm. However, C_e was 70.61% higher in plowed treatments than in no-till soil in 15 - 30 cm depth. These results reflected the different availability of easily decomposable substrates between no-till and plowed soils. These results were similar to those of Franzluebbers et al. (1994; 1995) who observed that carbon mineralization was higher in conventional tillage for the 0 – 20 cm depth than in no-till soil.

Conclusion

The results of this study showed significant effect of tillage on soil P, K, Zn, soil C/N, soluble C and cumulative N mineralized. Also cumulative N mineralization was significantly affected by tillage and very closely related to levels of soil chemical properties. There were large differences in N mineralization potentials (N_0) in plowed treatments and no-till treatments. Differences in rate constant (Kn) increased in no-till soil as compared to plowed soil. Results obtained in our study indicated that C mineralized (C_m) and C mineralization potential (C_0) were not affected by tillage. The data showed that soil structural stability was not affected by tillage and manure application.

Generally, there were significant effect of liquid manure on soil pH, P, Ca, Mg, Zn and soluble C. The data showed that cumulative N mineralization (N_m) and N mineralization potentials (N_0) were greater at 0 - 15 cm depth in soils receiving manure. However, cumulative N mineralized decreased with increasing manure rates within 15 - 30 cm depth. The Kn and Kc values decreased with manure application. Potentially mineralizable C and N (C_0 , N_0) were higher in the surface (0 – 15 cm) than in the deeper (15 – 30 cm) layers.

These results indicated that tillage and liquid manure application have significantly influenced N and C mineralization parameters and some soil chemical properties. The results indicated the presence of close relationships between N and C mineralization parameters and some soil chemical properties and suggested that tillage and manure application influence to some extent the soil characteristics. Furthermore, both tillage and manure influence soil organic matter dynamics and quality.

ACKNOWLEDGMENTS

This work was supported by grants of the Ministère de l'Éducation du Québec and Service des sols, Ministère de l'Agriculture, des Pêcheries et de l'Alimentation du Québec. The authors wish to thank M. R. Laverdière and A. N'dayegamiye for reviewing the early drafts of the manuscript.

REFERENCES

- Anderson JPE (1982). Soil respiration. In A.L. Page ed. Methods of soil analysis. Part 2. 2nd Agronomy Monogr. 9, ed. Am. Soc. Agron. Madison, WI 831-871.
- Angers DA, Mehuys GR (1993). Aggregate stability to water. In M.R. Carter (ed) soil sampling and methods of analysis. Can. Soc. Soil Sci. Lewis Publ. 61: 651-658.
- Antoun H, Visser SA, Cescas MP, Joyal P (1985). Effect of liquid hog slurry manure application rates on silage corn yield and nutrient uptake. Can. J. Plant. Sci. 65: 63-70.
- Bennett OL, Stanford G, Mathias EL, Lundberg PE (1975). Nitrogen conservation under corn planted in quack-grass sod. J. Environ. Qual. 4: 107-110.
- Blevins RL, Thomas GW, Smith MS, Frye WW, Cornelius PL (1983). Changes in soil properties after 10 years continuous non-tilled and conventionally tilled corn. Soil tillage Res. 3: 135-146.
- Boyle M, Paul EA (1989). Carbon and nitrogen mineralization kinetics in soil previously amended with sewage sludge. Soil. Sci. Soc. Am. J. 53: 99-103.
- Campbell CA, Schnitzer M, Stewart JWB, Blederbeck VO, Selles F (1986). Effect of manure and P fertilizer on properties of a Black Chernozem in Southern Saskatchewan. Can. J. Soil. Sci. 66: 601-613.
- Campbell CA, Myers RJK, Weier K (1981). Potentially mineralizable nitrogen, decomposition rates and their relationship to temperature for five Queensland Soils. Aus. J. Soil. Res. 19: 323-332.
- Carter MR, Rennie DA (1984). Nitrogen transformation under zero and shallow tillage. Soil. Sci. Soc. Am. J. 48: 1077-1081.
- Carter MR, Rennie DA (1982). Changes in soil quality under zero tillage farming systems: Distribution of microbial biomass and mineralizable C and N potentials. Can. J. Soil. Sci. 62: 587-597.
- Cassman KG, aMunns DN (1980). Nitrogen mineralization as affected by soil moisture, temperature and depth. Soil. Sci. Soc. Am. J. 44: 1233-1237.
- Doran JW (1980a). Microbial changes associated with residue management with reduced tillage. Soil. Sci. Soc. Am. J. 44: 518-524.
- Doran JW (1980b). Soil microbial and biochemical changes associated with reduced tillage. Soil. Sci. Soc. Am. J. 44: 765-771.
- Doran JW (1987). Microbial biomass and mineralizable nitrogen distributions in no-tillage and plowed soils. Biol. Fertil. Soils 5: 68-75.
- Dormaar JF, Johnston A, Smoliak S (1984). Seasonal changes in carbon content and dehydrogenase, phosphatase and urease activities in mixed prairie and fescue grassland Ah horizons. J. Range Manage. 37: 31-35.
- Ellert BH, Bettany JR (1988). Comparison of kinetic models for describing net sulfur and nitrogen mineralization. Soil. Sci. Soc. Am. J. 52: 1692-1702.
- Ellert BH, Bettany JR (1992). Temperature dependence of net nitrogen and sulphur mineralization. Soil. Sci. Soc. Am. J. 56: 1133-1141.
- El-Haris MK, Cochran VL, Elliot LF, Bezdicke DF (1983). Effect of tillage, cropping and fertilizer management on soil nitrogen mineralization potential. Soil. Sci. Soc. Am. J. 47: 1157-1161.
- Flowers H, Arnold PW (1983). Immobilization and mineralization of nitrogen in soils incubated with pig slurry or ammonium sulphate. Soil Biol. Biochem. 15: 329-335.
- Franzluebbers AJ, Hons FM, Zuberer DA (1995). Tillage and crop effect on seasonal soil carbon and nitrogen dynamics. Soil. Sci. Soc. Am. J. 59: 1618-1624.
- Franzluebbers AJ, Hons FM, Zuberer DA (1994). Long-term changes in soil carbon and nitrogen pools in wheat management systems. Soil. Sci. Soc. Am. J. 58: 1639-1645.
- Grant CA, Bailey LD (1994). The effect of tillage and KCl addition on pH, conductance, NO₃-N, P, K and Cl distribution in the soil profile. Can. J. Soil Sci. 74: 307-314.
- Grant CA, Lafond GP (1994). The effects of tillage systems and crop rotations on soil chemical properties of a Black Chernozemic soil. Can. J. Soil. Sci. 74: 301-306.
- Hu I, Youngberg Ct, Gilmour CM (1972). Readily oxidizable carbon: An

- index of decomposition and humification of forest litter. *Soil. Sci. Soc. Am. J.* 36: 959-961.
- Karlen DL, Wollenhaupt NC, Erbach DC, Berry EC, Swan JB, Eash NS, Jordahl JL (1994). Long-term tillage effects on soil quality. *Soil Tillage Res.* 32: 313-327.
- Lindemann WC, Cardenas M (1984). Nitrogen mineralization potential and nitrogen transformation of sludge amended soil. *Soil. Sci. Am. J.* 48: 1072-1077.
- Macdonald NW, Zak DR, Pregitzer K (1995). Temperature effects on kinetics of microbial respiration and net nitrogen and sulfur mineralization. *Soil. Sci. Soc. Am. J.* 59: 233-240.
- McGill WB, Cannon KR, Robertson JA, Cook FD (1986). Dynamics of soil microbial biomass and water soluble organic C in Breton L after 50 years of cropping to two rotations. *Can. J. Soil. Sci.* 66: 1-19.
- Mehlich A (1984). Mehlich-3 soil test extractant: a modification of Mehlich-2 extractant. *Commun. Soil. Sci. Plant. Anal.* 15: 1409-1416.
- N'dayegamiye A, Coté D (1989). Effect of long-term pig slurry and solid cattle manure application on soil chemical and biological properties. *Can. J. Soil Sci.* 69: 39-47.
- N'dayegamiye A (1990). Effets à long terme d'apports de fumier solide de bovins sur l'évolution des caractéristiques chimiques du sol et de la production de maïs -ensilage. *Can. J. Plant. Sci.* 70: 767-775.
- Nelson DW, Sommers LE (1982). Total carbon, organic carbon and organic matter. Pages 539-579. In A.L. Page ed. *Methods of soil analysis Part 2. 2nd Agronomy Monogr.* 9, ed Am. Soc. Agron. Madison, WI.
- Richter DD, Babbar LI, Huston MA, Jaeger M (1990). Effects of annual tillage on organic carbon in a fine-textured Udalf: The importance of root dynamics to soil carbon storage. *Soil Sci.* 149: 78-83.
- SAS Institute (2004). *SAS User's guide: Statistics.* SAS Institute Inc. Cary, North Carolina.
- Simard RR, N'dayegamiye A (1993). Nitrogen mineralization potential of meadow soils. *Can. J. Soil Sci.* 73: 27-38.
- Sommerfeldt TG, Chang C (1985). Changes in soil properties under annual applications of feedlot manure and different tillage practices. *Soil Sci. Soc. Am. J.* 49: 983-987.
- Spratt ED, McCurdy EV (1966). The effect of various long-term soil fertility treatments on the phosphorus status of a clay Chernozem. *Can. J. Soil Sci.* 46: 29-36.
- Stanford G, Smith SJ (1972). Nitrogen mineralization potentials of soils. *Soil. Sci. Soc. Am. J.* 36: 465-472.
- Sweeten JM, Mathers AC (1985). Improving soils with livestock manure. *J. Soil. Water Conserv.* 40: 206-210.
- Torben AB, Rosswal T (1987). Seasonal variation of potentially mineralizable nitrogen in four cropping systems. *Soil. Sci. Soc. Am. J.* 51: 1508-1514.
- Tracy PW, Westfall DG, Elliot ET, Peterson GA, Cole CV (1990). Carbon, nitrogen, phosphorus and sulfur mineralization in plow and no-till cultivation. *Soil. Sci. Soc. Am. J.* 54: 457-461.
- Wakley A, Black CA (1934). An examination of the degtjareff method for determining soil organic matter and a proposal modification of the chromic acid titration method. *Soil. Sci.* 37: 29-38.