

Full Length Research Paper

# Fatty acid, amino acid, mineral and antioxidant contents of acha (*Digitaria exilis*) grown on the Jos Plateau, Nigeria

Robert H. Glew<sup>1</sup>, Emmanuel P. Laabes<sup>2</sup>, Jack M. Presley<sup>3</sup>, John Schulze<sup>3</sup>, Ronnee Andrews<sup>4</sup>, Yuan-Chen Wang<sup>5</sup>, Yu-Chen Chang<sup>5</sup> and Lu-Te Chuang<sup>5\*</sup>

<sup>1</sup>Department of Biochemistry and Molecular Biology, School of Medicine, University of New Mexico, Albuquerque, New Mexico, USA.

<sup>2</sup>Department of Family Medicine, Jos University Teaching Hospital, Jos, Nigeria.

<sup>3</sup>Genome Center Proteomics Core, Molecular Structure Facility, University of California, Davis, California, USA.

<sup>4</sup>National Institute of Occupational Health and Safety, Ohio, USA.

<sup>5</sup>Department of Biotechnology, Yuanpei University, Hsinchu, Taiwan.

Accepted 4 December, 2012

***Digitaria exilis* (Kippist) Stapf (also known as acha, hungry rice) has been cultivated for millennia in the dry savannahs of West Africa, but much remains to be learned about its nutritional properties. Acha was collected in four villages in Northern Nigeria and analyzed for fatty acids, minerals, amino acids and antioxidant content. Fatty acids accounted for 1.91% of the dry weight, with 47.4% linoleic acid and 30.5% oleic acid. The content of the essential minerals, copper, magnesium, molybdenum, zinc and calcium averaged 4.88, 1060, 0.23, 23.0 and 172 µg/g, respectively. The protein content was 6.53% and the essential amino acid pattern, except for lysine, compared favorably to a World Health Organization (WHO) reference protein. The total polyphenolic content of methanolic extracts of acha matched that of common cereals (for example, maize, rice, wheat) and the extracts contained substantial amounts of free-radical scavenging substances. Thus, acha is a source of many nutrients critical to human health.**

**Key words:** Acha, fatty acids, minerals, amino acids, polyphenols.

## INTRODUCTION

Cereals are an important source of many nutrients for populations in all regions of the world, but especially for people living in underdeveloped countries where the lack of protein-rich foods and economic constraints compel them to rely upon cereals such as maize, sorghum, millet or rice as the staple of their subsistence diets (Burkill, 1994; National Research Council, 1996; Protabase Record available at [http://database.prota.org/dbtw-wpd/exec/dbtwpub.dll?AC+QBE\\_QUERY&BU=http://atab](http://database.prota.org/dbtw-wpd/exec/dbtwpub.dll?AC+QBE_QUERY&BU=http://atab)

ase; Coda et al., 2010).

Acha (*Digitaria exilis* (Kippis) Stapf), also called pene, fonio, petit mil, fundi and hungry rice in different regions of Africa, is a member of the cereal family that includes maize and millet (Burkill, 1994; National Research Council, 1996; Jideani and Jideani, 2011; Philip and Itodo, 2006, 2012). The plant is grown from Cape Verde to Lake Chad and in other regions of sub-Saharan Africa where it provides dry-savannah populations with a cereal staple or major dietary component that compares favorably with rice, sorghum, maize and millet in terms of its content of protein, crude fat, carbohydrate and essential minerals (Leung et al., 1968; Burkill, 1994; National Research Council, 1996; Irving and Jideani, 1997; Jideani, 1999). Acha (funde) was imported into the New

\*Corresponding author. E-mail: ltchuang@mail.ypu.edu.tw. Tel: (+886) 3-5381183 ext. 8151. Fax: (+886) 3-6102312.

World from West Africa in the 15th Century and is cultivated and consumed by populations in the Dominican Republic (Dieve, 1974).

Acha is often referred to as “hungry rice” by the indigenous people of West Africa who consume this grain; however, this is a misleading term, implying it is a ‘famine food’ consumed only during times of food scarcity. In fact, acha is prized for its taste and plays important economic and cultural roles in West Africa (National Research Council, 1996). Acha is of considerable importance in Nigeria where it is commonly eaten, often in preference to other cereals, as many as three times a day as a porridge, couscous or non-alcoholic beverage (National Research Council, 1996; Jideani, 1999). It is also valued as a weaning food because of its low bulk and high caloric density. Acha requires minimal processing and can be cooked quickly. Other attributes of acha are that it grows even where rainfall and soil fertility are poor (Jideani, 1999) and can be stored in closed containers for many years without need of preservatives. Furthermore, contrary to the misconception mentioned above that some regard acha as a food only for poor populations, in Nigeria, it is widely regarded as a prestige food.

Although the literature contains reports of the content of certain nutrients in acha (Leung et al., 1968; Dieve, 1974; Temple and Bassa, 1991; Barikmo et al., 2004), it should be kept in mind that nutrient data for a particular plant food from one region of Africa may not necessarily be representative for the same plant grown in some other region where the climate, soil conditions and farming practices may be different.

Despite its popularity as a food item for millions in Africa, and the availability of informative reports of the nutrient composition of acha (Jideani et al., 1994; Philip and Itodo, 2006; Fogny-Fanou et al., 2009; Jideani and Jideani, 2011), there remains a need for additional information regarding its content of fatty acids, minerals and antioxidants. Furthermore, the nutrient content of a particular plant food can vary considerably between countries and even regions within a country due to variations in temperature, rainfall, fertilizer use and the nutrient content of the soil (Greenfield and Southgate, 1992). Chukwu and Abdul-Kadir (2008) recently conducted a study of the proximate chemical composition of acha in which they determined the content of 18 amino acids and 8 essential minerals in a sample of acha purchased in the Central Market in Minna, Nigeria. These investigators found that acha contained more of methionine and certain essential minerals, and trace elements (calcium, magnesium, iron and copper) than most cereals.

The main purpose of the present study was to replicate and extend the study by Chukwu and Abdul-Kadir (2008) by determining the fatty acid composition, amino acid content and the amounts of 13 minerals and trace elements in some specimens of acha purchased in markets in four different villages in Plateau State, Nigeria. In addition, we estimated the quantity of phenolic com-

pounds in methanolic extracts of acha and the capacity of these extracts to scavenge free radicals.

## MATERIALS AND METHODS

### Collection of acha

Acha was purchased in 2009 from retailers in open-air markets located in four villages (Rantya, Bokkos, Ganawuri and Hoss) in Plateau State in north-central Nigeria. The approximate distances of these villages from the center of the city of Jos range from 10 km (Rantya) to 150 km (Bokkos). Botanical documentation of the identity of the four specimens of acha was provided by Dr. Timothy K. Lowrey, Curator of the University of New Mexico Herbarium Museum of Southwestern Biology, Albuquerque, New Mexico, USA. The grains were hulled and milled locally and then sun-dried and ground to powder with the aid of a mortar and pestle. Immediately prior to analyzing the four acha specimens for their content of fatty acids, minerals and trace elements, amino acids and antioxidants, they were dried to a constant weight as specified below. Each of the four acha specimens was analyzed in triplicate and results reported as mean  $\pm$  standard deviation. All analyses were performed in accredited laboratories in Taiwan and the United States.

### Lipid extraction and fatty acid analysis

Prior to lipid extraction, powdered acha was vacuum-dried for 12 h using an Eyela centrifugal evaporator CVE-1000 (Tokyo, Japan). Total lipids were extracted using the method of Folch and coworkers (1975), with minor modifications (Glew et al., 2009). The extracted lipid was then reconstituted to 0.5 mL with chloroform. To prepare fatty acid methyl esters, a 0.1 mL aliquot of the chloroform solution of the total lipid fraction was evaporated under a stream of nitrogen, and then reacted with 14% (w/v) boron trifluoride methanol complex (BF<sub>3</sub>) (Morrison and Smith, 1964) for 20 min at 95°C. The fatty acid methyl esters were extracted with 1 mL of hexane. The profile of the fatty acids in the hexane extract (1  $\mu$ L) was determined by gas chromatography (GC) as previously described (Glew et al., 2009).

### Mineral analysis

Samples (0.2 g) of vacuum desiccator-dried and powdered acha were weighed into 125 mL Phillips beaker, covered with watch glass and digested at 150°C for 30 min with 10 mL of concentrated nitric acid and 0.6 mL of 60% perchloric acid. The watch-glass cover was then removed and the samples were brought to near dryness at 120°C. The digested samples were cooled to room temperature and brought to 10 mL with 4% nitric acid/1% perchloric acid. The digested samples were analyzed for metal content by inductively coupled plasma optical emission spectrometry (ICP-OES) as described earlier (Fernandez et al., 2003).

### Amino acid analysis

Twenty milligrams of desiccator-dried acha were hydrolyzed in 6 N HCl containing 1.0% (w/v) phenol at 110°C for 24 h *in vacuo*, and the resultant amino acids were separated and quantified using a Hitachi Amino Acid Analyzer L8900 (Tokyo, Japan) according to the manufacturer's instructions (Dionex Corporation, 2001) and previously described methods (Ozols, 1990; Clark et al., 1999; Jadnick et al., 1999). For the determination of methionine and cysteine, samples were oxidized with performic acid (Hirs, 1967) prior to acid hydrolysis. The coefficient of variation of the method ranged from 0.6 to 11% for the amino acids reported. Tryptophan

**Table 1.** Mass content (mg/g dry weight of sample) of fatty acids in four specimens of acha.

Fatty acid	Rantya	Bokkos	Ganawuri	Hoss	Mean
C16:0	2.82 (0.23) <sup>a</sup>	2.53 (0.39)	2.36 (0.26)	3.81 (0.08)	2.88 (0.24)
C16:1 n-7	0.04 (0.01)	0.05 (0.01)	0.04 (0.01)	0.06 (0.01)	0.05 (0.01)
C18:0	0.59 (0.05)	0.52 (0.13)	0.48 (0.05)	0.73 (0.01)	0.58 (0.06)
C18:1 n-9	5.69 (0.47)	5.37 (0.61)	4.45 (0.53)	7.87 (0.18)	5.85 (0.45)
C18:1 n-7	0.17 (0.00)	0.20 (0.06)	0.13 (0.03)	0.25 (0.01)	0.19 (0.02)
C18:2 n-6	8.78 (0.72)	8.22 (1.07)	7.14 (0.87)	12.2 (0.27)	9.08 (0.47)
C18:3 n-3	0.24 (0.02)	0.22 (0.03)	0.15 (0.02)	0.28 (0.01)	0.22 (0.02)
C20:0	0.15 (0.01)	0.16 (0.02)	0.13 (0.01)	0.21 (0.00)	0.16 (0.01)
C20:1	0.07 (0.00)	0.08 (0.02)	0.06 (0.01)	0.10 (0.01)	0.08 (0.01)
C22:0	0.08 (0.01)	0.06 (0.01)	0.09 (0.04)	0.11 (0.01)	0.06 (0.02)
Total	18.6	17.4	15.0	25.6	19.1 (3.1)

<sup>a</sup>The number in parentheses indicates one standard deviation.

was not measured because hydrolysates of acha were incompatible with the particular chromatography column used to resolve and quantify amino acids.

#### Determination of total phenolic compounds

A one gram sample of acha was extracted three successive times for 24 h at 25°C with 20 ml of methanol in a shaker incubator at 125 rpm and then centrifuged at 1,000 × *g* for 10 min. The supernatants were combined, filtered through type 5A filter paper (Advantec, Tokyo, Japan) and evaporated using a vacuum concentrator. Finally, the acha extract was lyophilized and the dry weight of the extract determined. The total phenolic acid content of the methanol extract obtained above was estimated using the Folin-Ciocalteu colorimetric assay (Singleton et al., 1999). Gallic acid was used as the standard (0 to 40 mg/L), and results were expressed as mg of gallic acid equivalent (GAE) per gram dry weight of original acha specimen. These measurements were performed in triplicate.

#### The 1,1-diphenyl-2-picrylhydrazyl free-radical scavenging assay

The ability of the methanol extracts of acha to neutralize the free radical 1,1-diphenyl-2-picrylhydrazyl (DPPH) (Sigma Chemical Co.) was determined using the method of Yu et al. (2002). The lyophilized methanol extract was dissolved in distilled water and diluted to the appropriate concentration. One hundred microliters of the diluted solutions were aliquoted into the wells of a 96-well plate and mixed with 0.1 mL of 0.2 mM DPPH reagent. Distilled water served as control, and DPPH was replaced with distilled water to provide the blank. The plate was left to stand in the dark at 25°C for 30 min after which the absorbance was measured at 517 nm with the aid of an Anthos Zenyth 3100 Enzyme-linked immunosorbent assay (ELISA) reader (Anthos Labtec Instruments, Salzburg, Austria). The DPPH scavenging activity was calculated as: DPPH scavenging (%) = [1 – (sample absorbance / control absorbance)] × 100%. The EC<sub>50</sub> is the concentration (mg/mL) of the methanolic extract of acha required to scavenge 50% of the DPPH radicals in the assay.

#### Statistical analyses

Data were analyzed by analysis of variance (ANOVA) and Duncan's multiple range test using Number Cruncher statistical software

(version 6, 2004, Kaysville, UT, USA). A *p* value less than 0.05 was considered statistically significant.

## RESULTS AND DISCUSSION

### Fatty acid composition and content

Since the fatty acid percentages for the acha specimens gathered at the four different locations were statistically indistinguishable, the four data sets were averaged (Table 1). Fatty acid accounted for 15.0 to 25.6 mg/g of the total dry weight of the four acha specimens and averaged 1.91% of the dry weight (Table 2). The acha from Hoss contained the highest amount of total fatty acid (25.6 mg/g dry weight) and that from Ganawuri contained the lowest amount (15.0 mg/g dry weight). Saturated fatty acids (mainly palmitic acid and stearic acid) represented, on average, only 18.1% of the total fatty acids. Noteworthy was the finding that, on the basis of percentage or absolute amount, the essential omega-6 fatty acid linoleic acid was the most abundant fatty acid in all four of the acha specimens. The acha contained very little  $\alpha$ -linolenic acid. Furthermore, the average omega-6/omega-3 ratio for the four specimens was 41/1.

Acha, like most whitened cereals, does not contain very much fatty acid (Table 2) (mean, 1.91% of dry weight). The daily recommended intakes of linoleic acid and  $\alpha$ -linolenic for an adult are 13 to 17 g and 1.1 to 1.6 g, respectively (World Health Organization, 2007). Fifty grams dry weight of acha would provide only about 0.4 g of linoleic acid and 0.01 g of  $\alpha$ -linolenic acid. The quantity of  $\alpha$ -linolenic acid in acha represents 2% or less of the daily requirement for an adult (World Health Organization, 1985). Furthermore, the fatty acid profile of acha is disadvantageous in respect of human nutrition because the ratio of linoleic acid/ $\alpha$ -linolenic acid is relatively high at 41/1.

To put this ratio into perspective, consider that the linoleic/ $\alpha$ -linolenic acid ratio in many western diets is in the range of 10/1 to 20/1. A linoleic acid/ $\alpha$ -linolenic acid

**Table 2.** Mass percentage of fatty acids in four specimens of acha.

Fatty acid	Rantya	Bokkos	Ganawuri	Hoss	Mean
C16:0	15.1 (0.06) <sup>a</sup>	14.5 (0.66)	15.7 (0.24)	14.9 (0.05)	15.1 (0.25)
C16:1 n-7	0.23 (0.03)	0.27 (0.02)	0.24 (0.02)	0.25 (0.03)	0.25 (0.03)
C18:0	3.15 (0.02)	2.95 (0.36)	3.17 (0.08)	2.84 (0.02)	3.03 (0.12)
C18:1 n-9	30.6 (0.09)	30.9 (0.69)	29.6 (0.11)	30.8 (0.10)	30.5 (0.25)
C18:1 n-7	0.94 (0.08)	1.17 (0.40)	0.88 (0.06)	0.96 (0.04)	0.99 (0.15)
C18:2 n-6	47.2 (0.07)	47.2 (0.83)	47.5 (0.05)	47.6 (0.03)	47.4 (0.25)
C18:3 n-3	1.30 (0.01)	1.25 (0.07)	0.99 (0.02)	1.08 (0.03)	1.16 (0.03)
C20:0	0.79 (0.02)	0.92 (0.01)	0.88 (0.05)	0.81 (0.02)	0.85 (0.02)
C20:1	0.35 (0.02)	0.44 (0.10)	0.40 (0.04)	0.38 (0.03)	0.39 (0.05)
C22:0	0.43 (0.07)	0.34 (0.03)	0.60 (0.16)	0.43 (0.02)	0.45 (0.07)

<sup>a</sup>The number in parentheses indicates one standard deviation.

ratio greater than 5/1 is widely regarded as pro-inflammatory and generally unhealthful (Simopoulos et al., 1999; Simopoulos, 2002), especially with regard to cardiovascular disease and certain cancers (Ozols, 1990; Jadnick et al., 1999). In contrast to acha, the linoleic acid/ $\alpha$ -linolenic acid ratio of black finger millet is 5.5/1 (Masum-Akond et al., 2002). To our knowledge, there are no fatty acid data in the literature for acha against which we might compare our data; however, Jideani (1999) reported that acha contains 2.5% crude fat, a value that is consistent with our finding of an average fatty acid content of 1.91% (Table 1). Others (National Research Council, 1996) have reported a fat content of 1.8% for acha.

#### Mineral and trace element content

The mean calcium content of the four acha samples was 172  $\mu\text{g/g}$  dry weight and the mean copper content was 4.88  $\mu\text{g/g}$  (Table 3). The magnesium content of the four acha samples ranged from 826  $\mu\text{g/g}$  in the acha from Ganaweri to 1520  $\mu\text{g/g}$  in Hoss. There was a very wide range (11.9 to 121  $\mu\text{g/g}$ ) in the iron content of acha. The mean manganese content was 14.8  $\mu\text{g/g}$ . All four acha specimens contained high levels of zinc (20 to 26  $\mu\text{g/g}$ ) while selenium (a component of glutathione peroxidase known to defend the body against oxidative stress related diseases) was not detected in any of the four samples of acha. As for unhealthful elements, all of the plants contained significant amount of strontium (0.70 to 1.35  $\mu\text{g/g}$ ) but lead was not detectable (data not shown). We were interested in estimating the extent to which acha grown on the Jos Plateau might contribute to satisfying the calcium and iron requirements of humans. The recommended daily intakes of calcium (Bhatia, 2008) and iron (Andrews, 1999) for an adult male are 1,000 and 10 mg, respectively. From the data in Table 3, one can estimate that 50 g dry weight of acha would provide only about 10 mg of calcium or 1% of the daily recommended

amount but 2.5 mg or 25% of an individual's daily iron need. Acha could also contribute significantly to an adult's daily needs of copper, magnesium and zinc. As for minerals and trace elements, acha obtained in Minna and that purchased in Plateau State contained similar amounts of calcium and magnesium (Chukwu and Abdulkadir, 2008); however, the acha gathered in Minna contained one-fourth less copper but 3 to 4-fold more iron than the acha from the Jos Plateau that we analyzed in the present study.

#### Amino acid composition and content

As shown in Table 4, the protein content (obtained by summing all the amino acids together) of the four acha specimens ranged from 4.65 to 8.02% (mean, 6.53%). Temple and Bassa (1991), using a direct method for estimating protein, reported that acha contains 7% crude protein. To assess the quality of the protein in the acha specimens, we compared their proportions of essential amino acid (except tryptophan) to the proportions of the same amino acids in a WHO reference protein (World Health Organization, 1985). The data in Table 5 show that acha scored above 100 for five of seven essential amino acid categories. The scores for lysine and threonine were 42 and 90%, respectively, of the corresponding amino acids in the WHO reference protein. The sulfur amino acid (that is, cysteine plus methionine) score for acha was 231, which is high relative to that for other West Africa staple cereals (rice, maize, millet and sorghum). With regard to the other essential amino acids or amino acid pairs; isoleucine, leucine, threonine, valine and phenylalanine plus tyrosine, the essential amino acid scores for acha ranged between 106 and 150 (Table 5).

Thus, in terms of protein content and its pattern of amino acids relative to the WHO standard, the acha samples from four different villages on the Jos Plateau of North-central Nigeria contained moderate amounts of protein and, except for lysine, a pattern of essential amino

**Table 3.** Amino acid content (mg/100 mg dry weight) of acha grown in Nigeria.

Amino acid	Rantya	Bokkos	Ganawuri	Hoss	Mean
Aspartic acid	0.487 (0.040) <sup>d</sup>	0.511 (0.066)	0.325 (0.033)	0.405 (0.032)	0.432
Threonine	0.250 (0.015)	0.271 (0.023)	0.192 (0.018)	0.217 (0.023)	0.233
Serine	0.419 (0.040)	0.477 (0.064)	0.239 (0.024)	0.315 (0.023)	0.369
Glutamic acid	1.78 (0.14)	1.81 (0.02)	0.95 (0.11)	1.29 (0.11)	1.46
Proline	0.420 (0.044)	0.480 (0.008)	0.248 (0.019)	0.342 (0.037)	0.373
Glycine	0.191 (0.009)	0.188 (0.008)	0.140 (0.014)	0.172 (0.013)	0.173
Alanine	0.694 (0.047)	0.751 (0.012)	0.413 (0.044)	0.525 (0.040)	0.596
Valine	0.417(0.018)	0.413 (0.030)	0.247(0.024)	0.301(0.029)	0.345
Isoleucine	0.322(0.097)	0.332 (0.060)	0.173 (0.019)	0.301(0.029)	0.282
Leucine	0.737 (0.025)	0.910 (0.019)	0.444 (0.054)	0.554 (0.054)	0.536
Tyrosine	0.200 (0.021)	0.231 (0.020)	0.124 (0.009)	0.160 (0.037)	0.179
Phenylalanine	0.463 (0.012)	0.555 (0.016)	0.263 (0.021)	0.355 (0.034)	0.409
Histidine	0.170 (0.010)	0.179 (0.014)	0.098 (0.012)	0.131 (0.010)	0.145
Lysine	0.160 (0.006)	0.149 (0.019)	0.121 (0.019)	0.159 (0.014)	0.147
Arginine	0.160 (0.037)	0.262 (0.038)	0.178 (0.034)	0.239 (0.029)	0.211
Cysteine <sup>e</sup>	0.180	0.163	0.146	0.168	0.164
Methionine <sup>e</sup>	0.384	0.342	0.342	0.378	0.362
Total	7.54 <sup>a</sup>	8.02 <sup>a</sup>	4.65 <sup>b</sup>	5.93 <sup>c</sup>	6.53

<sup>a</sup>Different from groups 3 and 4, <sup>b</sup>different from groups 1, 2 and 4, <sup>c</sup>different from groups 1, 2 and 3, <sup>d</sup>the number in parentheses is one standard deviation, <sup>e</sup>only a single determination was made.

**Table 4.** Comparison of the mineral content ( $\mu\text{g/g}$  dry weight) of four specimens of acha and black finger millet.

Mineral	Acha specimen				Mean	Black finger millet <sup>a</sup>
	Rantya	Bokkos	Ganawuri	Hoss		
Calcium	187	86	100	314	172 (105) <sup>b</sup>	4010
Chromium	n.d.	0.19	0.22	0.32	0.21 (0.09)	2.23
Copper	4.57	5.44	4.12	5.39	4.88 (0.64)	7.47
Iron	19.0	40.9	11.9	121	48.2 (50.1)	182
Potassium	2430	2200	2250	3070	2490 (401)	5410
Magnesium	1010	826	894	1520	1060 (314)	1740
Manganese	22.9	10.6	8.95	16.6	14.8 (6.30)	292
Molybdenum	0.16	0.30	<0.11 <sup>c</sup>	<0.11 <sup>c</sup>	-	0.15
Sodium	<41 <sup>c</sup>	<41 <sup>c</sup>	<41 <sup>c</sup>	<41 <sup>c</sup>	-	148
Phosphorus	2180	1950	2040	3310	2370 (634)	2760
Zinc	24	21	20	26	23 (3)	27.1

<sup>a</sup>Data from Glew et al. (2008), <sup>b</sup>the number in parentheses indicates one standard deviation, <sup>c</sup>lower limit of detection. Selenium was not detected (<4.2  $\mu\text{g/g}$  dry weight). n.d = no determination.

amino acids that compared favorably to a WHO reference protein (World Health Organization, 1985). The amounts and proportions of amino acids in the acha obtained in four different villages in Plateau State are similar to values reported by Chukwu and Adbul-kadir (2008), Temple and Bassa (1991), and others (National Research Council, 1996) for acha. As others have noted previously (World Health Organization, 1985; Chukwu and Adbul-kadir, 2008; Carbenier et al., 1960; Vodouhe et al., 2003), it was also found in this study that acha protein contains

relatively high proportions of the sulfur amino acids (methionine and cysteine), and glycine, glutamate/glutamine, proline and leucine (Table 3).

The protein in a typical serving of acha could satisfy about 10% of the daily needs of an adult for all of the essential amino acids except lysine which scored only 42% relative to the WHO standard protein (Table 5). For example, consumption of 50 g dry weight of acha (about 65 g of freshly-harvested acha) with a protein content of 7% would provide 3.5 g of protein, or a little less than

**Table 5.** Comparison of the amino acid composition of acha to black finger millet and the WHO reference protein (World Health Organization, 1985)

Amino acid	WHO ideal <sup>a</sup> (% of total)	Black finger millet <sup>b</sup> (% of total)	Acha <sup>c</sup> (% of total)	Acha × 100/ WHO ideal
Isoleucine	4.0	4.8	4.3	108
Leucine	7.0	10.4	8.2	117
Lysine	5.5	2.6	2.3	42
Methionine plus cysteine	3.5	6.2	8.1	231
Phenylalanine plus tyrosine	6.0	6.7	9.0	150
Threonine	4.0	3.9	3.6	90
Valine	5.0	6.8	5.3	106

<sup>a</sup> WHO (World Health Organization, 1985), <sup>b</sup> data from Glew et al. (2008), <sup>c</sup> averages from Table 4 of this report.

populations relying heavily on this grain should be advised to consume other food sources such as soy beans that would supplement their lysine intake.

### Polyphenol content

The mean polyphenol content of the acha specimens from the four villages was 2.0 (0.01) mg/g dry weight, which is similar to the polyphenol content of 2.2 to 3.2 mg/g reported for rice (Chotimarkon et al., 2008), 1.29 mg/g for wheat (Masum-Akond et al., 2002) and 0.8 to 1.4 mg/g for oats (Adom and Liu, 2002). The DPPH radical-scavenging activity of the methanol extracts of the four acha specimens, expressed as the EC<sub>50</sub> value, was 0.51 units/mL which is similar to values reported for several varieties of winter wheat (Kähkönen et al., 1999). The total phenolic content of methanolic acha extracts, estimated as gallic acid equivalents (GAEs), is in the range of values reported for other cereals, including rice, wheat, maize and oats (Adom and Liu, 2002; Masum-Akond et al., 2002; Chotimarkon et al., 2008). However, although the ability of methanolic extracts of our four acha specimens to neutralize free radicals was poor relative to similar extracts of green leafy vegetables, the antioxidant content of acha was similar to that of other grains which, in general, are not rich in these types of substances.

The overall result of this study was the finding that acha grown on the Jos Plateau contains useful amounts of the essential fatty acid linoleic acid (Table 2), a number of essential minerals and trace elements, including copper, iron, magnesium, molybdenum and zinc (Table 3), and nutritionally-significant amounts of good-quality protein (Tables 4 and 5). However, compared to another common cereal grown on the Jos Plateau, namely black finger millet (Glew et al., 2008), acha appears to be a poor to moderate source of  $\alpha$ -linolenic acid, calcium, chromium, manganese and lysine. It has long been recognized that acha is a poor source of lysine (Food and Agriculture Organization, 1970; Chukwu and Abdul-kadir, 2008).

The wide variation we found for the iron content of the four acha specimens could be due to contamination introduced by the tools used by the local farmers to gather and mill acha. In contrast, the percentages of individual fatty acids between the four different specimens of acha analyzed in the present study was small (less than 16%) relative to the large variations in the content of various minerals which ranged from 13 to 117% (Table 3). This observation indicates that the fatty acid composition of acha grown in different locations on the Jos Plateau was much less dependent on soil conditions and climate than the mineral content of acha. Nevertheless, it would be useful to test under controlled conditions the effects of light, temperature, rainfall and fertilizer use on the nutrient composition of acha.

In light of the relatively high incidence of malnutrition (for example, stunting, underweight, wasting) and deficiencies of zinc and other micronutrients in Nigeria and many other regions of West Africa (Carbenier et al., 1960; Thacher et al., 1999; Agyei-Frempong et al., 2001; Lutter and Rivera, 2003; Vodouhe et al., 2003; Adu-Afarwuch et al., 2008), the information in this study should serve to both underscore the significant contributions acha can make to the human diet; however, it also points out certain nutritional limitations of this cereal grain, in particular, its low content of lysine, the minerals and trace elements calcium, manganese, selenium and the essential fatty acid ( $\alpha$ -linolenic acid).

Future studies should be aimed at identifying means and growing conditions that would increase the content of these particular nutrients in acha. There is also a need for studies of the bioavailability of various nutrients in acha and on the potential thyroid toxicity of flavonoids contained in the seed (Startelet et al., 1996). Furthermore, the suggestion by Jideani (1999) that acha might be a useful component of special diets for individuals with diabetes remains to be evaluated. Finally, since the Jos University Teaching Hospital in Jos, Nigeria (Fernandez et al., 2002; Bond et al., 2005) and institutions elsewhere in West Africa (Agbon et al., 2009) have a long-standing interest in developing cereal-based weaning foods and

complimentary foods for malnourished children and adults, the information provided herein regarding the nutrient content of acha should be helpful to dieticians in Nigeria and elsewhere in the world, including the Caribbean region where acha is also grown (Leung et al., 1968), who are interested in developing the next generation of complementary foods.

## REFERENCES

- Adom KK, Liu RH (2002). Antioxidant activity of grains. *J. Agric. Food Chem.* 50:6182-6187.
- Adu-Afarwuch S, Lartey A, Brown KH, Zlotkin S, Briend A, Dewey KG (2008). Home fortification of complementary foods with micronutrient supplements is well accepted and has positive effects on iron status in Ghana. *Am. J. Clin. Nutr.* 87:929-938.
- Agbon CA, Oguntona CRB, Mayaki TF (2009). Micronutrient content of traditional complementary foods. *The Forum for Family and Consumer Issues* 14 (2) <http://ncsu.edu/ffci/publications/v14-n2-2009-summer/index-v14-n2-summer-2009.ph>
- Agyei-Frempong MT, Asare G, Owiredu WKBA, Yeboah FA (2001). Iron deficiency in rural Ghanaian children. *East Afr. Med. J.* 78:246-249.
- Andrews NC (1999). Disorders of iron metabolism. *N. Engl. J. Med.* 342:1986-1995.
- Barikmo I, Ouattara F, Oshaug A (2004). Protein, carbohydrate and fibre in cereals from Mali- How to fit the results in a food composition table and database. *J. Food Comp. Anal.* 17:291-300.
- Bhatia V (2008). Dietary calcium intake - a critical reappraisal. *Indian J. Med. Res.* 127:269-273.
- Bond B, Fernandez DR, VanderJagt DJ, Williams M, Huang Y-S, Chuang L-T, Millson M, Andrews R, Glew RH (2005). Fatty acid, amino acid and trace mineral analysis of three complementary foods from Jos, Nigeria. *J. Food Comp. Anal.* 18:675-690.
- Burkill HM (1994). *The Useful Plants of West Africa*, 2nd Edition, Volume 2. Families E-I, Royal Botanic Gardens, Kew, Richmond, United Kingdom. P 636.
- Carbenier R, Jaeger P, Busson F (1960). Study of the protein fraction of fonio, *Digitaria exilis* (Kippist) Stapf., a protein exceptionally rich in methionine. *Ann. Nutr. Aliment* 14:165-169.
- Chotimarkon C, Benjakul S, Silalai N (2008). Antioxidant components and properties of five long-grained rice bran extracts from commercial available cultivars in Thailand. *Food Chem.* 111:636-641.
- Chukwu O, Abdul-Kadir AJ (2008). Proximate chemical composition of acha (*Digitaria exilis* and *Digitaria iburu*) grains. *J. Food Technol.* 6:214-216.
- Clark AP, Jadnick P, Rocklin RD, Liu Y, Avdalovic N (1999). An integrated amperometry waveform for direct, sensitive detection of amino acids and amino sugars following anion-exchange chromatography. *Anal. Chem.* 71:2774-2781.
- Coda R, Di Cagno R, Edema MO, Nionelli L, Gobetti M (2010). Exploitation of acha (*Digitaria exilis*) and iburu (*Digitaria iburu*) flours: Chemical characterization and their use for sourdough fermentation. *Food Microbiol.* 27:1043-1050.
- Dieve CE (1974). Glosario de afronegrismos en la toponimia y el espanol hablado en Santo Domingo, Boletín del Museo del Hombre Dominicano 5. Santo Domingo, Dominican Republic.
- Dionex Corporation (2001). Determination of the amino acid content of peptides by AAA-Direct, Dionex Corporation Technical Note 50, Dionex Corporation, Sunnyvale, CA.
- Fernandez DR, VanderJagt DJ, Williams M, Huang Y-S, Chuang L-T, Millson M, Andrews R, Pastuszyn A, Glew RH (2002). Fatty acid, amino acid, and trace mineral analyses of five weaning foods from Jos, Nigeria. *Plant Foods Hum. Nutr.* 57:257-274.
- Fernandez DR, VanderJagt DJ, Millson M, Huang YS, Chuang LT, Pastuszyn A, Glew RH (2003). Fatty acid, amino acid and trace mineral composition of *Eleusine corocana* (Pwana) seeds from Northern Nigeria. *Plant Foods Hum. Nutr.* 58:1-10.
- Fogny-Fanou N, Koreissi Y, Dossa RAM, Brouwer ID (2009). Consumption of and beliefs about fonio (*Digitaria exilis*) in urban area in Mali. *Afr. J. Food Agric. Nutr. Dev.* 9:1927-1944.
- Folch J, Lees M, Stanley GH (1975). A simple method for the isolation and purification of total lipids from animal tissues. *J. Biol. Chem.* 226:497-509.
- Food and Agriculture Organization (1970). Amino-acid content of foods and biological data on proteins. *FAO Nutrition Studies No. 24*, Rome, Italy.
- Glew RS, Chuang LT, Roberts JL, Glew RH (2008). Amino acid, fatty acid, and mineral content of black finger millet (*Eleusine corocana*) cultivated on the Jos Plateau. *Food* 2:115-118.
- Glew RS, Amoako-Atta B, Ankar-Brewoo G, Presley J, Chuang LT, Millson M, Smith BR, Glew RH (2009). Non-cultivated plant foods in West Africa: Nutritional analysis of the leaves of three indigenous leafy vegetables in Ghana. *Food* 3: 39-42.
- Greenfield H, Southgate DAT (1992). *Food Composition Data-Production, Management and Use*. Elsevier Science Publishers LTD, Barking.
- Hirs CWH (1967). Performic acid oxidation. *Method Enzymol.* 11:197-199.
- Irving DW, Jideani IA (1997). Microstructure and composition of *Digitaria exilis* Stapf (acha): A potential crop. *Cereal Chem.* 74:224-228.
- Jadnick P, Clark AP, Avdalovic N, Andersen DC, Cacia J (1999). Analyzing mixtures of amino acids and carbohydrates using bi-modal integrated amperometric detection. *J. Chromatogr.* 732:193-201.
- Jideani AI, Apentwn RKO, Miller HG (1994). The effect of cooking on proteins from acha (*Digitaria exilis* Stapf.). *J. Sci. Food Agric.* 65:465-476.
- Jideani I, Jideani V (2011). Development on the cereal grains *Digitaria exilis* (acha) and *Digitaria iburu*. *J. Food Sci. Technol.* 48:251-259.
- Jideani LA (1999). Traditional and possible technological uses of *Digitaria exilis* (acha) and *Digitaria iburu* (iburu): A review. *Plant Foods Hum. Nutr.* 54:363-374.
- Kähkönen MP, Hopia AI, Vuorela HJ, Rahua JP, Pihlaga K, Kujala TS, Heinonen M (1999). Antioxidant activity of plant extracts containing phenolic compounds. *J. Agric. Food Chem.* 47:3954-3962.
- Leung W-TW, Busson F, Jardin C (1968). *Food Composition Table for Use in Africa*. U.S. Department of Health, Education and Welfare and the Food and Agriculture Organization of the United Nations (FAO), Bethesda, Maryland.
- Lutter CK, Rivera JA (2003). Nutritional status of infants and young children and characteristics of their diets. *J. Nutr.* 133:2941S-2949S.
- Masum-Akond ASMG, Khandakar L, Hossain KG, Furuta Y (2002). Total polyphenol, polyphenol oxidase, antioxidant activity and color profiles of some wheat varieties from Bangladesh. *Res. J. Agri. Biol. Sci.* 6:186-190.
- Morrison WR, Smith LM (1964). Preparation of fatty acid methyl esters and dimethylacetals from lipids with boron trifluoride methanol. *J. Lipid Res.* 5:600-608.
- National Research Council, National Academy of Sciences (1996). *Fonio (Acha)*. In: *Lost Crops of Africa, Volume I, Grains*. National Academy Press, Washington, D.C., USA. pp 59-75.
- Ozols MJ (1990). Amino acid analysis. *Methods Enzymol.* 182:587-601.
- Philip T, Itodo I (2006). Acha (*Digitaria spp.*) a "Rediscovered" Indigenous Crop of West Africa. *Agricultural Engineering International: The CIGR e-journal Invited Overview No. 23*, Vol.VIII December.
- Philip TK, Itodo IN (2012). Demographic characteristics, agricultural and technological profile of acha farmers in Nigeria. *Agric. Eng. Int.* 14:89-93.
- Protabase Record display, *Digitaria exilis* (Kippist) Stapf", available at [http://database.prota.org/dbtw-wpd/exec/dbtwpub.dll?AC+QBE\\_QUERY&BU=http://atabase](http://database.prota.org/dbtw-wpd/exec/dbtwpub.dll?AC+QBE_QUERY&BU=http://atabase) (accessed 2 May 2009).
- Simopoulos AP, Leaf A, Salem N (1999). Workshop on the essentiality of and recommended dietary intake of omega-3 fatty acids, 7-9 April, National Institutes of Health (NIH) in Bethesda, Maryland, USA.
- Simopoulos AP (2002). The importance of the ratio of omega-6/omega-3 essential fatty acids. *Biomed. Pharmacother.* 56:365-379.
- Singleton VL, Orthofer R, Lamuela-Raventos RM (1999). Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Method. Enzymol.* 299:152-178.
- Startelet H, Serghat S, Lobstein A, Ingenbleek Y, Anton R, Petitfrere E, Aguié-Aguie G, Martiny L, Haye B (1996). Flavonoids extracted from fonio millet (*Digitaria exilis*) reveal potent antithyroid properties.

- Nutrition 12:100-106.
- Temple VJ, Bassa JD (1991). Proximate chemical composition of acha (*Digitaria exilis*) grain. J. Sci. Food Agric. 56:561-563.
- Thacher T, Glew RH, Isichei C, Lawson JO, Scariano JK, Hollis BW, VanderJagt DJ (1999). Rickets in Nigerian children: Response to calcium supplementation. J. Trop. Pediatr. 45:202-207.
- Vodouhe SR, Zannou A, Achigan-Dako GE (2003). Proceedings of the 1st Workshop on Genetic Diversity of Fonio (*Digitaria exilis* Staph.) in West Africa, Guinea Conakry, 4-6 August, 1998. IPGRI, Rome, Italy.
- World Health Organization (1985). WHO/FAO Report: Energy and protein requirements. WHO technical report series No. 724, WHO Geneva.
- World Health Organization (2007). Protein and amino acid requirements in human nutrition. Report of a joint WHO/FAO/UNU Expert Consultation. United Nations University.
- Yu L, Haley S, Perret S, Harris M, Wilson J, Qian M (2002). Free radical scavenging properties of wheat extracts. J. Agric. Food Chem. 50:1619-1624.