

Review

# Tomato leafminer [(*Tuta absoluta* Meyrick) (Lepidoptera: Gelechiidae)] and its current ecofriendly management strategies: A review

Birhan Aynalem

Department of Biotechnology, Debre Markos University, Ethiopia.

Received 17 November, 2017; Accepted 13 December, 2017

Continue climatic changes in the world enhance the emergence of invasive agricultural insect pests. *Tuta absoluta* is one of the most devastating tomato leafminer and it spreads extensively in almost all parts of the world. It is a serious threat for tomato production and it results in highest damage. *T. absoluta* has fast growth rate with developmental stages of egg, larvae, pupa, and adult and it adapts to different environmental conditions. A larva is the most destructive one that consumes preferably leaves, stems and fruits of the tomato plant by hiding within mesophyll of the plant tissues. Although different environmentally hazardous pesticides are used against *T. absoluta*, none of the insecticides give full control due to the resistance development. This problem calls an urgent search for eco-friendly alternatives to control this invasive insect. Recent investigations are focused on different environmental friendly approaches that inclusively termed as IPM strategies. These are cultural practices, biopesticides, and biological using parasitoids, predators, microbial cells, microbial products, RNAi, inheritance sterility development in insect and pest resistant plant cultivar production are considered as promising alternatives to control *T. absoluta*. Therefore, this review briefly describes the current status of these methods used to design suitable and sustainable management strategies against *T. absoluta*.

**Key words:** Tomato leafminer, insecticide resistances, biopesticides, biological control.

## INTRODUCTION

Alarmingly increasing climatic changes in the world expands the opportunity for the emergence of different invasive insect pests (Robinet and Roques, 2010). This is currently a serious threat to the agricultural industry. The insect [*Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae)]

is a worldwide (Desneux et al., 2010) devastating Solanaceous plant pest. It is first discovered from Peru in 1917 and recognized as leafminer of tomato (*Solanum lycopersicum* L.) in South America in 1960 (Guedes and Picaco, 2012). The notorious manifestation of *T. absoluta*

E-mail: [berha.bat@gmail.com](mailto:berha.bat@gmail.com).

Author(s) agree that this article remain permanently open access under the terms of the [Creative Commons Attribution License 4.0 International License](https://creativecommons.org/licenses/by/4.0/)

on tomato was detected from Spain in 2006 (Urbaneja et al., 2007). This insect has alarmingly spread into Asian countries, Europeans, Mediterranean shores and recently it is also extending into African countries (Karadjova et al., 2013; Zekeya et al., 2016).

In African, *T. absoluta* was firstly reported in Algeria, Morocco, and Libya in 2008 and 2009 (Harbi et al., 2012) and continued to invade Egypt in 2010 (Moussa et al., 2013), then reached Sudan and South Sudan in 2011 (Pfeiffer et al., 2013; Brevault et al., 2014). The pest had accessed to Ethiopia in 2012 most probably from Sudan or Yemen (Goftishu et al., 2014) and Kenya in 2013 (Mohamed et al., 2015), Tanzania and Senegal in 2014 (Biondi et al., 2015; Tonnang et al., 2015), Uganda in 2015 (Tumuhaise et al., 2016), South Africa in 2016 (Visser et al., 2017). The main spread vehicle for this invasive insect species was natural dispersal means, such as wind (Gontijo et al., 2013; Sridhar et al., 2014) and human fruit and vegetable translocation processes in local and abroad markets (Karadjova et al., 2013).

The possible entry site for *T. absoluta* into Ethiopia was expected from nearby Yemen, where severe tomato leafminer infestations was recorded (Gray et al., 2013) and the border of Sudan through Humera (Retta and Berhe, 2015) in the Northern part of the country. The spread of this pest was continued into Eastern (Somali and Oromia) regions and parts of shewa, then invaded the Central Refit Valley (Fortune, 2013). The distribution of *T. absoluta* in Ethiopia was terribly occupying the main tomato farming locations. Similarly, the abundance is also high when detected from moth captured using pheromone trapping method, ranging from 27 to 47 in open field and 103 to 255 in the greenhouse (Goftishu et al., 2014).

The reproduction potential of this insect is also high with a life cycle that is completed within 30 to 35 days based on the environmental conditions (Harizanova et al., 2009). Such fast reproductive character of the insect makes it more dangerous and notorious in tomato farming system.

The insect is time selective on their activity mostly concentrated in the early morning and dusk, however, in the daytime, they remain hidden among the leaves of the plant (Arnó and Gabarra, 2011). The lifespan of adult insect ranged between 10 and 15 days for females and 6 and 7 days for males. The female moth lays separately dispersed eggs on the underside of leaves especially at the apical shoot (Cherif et al., 2013), stems and sepals to facilitate fast pest distribution. The egg size is small (0.35 mm long) and each moth usually lays 250 to 300 eggs and the insect exhibits 10 to 12 generations in a year (Desneux et al., 2010). The larva is the feeding stage of this pest and can cause 80 to 100% losses in tomato if it is not controlled.

The adaptation ability of *T. absoluta* was also mostly fitted with different environmental conditions. For instance, Van Damme et al. (2015) tested the cold

hardiness and overwintering potential of insect to detect the supercooling point of larvae, pupae, and adults for a lower lethal time at 0 and 5°C, reported diapause incidence at 18 and 25°C. Then they determined the mean supercooling point for pupae (-16.7°C) that was higher than for adults (-17.8°C) and larvae (-18.2°C) in which the most agro-ecology is not situated in such condition. *T. absoluta* also adapts the temperature higher than 49°C in summer as findings showed in Sudan (Tonnang et al., 2015) and able to tolerate hot and dry conditions.

*T. absoluta* is not easily manageable by using chemical pesticides due to two reasons. First the larval stage completes its development inside the leaves mesophyll tissue. Therefore, larvae are not easily exposed to chemical insecticides (Ayalew, 2015). The second challenging thing is the resistance ability of insects towards different pesticides used (Cuthbertson et al., 2013; Ponti et al., 2015). Such pesticide resistance in insect enforces the farmer to try a number of inappropriate chemicals through spraying on their tomato farm. In fact, different types of pesticides have been used for *T. absoluta*, but none of them are significantly successful to control it (Abbes and Chermiti, 2012; Guedes and Picanço, 2012; Guedes and Siqueira, 2013). This continuous application of inappropriate agrochemicals for trial on the farmland has been aggravating residual accumulation in the environment and imposing human health problems (Abdel-Raheem et al., 2015).

Thus, to overcome such complicated problems, different findings are conducted to develop environmental friendly Integrated Pest Management (IPM) strategies. This includes mass trapping of male moths by using sex pheromone (Braham, 2014), using of phytochemicals, using biological agents such as parasitoids, predators, microbial cells, microbial products, small RNA interference (RNAi), inheritance sterility (IS) development in insect and pest resistant plant cultivar production approaches were applied to control *T. absoluta* (Van Lenteren and Bueno, 2003; Abbes et al., 2012; Cagnotti et al., 2012; Camargo et al., 2016). Thus, the main purpose of this review was to compile important findings conducted on eco-friendly control of *T. absoluta* for sustainable healthy farming system.

## HOST PLANT SELECTION MECHANISM AND LIFE CYCLE OF *T. ABSOLUTA*

Several insect pests use volatile attractant secretions that are emitted from the plant for their feeding and oviposition during host selection (McCormick et al., 2012; Beyaert and Hilker, 2014; Bawin et al., 2015). *T. absoluta* has such behavior to select their host plants (Megido et al., 2014) with a high preference towards tomato cultivars (Proffit et al., 2011; Shehata et al., 2016) and mostly

underside of leaves for their oviposition (Cherif et al., 2013). Although tomato is the main host for *T. absoluta*, the other solanaceous plant species also served as alternative hosts.

Mostly, mated females of *T. absoluta* prefer leaves especially apex part for oviposition than the other parts of the tomato plant (Torres et al., 2001; Leite et al., 2004; Proffit et al., 2011; Cherif et al., 2013). This oviposition site preference could be associated with a lower calcium content of apical plant leaves than that of middle and basal leaves of tomato (Leite et al., 1999). The oviposition of *T. absoluta* was mostly mediated by volatile signals generated from their host plant (Proffit et al., 2011). Tomato leaf has volatile compounds such as terpenoid that attracts mated females for oviposition. Plant cultivars, rich in 2-tridecane or zingiberene that are chemicals used to give tomato resistance against beet armyworm are not preferred for oviposition by *T. absoluta* (De Oliveira et al., 2012). However, plants with the absence or less content of terpenes that is herbivores repellent compound are preferable to *T. absoluta* oviposition (Proffit et al., 2011; Shehata et al., 2016). The oviposition site selection is not only determined by plant volatile chemical attractants but also their internal physiological changes during egg laying stage such as ovarian maturation, egg production, and mating status can influence the response of the insect to host signaling (Masante-Roca et al., 2007).

Tomato plants bearing *T. absoluta* eggs produce hexanal and (Z)-3-Hexen-1-ol that is secreted in response to plant damage (Anastasaki et al., 2015). Because, the chemical compound (Z)-3-Hexen-1-ol, identified from the headspace of intact plants (Proffit et al., 2011) is found in very limited concentrations from normal tomato leaves. But this chemical compound was increased dramatically after damaging the leaves by different factors (Buttery et al., 1987). Similarly, several volatile compounds in tomato plants become much more abundant when *T. absoluta* lays its eggs on it. For instance, terpinolene was found only on insect-infested plants but not in normal plants (Angeles Lopez et al., 2012). Furthermore, a number of sesquiterpenes such as  $\delta$ -elemene have been found to increase in the headspace of tomato plants with *T. absoluta* eggs (Anastasaki et al., 2015). However, these compounds are reported to play a role in tomato resistance against insects other than *T. absoluta* (Van der Hoeven et al., 2000; Bleeker et al., 2009). The production of such compound is aggravated when the environmental conditions are changed.


For instance, elevated emissions of sesquiterpenes are found after exposure of tomato plants under cold and hot stress (Copolovici et al., 2012). The discriminate analysis showed that these differences in ratios of similar blends can differentiate control against oviposited tomato plants, indicating an alteration of volatile blend after the oviposition with *T. absoluta* (Anastasaki et al., 2015).

The oviposition of *T. absoluta* shows significant preference and survival longevity towards tomato plants than other species of plants (Bawin et al., 2015). Tomato plants also give higher larval survivability and short development time as compared to the other plants (Bawin et al., 2015; Shehata et al., 2016). Additionally, the leaf contact also significantly increased the number of eggs laid, thus leaf surface morphology and chemistry accordingly have importance for oviposition of *T. absoluta* (Proffit et al., 2011). The leaf trichomes also provide chemical and mechanical stimuli for egg laying by female insect (Torres et al., 2001). The egg develops into first instar larva which starts to wound the leaf to enter into the mesophyll tissue of leaves or other parts of the host plant. The larval (feeding stages) development depends on the nutritional quality and/or production of plant metabolites (Bawin et al., 2015). The matured larvae pupate in the soil, on leaf and other parts of the plant. The pupa is initially soft, cylindrical, and green, later becomes dark brown (Genc, 2016). The pupal stage of the leafminer requires 8 to 15 days under controlled laboratory conditions. Adults become gray-brown and males are darker than females (Figure 1).

## **INSECTICIDE RESISTANCE MECHANISMS OF *T. ABSOLUTA***

Since *T. absoluta* was reported as leafminer of tomato in South America, insecticide spray was the primary controlling method. However, the insect becomes highly resistant towards synthetic insecticide and shows the reduced effectiveness of control (Lietti et al., 2005; Silva et al., 2011). This enforces farmers to apply different pesticides repeatedly on their farmland (Reyes et al., 2012). Such continuous and inappropriate chemical application on farmland aggravates the resistance problem on *T. absoluta* (Silva et al., 2011; Reyes et al., 2012). For instance, *T. absoluta* scored the higher resistant values 8-, 3.79-, 6.4- and 1.84-fold against four insecticides namely indoxacarb, metaflumizone, spinosad, and chlorantraniliprole in Turkey, respectively (Yalcin et al., 2015).

The chemical resistance of insect pest is mediated by enzymatic activities. Biochemical experiments suggest that the enzymatic systems might be involved in the loss of susceptibility of the tomato leafminer to different insecticides. For instance, insecticide spinosad that was registered in Chili has a unique mode of action against Lepidoptera larvae (Cleveland et al., 2002; Wang et al., 2009) and demonstrated low effectiveness towards *T. absoluta*. This is due to resistance enhanced by detoxifying enzymes such as mixed-function oxidases, glutathione-S-transferases, and esterases (Reyes et al., 2012). Those enzymes are found in several insects and make them resistant to different insecticides (Rodriguez et al., 2010; El-Latif et al., 2010; Wu et al., 2011).



**Figure 1.** Life cycle of *T. absoluta*, photographs compiled from Arno et al. (2010) and developmental periods from Erdogan and Babaroglu (2014)

Esterases is non-specific enzyme implicated as an insecticide resistance mechanism in several insect pests due to their ability to hydrolyze insecticidal esters as organophosphates and pyrethroids (Solenio et al., 2008; Rodriguez et al., 2010; El-Latif and Subrahmanyam, 2010; Wu et al., 2011). The glutathione-S-transferase (GST) enzyme activity was determined to facilitate 1.5-fold higher resistance of *T. absoluta* against four insecticides, indoxacarb, metaflumizone, spinosad, and chlorantraniliprole. The esterase (EST) enzyme had a detoxifying activity of these four insecticides in equal performance (Yalcin et al., 2015). In another study, the toxicity of chlorpyrifos is synergized by diethyl maleate and triphenylphosphate. The synergistic effect ranged from 1.3 to 1.9-fold in 24 h and 1.2 to 1.5-fold in 48 h. The exposure with chlorpyrifos for 24 and 48 h significantly increased the activities of esterase. This suggested that esterase is involved in resistance through detoxification of chlorpyrifos in *T. absoluta*. It is

confirmed in field populations of *T. absoluta* from Iran that displayed less susceptibility to chlorpyrifos (Zibae et al., 2016). Furthermore, larval mortality in field was significantly lower by spinosad (47%) as compared to laboratory (91.7%) and the esterase effect ratio varied from 1.7 to 14.7 in the Chili (Reyes et al., 2011).

Moreover, diamide insecticides resistance in *T. absoluta* was reported and correlated to ryanodine receptors (molecular target-site) modification (mutation). It was investigated by reciprocally crossing resistant strain with a susceptible strain and revealed an autosomal incompletely recessive mode of inheritance. Genotyping of individuals of resistant and field-collected strains showing different levels of diamide resistance revealed that the presence of G4903E and I4746M RyR target-site mutations. The two novel amino acid mutations, G4903V and I4746T, in some of the resistant *T. absoluta* strains were detected. The functional performance evidence of these mutations is detected to

alter the affinity of the RyR to diamides effect on *T. absoluta* (Roditakis et al., 2017).

## IPM STRATEGY AND ITS EFFECTIVENESS

Environmental friendly controlling strategies have been developed for protection from *T. absoluta*. This includes, cultural control measures (crop rotation, selective removal and destruction of infested plant material) (Korycinska et al., 2009), using natural enemies (parasitoids, predators, entomopathogens and nematodes) (Desneux et al., 2010; Urbaneja et al., 2012), botanicals and production of resistant tomato cultivar varieties (De Oliveira et al., 2012). The integration of these methods with each other and if possible with less environmental hazardous insecticides are important to control *T. absoluta* without disturbing ecological world. Especially botanicals, natural enemies and microbial cells with their products are considered as promising alternatives than using synthetic pesticides.

### The use of pheromone trap

#### Monitoring and mass trapping

Sex pheromone of insect is serving as alternative to monitor male moth population through trapping (Braham, 2014). Sex pheromone is chemical secretion released in the form of fluid that triggers the opposite sexual interest (Medido et al., 2013; Illakwahhi and Srivastava, 2017). This sex pheromone is released by virgin female tomato leafminer and strongly attractant to male moths (Quiroz, 1978). Mostly male moths are attracted to the secreted pheromone (Braham, 2014). Therefore, mass trapping of male moths is encouraged by using pheromone and imperative to early warning of pest abundance and monitor insect population (Megido et al., 2013; Ghoneim, 2014; Retta and Berhe, 2015). The trapped insects can be damaged mechanically or by applying chemical. These techniques have been successfully applied in controlling leafminers on both greenhouses and open field (Chermiti and Abbes, 2012; El-aassar et al., 2015).

### Biochemical bio-pesticides (Plant extracts)

Botanicals are very important natural resources used to control different agricultural pests for long period of time. Crude extracts from seeds, leaves, bark, bulbs, and fruits of the different plant species have been extensively tested on agricultural pests for bioactivity worldwide (Isman and Seffrin, 2014). The phytochemical products, especially extracts or essential oils are suggested as potential alternatives than synthetic pesticides to control insect pests (Philogène and Vincent, 2005). Ethanolic leaf extract obtained from *Piper amalago* var. *medium*, for instance, caused 70% larval and pupal mortality in two-

day exposure through exhibiting acute toxicity at the concentration of 2,000 mg L<sup>-1</sup> (de Brito et al., 2015). The allelochemicals found in Piperaceae are diverse with insecticidal/insectistatic properties (Scott et al., 2008). From allelochemicals, piperamides are the major classes of compounds found in species of the genus *Piper* that exhibits dual biological activity on insects, neurotoxic and affect lipid metabolism (Dyer and Palmer, 2004) and promising to manage chemical resistant insect populations (Scott et al., 2004). Effects of the knockdown type, with paralysis and immediate death of the insect, has also been reported in insects treated with amides isolated from Piperaceae species (Scott et al., 2007).

The ethanolic extract from the Neem (*Azadirachta indica*) and petroleum ether extract from *Jatropha* (*Jatropha curcus*) seeds are tested on *T. absoluta* egg and larval stage as bio-insecticides (Kona et al., 2014). Thus, the ethanolic extract of Neem results in 24.5% egg and 86.7 to 100% larval mortality of *T. absoluta* at different concentration. The observed larval mortality is better than Neem Azal T/S 0.3% registered with active ingredient of azadiractin which kills 84 to 87% of first to fourth instars larvae after 14 days of exposure (Yankova et al., 2014). In the same way, the petroleum ether extract obtained from *Jatropha* also achieved 18 to 25% egg and 87 to 100% larval death on *T. absoluta* after being exposed for 4 days in different concentration (Moreno et al., 2011; Kona et al., 2014). The extracts obtained from jojoba (*Simmondsia chinensis*) seed at 100% concentration resulted in 75% mortality on 2nd larval instars of *T. absoluta* (Abdel-Baky and Al-Soqeer, 2017). The botanical extracts from "Tossegn" Garden thyme (*Thymus vulgaris*) and "Gulo" Castor bean (*Ricinus communis*) also caused 95 and 58% larval mortality of *T. absoluta*, respectively (Nilahyane et al., 2012). The aqueous extracts obtained from five different plant species showed moderate to high mortality on *T. absoluta* developmental stages. These extracts are from chinaberry (*Melia azedarach*) leaves and fruit caused (91%), geranium (*Pelargonium zonale*) (87%), garlic (*Allium sativum*) (85%), onion (*Allium cepa*) (80%), basil (*Ocimum basilicum*) flower (74%) and leaves (54%) mortalities on 2nd instars larvae at 6% concentration after 5 days exposure (Ghanim and Ghani, 2014). In another study, garlic also acts as a repellent for larvae in 37.5% and caused weight loss on first larval instars of *T. absoluta* (Salama and Shehata, 2017). This resulted in 78% 1st instars larval mortality after 4 days of exposure to 2000 ppm oil extract concentration. Clover (*Trifolium repens*) and *Eucalyptus* (*Eucalyptus camaldulensis*) oil extract also caused moderate (67 and 63%) mortality of 1st instars larvae at the highest (2000 ppm) concentration after 6 days exposure. Thus, crude extract concentration and exposure time are considered as the determinant factors for effectiveness of plant phytochemicals against *T. absoluta*.

The hexane and ethanol extracts from aerial parts of

*Acmella oleracea* exhibited the highest activity, causing (100 and 88%) mortality, respectively, in *T. absoluta* at a concentration of 10 µg of extract per mg of an insect after 6 h of exposure (Moreno et al., 2011). The bioactivity of *A. oleracea* is due to alkaloids present in the plant. The main active amide in the plant is an isobutylamide (2*E*, 6*Z*, 8*E*)-*N*-isobutyldeca-2, 6, 8-trienamide, commonly known as spilanthol (Ley et al., 2006; Wu et al., 2008). Three alkaloids were identified in the bioactive fractions of the hexane extract of *A. oleracea* [spilanthol, (*E*)-*N*-isobutylundeca-2-en-8, 10-dienamide and (*R*, *E*)-*N*-(2-methyl butyl) undeca-2-en-8,10-dienamide]. The results showed that all of the compounds isolated had a high insecticidal activity that was at least as toxic as permethrin, a pyrethroid recommended for control of *T. absoluta* (Moreno et al., 2011). The mechanism of action is that active alkaloids found in *A. oleracea* has not yet been determined. However, it is roughly proposed that it affects the nervous system, as evident from abnormal movement like uncoordinated muscular activity. This effect suggests that the compounds disturb nerve conduction somewhere. Furthermore, crude extracts of three plant *A. indica* seed, *Cymbopogon citrates*, and *A. sativum* also caused 98, 97 and 95% mortality at larval stage of *T. absoluta*, respectively in 7 days of exposure in Ethiopia (Shiberu and Getu, 2017).

## Biological control

### *The use of parasitoids and predators*

The entomophagous insects, predators and parasitoids are natural enemies of insect and considered as alternative agents for pest control. As Luna et al. (2012) reviewed, more than 50 species of natural enemies are categorized under predators, parasitoids, and entomopathogens and used for *T. absoluta* control. Among these enemies, numerous studies are focused on parasitoid that approximately accounts for 20 species and considered as effective in Lepidopteran pest control (Polack et al., 2011). Moreover, six parasitoid groups are found to attack *T. absoluta* eggs, larvae and/or pupae (Luna et al., 2012). Specifically, Hymenopteran wasps, Tachinidae (Diptera) species, mirid bug *Tupiocoris cucurbitaceus* (Spinola) and the parasitoid wasps *Dineulophus phtorimaeae* (de Santis) and *Pseudapanteles dingus* showed positive traits on biological control aspect on insects (Luna et al., 2010; Lopez et al., 2011).

As investigators reported, *Trichogrammatoidea bactrae* and *Trichogramma pretiosum* are egg parasitoids of *T. absoluta* (Lewis et al., 2003). However, the endoparasitoid, *P. dignus*, and the ectoparasitoid, *D. phtorimaeae* formed more than 50% of natural parasitism in the larval stage of *T. absoluta* (Sánchez et al., 2009; Luna et al., 2010) and result in larval death due to nutrient consumption. The

egg parasitoids are preferable since if they are effective, they can rapidly reduce the crop damage, whereas the larval and pupal parasitoids will control the pest once they result in the damage on the fruits, leaves and stems of the plant (Newton, 1998). Due to this, most of the reports are focused on egg parasitoids and they are dominantly categorized under Order Hymenoptera. The most important *T. absoluta* egg parasitoids are found in the families of Trichogrammatidae, Encyrtidae, and Eupelmidae (Ghoneim, 2014). About 210 species of *Trichogramma* are signaled as natural enemies of a variety of agricultural and forest pests in many regions of the world and at least 12 species are widely used commercially in biological control programs (Pratissoli et al., 2005; Kumar et al., 2009; Suckling and Brockerhoff, 2010). This parasitoid has short generation time and they can be easily reared on respective hosts (Mansour, 2010). *Trichogramma pretiosum* and *Trichogramma exiguum* are egg parasitoids under the family Trichogrammatidae and the most extensively used to control tomato leafminer *T. absoluta* (Desneux et al., 2010; Molla et al., 2011; Öztemiz, 2012; Vasconcelos, 2013). Furthermore, several reports indicated the existence of *T. absoluta* predators that participate in the population reduction. For instance, *Nesidiocoris tenuis* Reuter (Hemiptera: Miridae) regulates *T. absoluta* population due to its ability to prey efficiently on eggs (Molla et al., 2011). This predator species was able to prey on more than 100 eggs per individual per day (Arno et al., 2009). Two predator species, *Mesoglossus pygmaeus* and *N. tenuis* also reported as they reduced leaflet infestations up to 75 and 97% and fruit infestations up to 56 and 100%, respectively (Molla et al., 2009); though, preying on the egg and larvae of *T. absoluta* (Gadarrá and Arno, 2010). However, until now there are no any reports that indicate the existence of adult parasitoids of *T. absoluta* (Ghoneim, 2014).

The susceptibility of insect larvae and pupae to entomopathogenic nematodes (*Steinernema carpocapsae*, *Steinernema feltiae*, and *Heterorhabditis bacteriophora*) was reported (Grewal et al., 2005). This is indicated in leaf bioassays evaluation of tomato using the nematodes for the capability to reach the larvae and kill them within the galleries (Rodríguez et al., 2005). Such experimental evaluation showed high larval mortality (79 to 100%) and low pupae mortality of only 10% (Rodríguez et al., 2005; Batalla-Carrera et al., 2010). In the leaf bioassay, a high level of larval parasitization (77 to 92%) was recorded revealing the nematode's capacity to kill the larvae inside the galleries. The pot experiments indicated that nematode treatment reduced insect infection of tomato plants by 87 to 95% (Batalla-Carrera et al., 2010).

### *The use of microbial cell*

The plant growth promoting the effect of soil microbes

has been shown to affect plant-insect interactions (Pineda et al., 2010). Improvement of the nutrient composition increases plant nutritional quality, while it affects insect performance at several trophic levels (Schoonhoven et al., 2005; Zhang et al., 2008). Systemic responses that plant root microorganisms cause emission of volatiles at aerial parts of plants in multitrophic interactions that attract above-ground parasitoids (Bruce and Pickett, 2007). Furthermore, plants interacting with beneficial microbes can also benefit from an increase in tolerance to herbivory (Bennett et al., 2008; Vannette and Hunter, 2008). This can be achieved by improving nutrient and water uptake facility for better plant development. Furthermore, beneficial microbes can facilitate the regrowth of tissues after herbivory and promote plant tolerance, which is reflected in compensation of the loss of plant biomass in the presence of herbivores (Herman et al., 2008; Kempel et al., 2009). For instance, *Trichoderma* species is plant growth promoting fungus that is commercially available and widely used in agriculture that induces systemic resistance against microbial pathogens in plants (Harman et al., 2004; Trillas et al., 2009; Segarra et al., 2009). When the *Trichoderma* spp. interacts with the plants, they recognize the existence of the species through the signal communication and become ready to defend other invaders. However, the plant-mediated effects on insects are not assessed (Pineda et al., 2010).

There are several soil microbes that have the ability to induce plant resistance in systemic tissues (induced systemic resistance, ISR). Induced systemic resistance in the plant is resulted by plant growth promoting bacteria, *Pseudomonas* and *Bacillus* species (Van Wees et al., 2008; Van der Ent et al., 2009), root mycorrhizal fungi (Trillas et al., 2009), areal stem endophytic fungi (Stein et al., 2008) and soil-plant growth promoting fungi (Segarra et al., 2009). The ISR efficiency is proven in different plant species against pathogenic bacteria, fungi, viruses, nematodes and recently insect pests (Van Oosten et al., 2008; Stein et al., 2008). The priming defense genes show a higher expression systemically in the leaves only after pathogen or insects attack in responsiveness of the plant to the plant hormones jasmonic acid and ethylene (Van Wees et al., 2008; Van der Ent et al., 2009). The levels of these hormones remain unchanged, suggesting that ISR is based on increased sensitivity to plant hormones rather than on the increased production of such hormones (Van der Ent et al., 2009).

Priming is a phenomenon that provides plants with an enhanced capacity to rapidly and effectively mount defense responses to biotic and abiotic stresses (Conrath et al., 2006). It is elicited by beneficial microbes, but also by pathogens, herbivores and selected synthetic compounds such as benzothiadiazole and b-aminobutyric acid (Heil and Silva, 2007; Frost et al., 2008). Typically, priming is characterized by accelerated defense-related gene expression once primed plants are attacked by a

pathogen or an insect, resulting in an enhanced level of resistance against the invader (Pineda et al., 2010). This enhanced resistance is effective against a broad spectrum of attackers. The molecular basis of priming is poorly understood, but recent advances in the field of priming demonstrate that the accumulation of latent defense-related transcription factors (Van der Ent et al., 2009), mitogen-activated protein kinases (MAP kinases) (Beckers et al., 2009) and secondary metabolites such as azelaic acid (Jung et al., 2009) and volatile organic compounds (Heil and Silva, 2007) participate on establishment of the primed state.

### ***The use of microbial product***

The bacterium *Bacillus thuringiensis* is a gram-positive spore-forming bacterium that produces toxic crystal proteins (Glare, 2000). This protein is a delta-endotoxin protein that is eaten by insect larvae. Then the protoxin is dissolved in the parasporal bodies and activated under alkaline conditions in the mid-gut of target insects, thereby releasing the active peptides that bind to specific receptors in the insect's mid-gut epithelial cells and creating pores in the epithelial membrane and stop insect feeding and ultimately result in death due to starvation (Charles et al., 1996). Thus, this bacterium produces such protein crystals during sporulation and the product is toxic to insect larvae and causes rapid death (Rezaei and Taleai-Hassanloui, 2016). The pure protein crystals from five *B. thuringiensis* strains that were taken from dead or diseased larvae exhibited satisfactory efficacy against *T. absoluta* larvae (Medeiros et al., 2006).

*B. thuringiensis* that was isolated from local soil show three folds toxicity against *T. absoluta* larvae compared to one of commercial *B. thuringiensis* in Algeria (Hernandez-Fernandez et al., 2010). Furthermore, four (B1, B2, B3, B4) *Bacillus* spp. isolated from dead larvae in Egypt caused the larval mortality of *T. absoluta* at 4th instars and the effect ranged between 80 and 93% in seven days interval (Youssef and Hassan, 2013). The effectiveness was increased (96.7%) at a concentration (2 g/ml) of isolates (Youssef and Hassan, 2013). This is an indicator of crystal protein that as it increase the concentration of *Bacillus* spp., it increases the mortality. Toxins expressed by *Bacillus* spp. naturally colonize the phylloplane of tomato plants and survive for a period of 45 days on the tomato leaf surface (Theoduloz et al., 2003) that result in mortality on first to fourth larval instars, however, the death variation occurred due to feeding behavior difference (Giustolin et al., 2001; Youssef and Hassan, 2013). The first and second larval instars penetrate directly the leaves without much feeding. Therefore, they are exposed to a lower dose of bacterial spores and toxins. Older instars are more susceptible to treatments than younger ones due to their longer stay or larval movement in and out of the mines

and galleries several times during their development and at that moment they are very vulnerable to infection by the bacterial spores (Harizanova et al., 2009).

In the field condition, *B. thuringiensis* after 50 days of the application showed a significant average reduction of *T. absoluta* infestation to 12.4%, individuals as compared to  $26.5 \pm 11.34$  individuals in the control. Furthermore, after 90 days of the post applications, the means number of infestations significantly decreased to 46% as compared to 30.6 individuals in the control. The weight of the tomatoes was determined as better scored after harvest (Sabbour and Soliman, 2012).

The entomopathogenic fungus *Baeuveria bassiana* was considered as the best and effective candidate against larvae of *T. absoluta*. These fungal strain, with high dose treatment ( $4.75 \times 10^7$  conidia/ml), resulted in 100% larval death within 3 days, while low doses led in a fatality rate of 87% within 4 days (Ikram and Abdallah, 2011). Another investigator also reported that *B. bassiana* that caused 68% *T. absoluta* death at larval stage (Cabello et al., 2009) *Metarhizium anisopliae* also kills 37% of female moths (*T. absoluta*) (Cristina et al., 2008). In Egypt, the used *M. anisopliae* and *B. bassiana* reduced *T. absoluta* infestation by  $9.8 \pm 9.9$  and  $10.8 \pm 11.9$  individuals as compared to  $21.7 \pm 8.6$  individuals in the control under greenhouse conditions (Sabbour, 2014).

## RNA INTERFERENCE (RNAI)

The RNA interference (RNAi) is a mechanism that small RNAs can rapidly cause post-transcriptional specific gene silencing (Hannon, 2002). Such biotechnological approach is currently becoming the potential application for agricultural insect pest control (Gordon and Waterhouse, 2007; Katoch et al., 2013). Tomato pinworm, *T. absoluta* mitigation was initiated by targeting on gene silencing approaches using small RNA interference (Price and Gatehouse, 2008; Camargo et al., 2016). RNA interference (RNAi) gene silencing process is performed by providing double-stranded RNA (dsRNA) molecules that match a specific target gene sequences in a targeted organism (Koch and Kogel, 2014).

This approach works when the plant was engineered to express insect dsRNAs (Price and Gatehouse, 2008). However, obtaining sufficient, effective and essential genes used for insect development and survival was a critical step (Zhang et al., 2013). Thus, the first gene screening from *T. absoluta* was made using insect hormone biosynthetic pathways as potential targets for systemic silencing (Camargo et al., 2015). The production of transgenic plants expressing dsRNA molecules targeted to essential insect genes was providing a means of specific gene silencing in plant-feeding larvae. Then the larval phenotypes are developed loss of appetite to death (Camargo et al., 2016). The first experiment of such method on the *T. absoluta* was conducted by Camargo et al. (2016) using two

approaches to dsRNA delivery for larvae. The supply technique is allowing larvae to feed upon the tomato leaves that contains dsRNA obtained through absorption from aqueous solution through sap-sucking of *Bemisia tabaci* (Luan et al., 2013) and transient transcription of dsRNA by the host plant after the infiltration of *Agrobacterium* cells carrying binary plasmids that expressed hairpin versions of the target gene sequences ('agro-infiltration') (Leckie and Stewart, 2010). This method is designated as "*in planta-induced transient gene silencing*" (PITGS) that successfully delivered hairpin silencing genes in other pathogens (Panwar et al., 2012).

The Vacuolar ATPase catalytic subunit A gene (V-ATPase) and Arginine kinase were targeted gene for this experiment (Camargo et al., 2016). The H<sup>+</sup>-ATPase vacuolar pump is one of the essential enzymes present in almost all eukaryotic cells that are responsible for generating energy gradients in many membranes and organelles (Nelson et al., 2000; Wieczorek et al., 2009). Moreover, arginine kinase belongs to a transferase protein family that catalyzes the transfer of a high energy phosphate group from ATP to L-arginine to yield phosphor-arginine used for energy storage (Bragg et al., 2012; Kola et al., 2015). Then the report was indicating that silencing of the insect V-ATPase and Arginine kinase genes by dsRNA showed 60% reduction in target gene transcript accumulation and increase larval mortality (Camargo et al., 2016). Development of transgenic tomato lines expressing the hairpin version of these genes demonstrated that this approach can adversely affect insect development and viability by reducing the expression of insect target genes. Thus, leaf-absorbed dsRNA delivery avoids cloning steps into specific vectors and allows a better control of provided dsRNA amounts in the leaves (Luan et al., 2013; Camargo et al., 2016), whereas PITGS offers a more realistic trigger of the RNAi machinery in transgenic plants, since the dsRNA hairpin molecule will be transcribed and processed by the plant cells (Camargo et al., 2016).

The findings are suggesting that in addition to the easy provision of dsRNAs for insect pests, sufficient dsRNA accumulation is also required to produce gene silencing in *T. absoluta* at least minimum dose before triggering RNAi (Yu et al., 2012). This requires trans-membrane proteins, such as systemic RNA interference deficient-1 proteins (SID-1) that involved in dsRNA uptake and systemic spreading in the tissues (May and Plasterk, 2005) and then putative orthologous SID-1 genes in the *T. absoluta* transcriptome was identified by RNA-seq analysis (Camargo et al., 2015) for transportation.

## INHERITED STERILITY TECHNIQUES (IST)

Inherited sterility techniques are commonly practiced to control different agricultural insect pests (Bloem and

Carpenter, 2001; Carpenter et al., 2005; Boshra, 2007). It does not exert any effect on the environment and impotent approach to control especially invasive pest in an agricultural system like *T. absoluta* (Cagnotti et al., 2012). The radiation sterilization of insects and sterile male insect release into a wild population of the same species was expected to prevent the reproduction of wild females and practiced for pest control (Knipling, 1955). At the first time such sterilization has used  $\gamma$ -radiation with a highly radioactive material, although, it is currently not available in the market due to difficulties related to the transportation of co-sources (Mastrangelo et al., 2010; Mehta and Parker, 2011). Alternatively, X-irradiators are used for sterilization of insect pests. It is easier and independent of a radioactive source that is performed safely on insect sterility (Cagnotti et al., 2012).

The inheritance sterilization works in the way that the irradiated male insect is released into the natural population and mate with an untreated female to produce abnormality in offspring. In this case, most radiation-induced chromosomal breaks will not lead to the loss of chromosome fragments and remains stable during both mitosis and meiosis and can be transmitted through germ cells to the next generation (Marec et al., 2001; Carpenter et al., 2005). When partially sterile males mate with wild fertile females their deleterious effects induced by radiation are inherited and expressed for several generations, mainly in the first generation of descendants of irradiated adults (Carpenter et al., 2005). Such practice is firstly conducted by Cagnotti et al. (2012) on the *T. absoluta* and considered as an effective technique to control it. The optimum dose of X-radiation to make male moth sterile is between 200 and 250 Gy (Cagnotti et al., 2012). In the first generation, the egg hatching is reduced and the produced offspring becomes highly sterile (more than the irradiated parent moths) and most are male (Bloem and Carpenter, 2001; Seth and Sharma, 2001). Moreover, adult emergence decreased at doses of X-radiation increased the results into formation of malformed wings and bent legs at doses >350 Gy (Cagnotti et al., 2012).

Thus, the inherited sterility of insect pest technique is considered as a key strategy to manage *T. absoluta* and it is considered as the area-wide concept of integrated pest management system (Carpenter et al., 2005; Cagnotti et al., 2012). This inherited sterility way of pest management is compatible with the use of other area-wide control tactics such as mating disruption, biological control, cultural control methods, and the use of bio-rational pesticides (Bloem and Carpenter, 2001; Carpenter et al., 2005).

## SYNERGETIC RESISTANCE DEVELOPMENT OF HOST TOMATO

Insect resistant cultivar development is considered as

one of alternative approach than chemical pesticides for obtaining cost-benefit ratio for pest control (Pereira et al., 2008; Gonçalves et al., 2010; Maluf et al., 2010). In Brazil, tomato improvement was practiced to obtain pest-resistant cultivars by incorporating the alleles for resistance present in wild plants for commercialization and produce the allele-chemicals associated with resistance (Silva et al., 2009; Gonçalves et al., 2010; Maluf et al., 2010). Three types of allele-chemicals (acyl-sugars, zingiberene, and 2-tridecanone) have been associated as resistant to *T. absoluta* (Pereira et al., 2008; Silva et al., 2009; Maluf et al., 2010). However, studies comparing levels of pest resistance among strains rich in different allele-chemicals are few (De Oliveira et al., 2012).

The resistance of tomato strains rich in 2-tridecanone (2-TD), zingiberene (ZGB), and acyl sugars (AA) to the tomato moth, *T. absoluta* showed significant reduction on evaluated characteristics, such as oviposition rate, severity of damage to the plants, injuries to the leaflets, and the percentage of leaflets attacked (De Oliveira et al., 2012). The oviposition rate reduction of *T. absoluta* was directly associated with high concentrations of 2-TD, ZGB, and AA in the tomato cultivars (Silva et al., 2009; Gonçalves et al., 2010; Maluf et al., 2010). This is a very important strategy to prevent the tomato plant from insect attack. These three allele-chemicals had equivalent effects on oviposition, which was consistently lower on these strains than on the susceptible controls (De Oliveira et al., 2012).

## CONCLUSION

The agricultural pests are main threats for the vegetables, fruits, cereals and flower production in several countries. *T. absoluta* is one of the invasive tomato pest and it is devastating worldwide. The insect will result in 80 to 100% crop damage if it is not controlled. The entrances of larvae into leaf mesophyll prevent them from pesticides and save them from death. Furthermore, *T. absoluta* has developed resistance to different types of pesticides and become too difficult to control it using insecticides. The repeated spraying of synthetic pesticides on the farmland is becoming problematic to environmental safety.

The IPM strategy is very important and has dual purpose, environmental friendly and effective to manage insect pests. Biological control is one of IPM and potential to mitigate insect pests from farming system. Microbial groups both bacterial and fungal species are showing important implication on *T. absoluta* control. Furthermore, biotechnological techniques are also another alternative to control tomato leafminer *T. absoluta*. It is a newly coming technology mostly targeted on RNA interference. Small RNA interference that hairpin formation of introduced double serenaded RNA interfaces the

essential component synthesis in the insect and stop the insect growth. Thus, IPM strategy is then the promising alternative to control *T. absoluta* for healthy farming of the tomato vegetable.

## CONFLICT OF INTERESTS

The author has not declared any conflict of interests.

## REFERENCES

- Abbes K, Chermiti B (2012). Failure of the biological control of *Tuta absoluta* using the predator *Nesidiocoris tenuis* in a protected tomato crop: analysis of factors. IOBC/WPRS Bull. 80:231-236.
- Abbes K, Harbi A, Chermiti, B (2012). The tomato leafminer *Tuta absoluta* (Meyrick) in Tunisia: current status and management strategies. EPPO Bull. 42:226-233.
- Abdel-Baky FN, Al-Soqeer AA (2017). Controlling the 2nd instars larvae of *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae) by simmondsin extracted from Jojoba seeds in KSA. J. Entomol. 14:73-80.
- Abdel-Raheem A, Ismail A, Abdel-Rahman S, Abdel-Rhman E, Naglaa F (2015). Efficacy of Three Entomopathogenic Fungi on Tomato leaf miner, *Tuta absoluta* in Tomato crop in Egypt. Swift J. Agric. Res. 1:015-021.
- Anastasaki E, George B, Nikos E, Antonios N, Panagiotis G (2015). Oviposition induced volatiles in tomato plants. Phytochemistry Letters. 13:262-266.
- Angeles Lopez Y, Martinez-Gallardo N, Ramirez-Romero R, Lopez M, Sanchez-Hernandez C, Delano-Frier J (2012). Cross-kingdom effects of plant-plant signaling via volatile organic compounds emitted by tomato (*Solanum lycopersicum*) plants infested by the greenhouse whitefly (*Trialeurodes vaporariorum*). J. Chem. Ecol. 38:1376-1386.
- Arnó J, Gabarra R (2010). Controlling *Tuta absoluta*, a new invasive pest in Europe. Training in Integrated Pest Management. ENDURE, IRTA, Cabriels, Spain.
- Arno J, Sorribas R, Prat M, Matas M, Pozo C, Rodri'guez D, Garreta A, Gabarra, R. (2009). *Tuta absoluta*, a new pest in IPM tomatoes in the Northeast of Spain. IOBC/WPRS Bull. 49:203-208.
- Arno' J, Gabarra R (2011) Side effects of selected insecticides on the *Tuta absoluta* (Lepidoptera: Gelechiidae) predators *Macrolophus pygmaeus* and *Nesidiocoris tenuis* (Hemiptera: Miridae). J Pest Sci. 84:513-520
- Ayalew G (2015). Efficacy of selected insecticides against the South American tomato moth, *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae) on tomato in the Central Rift Valley of Ethiopia. Afr. Entomol. 23:410-417.
- Batalla-Carrera L, Morton A, Garcidel-Pino F (2010). Efficacy of entomopathogenic nematodes against the tomato leafminer *Tuta absoluta* in laboratory and greenhouse conditions. Bio Cont. 55:523-530.
- Bawin T, Dujeu D, De Backer L, Francis F Verheggen J (2015). Ability of *Tuta absoluta* (Lepidoptera: Gelechiidae) to develop on alternative host plant species. Can. Entomol. pp. 1-9.
- Bawin T, Dujeu D, De Backer L, Fauconnier M, Lognay G, Delaplace P, Francis F, Verheggen JF (2015). Could alternative solanaceous hosts act as refuges for the tomato leafminer, *Tuta absoluta*. Arthropod-Plant Interac. 9:425-435.
- Beckers GJ, Jaskiewicz M, Liu Y, Underwood WR, He S, Zhang S, Conrath U (2009). Mitogen-activated protein kinases 3 and 6 are required for full priming of stress responses in *Arabidopsis thaliana*. Plant Cell 21:944-953.
- Bennett AE, Alers-Garcia J, Bever JD (2008). Three-way interactions among mutualistic mycorrhizal fungi, plants, and plant enemies: hypotheses and synthesis. Am. Nat. 167:141-152.
- Beyaert I, Hilker M (2014). Plant odor plumes as mediators of plant-insect interactions. Biol. Rev. 89:68-81.
- Biondi A, Zappalà L, Desneux N, Aparo A, Siscaro G, Rapisarda C Garzia T (2005). Potential toxicity of  $\alpha$ -cypermethrin-treated nets on *Tuta absoluta* (Lepidoptera: Gelechiidae). J. Econ. Entomol. 108:1191-1197.
- Bleeker PM, Diergaarde PJ, Ament K, Guerra J, Weidner M, Schutz S, De Both MTJ, Harding MA, Schuurink RC (2009). The role of specific tomato volatiles in tomato-whitefly interaction. Plant Physiol. 151:925-935.
- Bloem S, Carpenter JE (2001). Evaluation of population suppression by irradiated Lepidoptera and their progeny. Fla. Entomol. 84:165-170.
- Boshra SA (2007). Effect of high-temperature pre-irradiation on reproduction and mating competitiveness of male *Sitotroga cerealella* (Olivier) and their F1 progeny. J. Stored Prod. Res. 43:73-78.
- Bragg J, Rajkovic A, Anderson C, Curtis R, Van Houten J, Begres B, Naples C, Snider M, Fraga D (2012). Identification and characterization of a putative Arginine kinase homology from *Myxococcus Xanthus* required for fruiting body formation and cell differentiation. J. Bacteriol. 194:2668-2676.
- Braham M (2014). Sex pheromone traps for monitoring the tomato leafminer, *Tuta absoluta*: effect of colored traps and field weathering of lure on male captures. Res. J. Agric. Environ. Manage. 3: 290-298.
- Brevault T, Sylla S, Diatte M, Bernadas G, Diarra K (2014). *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae): a new threat to tomato production in sub-Saharan Africa: Short communications. Afr. Entomol. 22:441-444.
- Bruce AJT, Pickett AJ (2007). Plant defense signaling induced by biotic attacks. Curr. Opi. Plant Biol. 10:387-392.
- Buttery RG, Teranishi R, Ling LC (1987). Fresh tomato aroma volatiles: a quantitative study. J. Agric. Food Chem. 35:540-544.
- Cabello T, Gallego RJ, Vila E, Soler A, Pino M, Carnero A, Hernández-Suárez E, Polaszek A (2009). Biological control of the South American tomato Pinworm, *Tuta absoluta* (Lep.: Gelechiidae), with releases of *Trichogramma Achaea* (Hym.:Trichogrammatidae) on tomato greenhouse of Spain, Centro Internacional de la Papa, Lima (PE).
- Cagnotti LC, Viscarret MM, Riquelme BM, Botto NE, Carabajal ZL, Segura FD, Lo'pez NS (2012). Effects of X-rays on *Tuta absoluta* for use in inherited sterility programmes. J. Pest Sci. 85:413-421.
- Camargo AR, Guilherme O, Possignolo P I, Peres PL, Lam E, Lima EJ, Figueira A, Marques-Souza H (2016). RNA interference as a gene silencing tool to control *Tuta absoluta* in tomato (*Solanum lycopersicum*). Peer J. 4:1-24.
- Camargo AR, Herai HR, Santos NL, Bento MF, Lima EJ, Marques-Souza H, Figueira A (2015). De novo transcriptome assembly and analysis to identify potential gene targets for RNAi-mediated control of the tomato leafminer (*Tuta absoluta*). BMC Genom. 16:635-642.
- Carpenter JE, Bloem S, Marec F (2005). Inherited sterility in insects. In: Dyck VA, Hendrichs J, Robinson AS (eds) Sterile insect technique principles and practice in area-wide integrated pest management, Chap 2.4. Springer, Netherlands, pp. 115-146.
- Charles JF, Nielson-LeRoux C, Deelucse A (1996). A *Bacillus sphaericus* toxins: Molecular biology and mode of action. Ann. Rev. Entomol. 41:451- 472.
- Cherif A, Mansour R, Grissa-Lebdi K (2013). Biological aspects of tomato leafminer *Tuta absoluta* (Lepidoptera: Gelechiidae) in conditions of Northeastern Tunisia: possible implications for pest management. Environ. Exp. Biol. 11:179-184.
- Chermiti B, Abbes K (2012). Comparison of pheromone lures used in mass trapping to control the tomato leafminer *Tuta absoluta* (Meyrick, 1917) in industrial tomato crops in Kairouan (Tunisia). OEPP/EPPO Bull. 42:241-248
- Cleveland BC, Mayes AM, Cryer S. (2002). An ecological risk assessment for spinosad use on cotton. Pest Manag. Sci. 58:70-84.
- Conrath U, Backers MJG, Flors V, Garcia-Agustin P, Jakob G, Mauch F (2006). Priming: Getting ready for battle. Mol. Plant- Microbe Interact. 19:1062-1071.
- Copolovici L, Kännaste A, Pazouki L, Niinemets Ü (2012). Emissions of green leaf volatiles and terpenoids from *Solanum lycopersicum* are quantitatively related to the severity of cold and heat shock treatments. J. Plant Physiol. 169:664-672.
- Cristina AF, Torres BJ, Fernandes VMA, Farias IMA (2008). Parasitism of *Tuta absoluta* in tomato plants by *Trichogramma pretiosum* riley in

- response to host density and plant structures. *Ciência Rural*, Santa Maria, 38: 1504-1509.
- Cuthbertson G, Mathers J, Blackburn F, Korycinska A, Luo W, Jacobson J, Northing P (2013). Population development of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) under simulated UK glasshouse conditions. *Insects* 4:185-197.
- De Brito FE, Baldin LLE, Silva MCR, Ribeiro PL, Vendramim DJ (2015). Bioactivity of *Piper* extracts on *Tuta absoluta* (Lepidoptera: Gelechiidae) in tomato. *Pesq. agropec. bras.* Brasília, 50: 196-202.
- De Oliveira CM, Junior CV, Maluf RW, Neiva PI, Maciel MG (2012). Resistance of tomato strains to the moth *Tuta absoluta* imparted by allelochemicals and trichome density. *Ciência e Agrotecnol.* 36:45-52.
- Desneux N, Wajnberg E, Wyckhuys KA, Burgio G, Arpaia S, Narvaez-Vasquez C A, Gonzalez-Cabrera J, Catalan Ruescas D, Tabone E, Frandon J, Pizzol J, Poncet Cabello T, Urbaneja A (2010). Biological invasion of European tomato crops by *Tuta absoluta*: ecology, geographic expansion and prospects for biological control. *J. Pest Sci.* 83:197- 215.
- Dyer LA, Palmer ADN (2004). *Piper*: a model genus for studies of phytochemistry, ecology, and evolution. New York: Kluwer Academic: Plenum Publishers, P 228.
- EI-aassar MR, Soliman MHA, Elaai AAA (2015). Efficiency of sex pheromone traps and some bio and chemical insecticides against tomato borer larvae, *Tuta absoluta* (Meyrick) and estimate the damages of leaves and fruit tomato plant. *Ann. Agric. Sci.* 60:153-156.
- EI-Latif AOA Subrahmanyam B (2010). Pyrethroid synergists suppress ester as emediated resistance in Indian strains of the cotton bollworm, *Helicoverpa armigera* (Hübner), Pesticide. *Biochem. Physiol.* 97:279-288.
- Erdogan P, Babaroglu EN (2014). Life table of the tomato leaf miner, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *JAFAG*, 31: 80-89.
- Fortune A (2013). Tomato in danger: newspaper report, 14:230-239.
- Frost CJ (2008). Plant defense priming against herbivores: getting ready for a different battle. *Plant Physiol.* 146:818-824.
- Gadarrá R, Arno J (2010). Resultados de las experiencias de control biológico de la polilla del tomate en cultivo de invernadero y aire libre en Cataluña. *Phytoma España*, 217:65-68.
- Genc H (2016). The tomato leafminer, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae): pupal key characters for sexing individuals. *Turk. J. Zool.* 40:801-805.
- Ghanim MN, Ghani ABS (2014). Controlling *Tuta absoluta* (Lepidoptera: Gelechiidae) and *Aphis gossypii* (Hemiptera: Aphididae) by aqueous plant extracts. *J. Life Sci.* 11:299-307.
- Ghoneim K (2014). Parasitic insects and mites as potential biocontrol agents for a devastating pest of tomato, *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae) in the world: a review. *Int. J. Adv. Res.* 2:81-115.
- Giustolin TA, Vendramim JD, Alves SB, Vieira SA, Pereira RM (2001). Susceptibility of *Tuta absoluta* (Meyrick) (Lep., Gelechiidae) reared on two species of *Lycopersicon* to *Bacillus thuringiensis* var. *kurstaki*. *J. Appl. Entomol.* 125: 551-556.
- Glare RT, Callaghan M (2008). *Bacillus thuringiensis*: Biology, ecology, safety and Larval Serpentine Leaf miners *Liriomyza acicerina* (Rondani, 1875) (Diptera: Agromyzidae) biocontrol. *J. Appl. Sci. Res.* 4:1191-1198.
- Goftishu M, Seid A, Dechassa N (2014). Occurrence and population dynamics of tomato leaf miner *Tuta absoluta* (Meyrick), Lepidoptera: Gelechiidae] in Eastern Ethiopia. *East Afr. J. Sci.* 8:59-64.
- Gonçalves NAC, Silva VDF, Maluf WR, Maciel GM, Nizio DA, Gomes LA, Azevedo SMD (2010). Resistência à traça-dotomateiro em plantas com altos teores de acilglicosídeos nas folhas. *Hortic. Bras.* 28:203-208.
- Gontijo PC, Picanço MC, Pereira EJJ, Martins JC, Chediak M, Guedes RC (2013). Spatial and Temporal Variation in the Control Failure Likelihood of the Tomato Leaf Miner, *Tuta absoluta*. *Annals Appl. Biol.* 162:50-59.
- Gordon K, Waterhouse PM (2007). RNAi for insect-proof plants. *Nature Biotech.* 25:1231-1232.
- Gray Q, Abu T, Teddy T (2013). Tomato production in Ethiopia Challenged by Pest GAIN Report. Global Agricultural Information Network (GAIN), Addis Ababa Ethiopia, Number: 1305.
- Grewal PS, Ehlers RU, Shapiro-Ilan DI (2005). Nematodes as biocontrol agents. CABI Publishing, Oxon, UK P 528.
- Guedes C, Siqueira A (2013). The tomato borer *Tuta absoluta*: insecticide resistance and control failure. *Plant Sci. Rev.* 7:1-7.
- Guedes RNC, Picanc,o MC (2012) The tomato borer *Tuta absoluta* in South America: Pest status, management and insecticide resistance. *EPPO Bull.* 42:211-216.
- Hannon GJ (2002). RNA interference. *Nature* 418:244-251.
- Harbi A, Abbes K, Chermiti B (2012). Evaluation of two methods for the protection of tomato crops against the tomato leafminer *Tuta absoluta* (Meyrick) under greenhouses in Tunisia. *EPPO. Bull.* 42:317-321.
- Harizanova V, Stoeva A, Mohamedova M (2009). Tomato leaf miner, *Tuta absoluta* (Povolny) (Lepidoptera: Gelechiidae)- First record in Bulgaria. *Agric. Sci. Technol.* 1:95- 98.
- Harman GE (2004). Trichoderma species-opportunistic, avirulent plant symbionts. *Nat. Rev. Microbiol.* 2:43-56.
- Heil M, Bueno JC (2007). Within-plant signaling by volatiles leads to induction and priming of an indirect plant defense in nature. *Proc. Natl. Acad. Sci.* 104:5467-5472.
- Herman MAB (2008). Effects of plant growth-promoting rhizobacteria on bell pepper production and green peach aphid infestations in New York. *Crop Prot.* 27:996-1002.
- Hernandez - Fernandez J, Ramirez NL, Fuentes SL, and Jimenez J (2010). Molecular and biological characterization of native *Bacillus thuringiensis* strain for controlling tomat leafminer (*Tuta absoluta* Meyrick) (Lepidoptera: Gelechiidae) in Colombia. *World J. Microbiol. Biotechnol.* 23:579- 590.
- Ikrum MB, Abdallah B (2011). The entomopathogenic fungus *Beauveria* sp., a promising way to fight against *Tutaabsoluta*Meyrick in Algeria". In: International symposium on management of *Tuta absoluta* (tomato borer) proceeding. Agadir, Morocco, pp.16-18.
- Illakwahhi TD, Srivastava LBB (2017). Control and management of tomato leafminer, *Tuta absoluta* (Meyrick) (Lepidoptera, Gelechiidae).A review. *OSR-JAC*, 10:4-22.
- Isman BM, Seffrin R (2014). Natural Insecticides from the Annonaceae: A Unique Example for Developing Biopesticides. *Adv.Plant Bioprocess.* 67:401-413.
- Jung HW (2009). Priming in systemic plant immunity. *Sci.* 324: 89–91.
- Karadjova O, Ilieva Z, Krumov V, Petrova E, Ventsislavov V (2013) *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae): Potential for entry, establishment and spread in Bulgaria. *Bulg. J. Agri. Sci.*19: 563–571.
- Katoch R, Sethi A, Thakur N, Murdock LL (2013). RNAi for insect control: current perspective and future challenges. *Appl. Biochem.Biotech.* 171:847-873.
- Kempal A (2009). Symbiotic soil microorganisms as players in aboveground plant-herbivore interactions -the role of rhizobia. *Oikos* 118:634-640.
- Knipling EF (1955). Possibilities in insect control or eradication through the use of sexually sterile males. *J. Econ. Entomol.* 48:459-462.
- Koch A, Kogel KH (2014). New wind in the sails: improving the agronomic value of crop plants through RNAi-mediated gene silencing. *J. Plant Biotechnol.*12:821-831.
- Kola VSR, Renuka P, Madhav MS, Mangrauthia SK (2015). Key enzymes and proteins of crop insects as a candidate for RNAi based gene silencing. *Front. Physiol.* 6:119-125.
- Kona MEN, Taha KA, Mahmoud EEM (2014). Effects of Botanical Extracts of Neem (*Azadirachta indica*) and *Jatropha* (*Jatropha curcus*) on Eggs and Larvae of Tomato Leaf Miner, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *Persian Gulf Crop Protect.* 3:41-46.
- Korycinska A, Moran H (2009). South American tomato moth (*Tuta absoluta*): plant pest factsheet. Sand Hutton, York, UK: FERA.
- Kumar GA, Jalali SK, Venkatesan T, Stouthamer R, Niranjana P, Lalitha Y (2009). Internal transcribed spacer-2 restriction fragment length polymorphism (ITS-2-RFLP) tool to differentiate some exotic and indigenous trichogrammati egg parasitoids in India. *Biol. Control*, 49:207-213.
- Leckie BM, Stewart JN (2010). Agro-infiltration as a technique for rapid assays for evaluating candidate insect resistance trans-genes in plants. *Plant Cell Reps* 30:325-334.

- Leite GLD, Picanço M, Azevedo AA, Gonring AR (1999). Efeito de tricomas, aleloquímicos nutrientes na resistência de *Lycopersicon hirsutum* à traça-do-tomateiro. *Pesq. Agropec. Bras.* 34:2059-2064.
- Leite GLD, Picanço M, Jham GN, Marquini F (2004). Intensity of *Tuta absoluta* (Meyrick, 1917) (Lepidoptera: Gelechiidae) and *Liriomyza* spp. (Diptera: Agromyzidae) attacks on *Lycopersicon esculentum* Mill. leaves. *Ciênc. Agropec.* 28:42-48.
- Lewis WJ, Vet LEM, Tumlimson JH, van Lenteren JC, Papaj DR (2003). Variations in natural enemy foraging behavior: essential element of a sound biological control theory. In: *Quality Control and Production of Biological Control Agents*. (Ed. van Lenteren JC), CABI Publishing, Wallingford (GB). pp. 41-58.
- Ley JP, Blings M, Krammer G, Reinders G, Schmidt CO, Bertram HJ, (2006). Isolation and synthesis of acmellonate, a new unsaturated long chain 2-ketol ester from *Spilanthesacmella*. *Nat. Prod. Res.* 20:798-804.
- Lietti MM, Botto E, Alzogaray RA, (2005). Insecticide resistance in Argentine populations of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *Neotrop. Entomol.* 34:113-119.
- Luan JB, Ghanim M, Czosnek H (2013). Silencing the ecdysone synthesis and signaling pathway genes disrupt nymphal development in the whitefly. *Ins. Biochem. Mol. Biol.* 43:740-746.
- Luna MG, Wada VI, Sánchez NE (2010). Biology of *Dineulophus phtorimaeae* (Hymenoptera: Eulophidae) and field interaction with *Pseudapanteles dignus* (Hymenoptera: Braconidae), larval parasitoids of *Tuta absoluta* (Lepidoptera: Gelechiidae) in tomato. *Annals Entom. Soci. Amer.* 103:936-942.
- Luna GM, Sánchez E.N, Pereyra CP, Nieves E, Savino V, Luft E, Virla E, Speranza S (2012). Biological control of *Tuta absoluta* in Argentina and Italy: evaluation of indigenous insects as natural enemies. *OEPP/EPPO Bull.* 42:260-267.
- Maluf WR Maciel MG, Gomes AAL, Cardoso GM, Goncalves DL, de Silva CE, Knapp M (2010). Broad-spectrum arthropod resistance in hybrids between high and low acylsugar tomato lines. *Crop Science*, 50:439-450.
- Mansour M (2010). Effects of gamma radiation on the Mediterranean flour moth, *Ephestia kuehniella*, eggs, and acceptability of irradiated eggs by *Trichogramma cacoeciae* females. *J. Pest Sci.* 83:243-249.
- Marec F, Tothová A, Sahara K, Traut W (2001). Meiotic pairing of sex chromosome fragments and its relation to atypical transmission of a sex-linked marker in *Ephestia kuehniella* (Insecta: Lepidoptera). *Heredity* 87: 659-671.
- Masante-Roca I, Anton S, Delbac L (2007). Attraction of the grapevine moth to host and non-host plant parts in the wind tunnel: effects of plant phenology, sex, and mating status. *Entomol. Exp. Appl.* 122:239-245.
- Mastrangelo T, Parker AG, Jessup A, Pereira R, Orozco-Da'vila D, Islam A, Dammalage T, Walder JMM (2010). A new generation of X-ray irradiators for insect sterilization. *J. Econ. Entomol.* 103:85-94.
- May RC, Plasterk RH (2005). RNA interference spreading in *C. elegans*. *Methods Enzymol.* 392:308-315.
- McCormick AC, Unsicker SB, Gershenson J (2012). The specificity of herbivore-induced plant volatiles in attracting herbivore enemies. *Trends Plant Sci.* 17:303-310.
- Medeiros NA, Burnette TD, Forscher P (2006). Myosin II functions in actin-bundle turnover in neuronal growth cones. *Nature Cell Biology*, 8:216-226.
- Megido RC, Haubruge E, Verheggen FJ (2013). Pheromone-based management strategies to control the tomato leafminer, *Tuta absoluta* (Lepidoptera: Gelechiidae). A review. *Biotechnol. Agro. Soc. Environ.* 17:475-482.
- Megido CR, De Backer L, Ettai' b R, Brostaux Y, Fauconnier LM, Delaplace P, Lognay G, Belkadi SM, Haubruge E, Francis F, Verheggen JF (2014). Role of larval host plant experience and solanaceous plant volatile emissions in *Tuta absoluta* (Lepidoptera: Gelechiidae) host finding behavior. *Arthropod-Plant Interactions* 8:293-304.
- Mehta K, Parker A (2011). Characterization and dosimetry of a practical X-ray alternative to Self-shielded gamma irradiators. *Radiat. Phys. Chem.* 80:107-113.
- Mohamed I, Mahmoud E, Elhaj M, Mohamed A, Ekesi S (2015). Host plants record for tomato leaf miner *Tuta absoluta* (Meyrick) in Sudan. *EPPO. Bull.* 45:108-111.
- Molla O, Gonzalez-Cabrera J, Urbaneja A (2011). The combined use of *Bacillus thuringiensis* and *Nesidiocoris tenuis* against the tomato borer *Tuta absoluta*. *BioControl*, 56:883-891.
- Molla O, Monton H, Vanaclocha P, Beitia F, Urbaneja A (2009). Predation by the mirids *Nesidiocoris tenuis* and *Macrolophus pygmaeus* on the tomato borer *Tuta absoluta*. *IOBC/WPRS Bull.* 49: 209-214.
- Moreno SC, Carvalho GA, Picanc MC, Morais EGF, Eriompereir R (2011). Bioactivity of compounds from *Acmella oleracea* against *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) and selectivity to two non-target species. *Pest Manag. Sci.* 8:23-32.
- Moussa A, Baiomy F, El-Adl E (2013). The status of tomato leafminer; *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in Egypt and potential effective Pesticides. *Acad. J. Entomol.* 6:110-115.
- Nelson N, Perzov N, Cohen A, Hagai K, Padler V, Nelson H (2000). The cellular biology of proton-motive force generation by V-ATPases. *J. Exper. Biol.* 203: 89-95.
- Newton PJ. (1998). False codling moth, *Cryptophlebia leucotreta* (Meyrick). In: "Citrus pests in the Republic of South Africa" (Bedford, E.C.G., van den Berg, M.A.; de Villiers, E.A., eds.). *Dynamic Ad., Nelspruit, South Africa.*
- Nilahyane A, Bouharroud R, Hormatallah A, Taadaouit NA (2012). Larvicidal effect of plant extracts on *Tuta absoluta* (Lepidoptera: Gelechiidae). "Working Group Integrated Control in Protected Crops, Mediterranean Climate". *IOBC-WRPS Bull.* 80: 305-310.
- Öztemiz S (2012). The tomato leafminer [(*Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae))] and its biological control. *KSU J. Nat. Sci.* 15: 47-57.
- Panwar V, McCallum B, Bakkeren G (2012). Endogenous silencing of *Puccinia triticina* pathogenicity genes through in planta-expressed sequences leads to the suppression of rust diseases on wheat. *Plant J.* 73:521-532.
- Pereira GVN (2008). Seleção para alto teor de acilacúcares em genótipos de tomateiro e sua relação com a resistência ao ácaro vermelho (*Tetranychus evansi*) e à traça (*Tuta absoluta*). *Ciência e Agrotecnologia*, 32:996-1004.
- Philogène BJR, Vincent C, (2005). Botanicals: yesterday's and today's promises. In: *Biopesticides of plant origin* (Regnault-Roger, C.; Philogène, B.J.R. and Vincent, C. (Ed.)), 1-15.
- Pineda A, Zheng S, van Loon AJ, Pieterse JM, Dicke M (2010). Helping plants to deal with insects: the role of beneficial soil-borne microbes. *Trends in Plant Sci.* 15:507-514.
- Polack LA, Pereyra PC, Sarandó'n SJ (2011). Control biológico de la polilla del tomate, *Tuta absoluta* (Lepidoptera: Gelechiidae) en invernáculos con flores asociadas en el noreste de la provincia de Buenos Aires, Argentina. *Seminario Internacional de Sanidad Agropecuaria. Reunión Conjunta de la International Organization for Biological Control.* Mayo 3-6, La Habana (CU) (in Spanish).
- Ponti L, Gilioli G, Biondi A, Desneux N, Gutierrez P (2015). Physiologically based demographic models streamline identification and collection of data in evidence-based pest risk assessment. *EPPO. Bull.* 45:317-322.
- Pratissoli D, Thuler RT, Andrade GS, Zanotti LCM, da Silva AF (2005). Estimate of *Trichogramma pretiosum* to control *Tuta absoluta* in stalked tomato. *Pesquisa Agropecuaria Brasileira*, 40:715-718.
- Price GRD, Gatehouse AJ (2008). RNAi-mediated crop protection against insects Review, *Trends Biotech.* 26:393-401.
- Proffitt M, Birgersson G, Bengtsson M, Reis R, Witzgall P, Lima E (2011). Attraction and oviposition of *Tuta absoluta* females in response to tomato leaf volatiles. *J. Chem. Ecol.* 37: 565-574.
- Quiroz C (1978). Utilización de trampas con hembras virgines de *Scrobipalpa absoluta* (Meyrick) (Lepidoptera: Gelechiidae) en estudios de dinámica de población. *Agric. Tech.* 38: 94-97.
- Retta NA, Berhe HD (2015). Tomato leafminer, *Tuta absoluta* (Meyrick), a devastating pest of tomatoes in the highlands of Northern Ethiopia: A call for attention and action. *Review, Res. J. Agric. Environ. Manag.* 4: 264-269.
- Reyes M, Rocha K, Alarcón L, Siegwart M, Sauphanor B (2011). Metabolic mechanisms involved in the resistance of field populations of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) to spinosad. *Pest. Biochem. Physiol.* 102:45-50.

- Reyes M, Franck P, Charmillot JP, Ioriatti C, Olivares J, Pasqualini E, Sauphanor B (2007). Diversity of insecticide resistance mechanisms and spectrum in European populations of the Codling moth, *Cydia pomonella*. *Pest Manag. Sci.* 63:890-902.
- Reyes M, Rocha K, Alarcón L, Siegwart M, Sauphanor B (2012). Metabolic mechanisms involved in the resistance of field populations of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) to spinosad. *Pestic. Biochem. Physiol.* 102:45-50.
- Rezaei AA, Talaei-Hassanlou R (2016). The Use of *Bacillus thuringiensis*-based Products in Biocontrol of Tomato Leaf Miner, *Tuta Absoluta* (Lepidoptera, Gelechiidae). *Int. J. Agric. Innov. Res.* 4:2319-1473.
- Robinet C, Roques A (2010). Direct impacts of recent climate warming on insect populations. *Integr. Zool.* 5:132-142.
- Roditakis E, Steinbach D, Moritz G, Emmanouil Vasakis E, Stavrakaki M, Ilias A, García-Vidal L, Martínez-Aguirre RM, Bielza P, Morou E, Silva EJ, Silva MW, Siqueira AH, Iqbal S, Troczka JB, Williamson SM, Bass C, Tsagkarakou A, Vontas J, Nauen R (2017). Ryanodine receptor point mutations confer diamide insecticide resistance in tomato leafminer, *Tuta absoluta* (Lepidoptera: Gelechiidae). *Ins. Bioch. Mol. Biol.* 80: 11:20.
- Rodriguez M, Bosch D, Sauphanor B, Avilla J (2010). Susceptibility to organophosphate insecticides activity of detoxifying enzymes in Spanish populations of *Cydia pomonella* (L.) (Lepidoptera: Tortricidae). *J. Econ. Entomol.* 103:482-491.
- Sabbour MM, Soliman N (2012). Evaluations of Three *Bacillus thuringiensis* Against *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in Egypt. *Int. J. Sci. Res.* 3:2067-2073.
- Sabbour MM (2014). Biocontrol of the tomato pinworm *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in Egypt. *Mid. E. J. Agric. Res.* 3:499-503.
- Salama SH, Shehata EI (2017). Bioactivity and Repellency of Some Plant Extracts Against the Tomato Leaf Miner *Tuta absoluta* (Meyrick 1917) (Lepidoptera: Gelechiidae). *RJPBCS*, 8:1021-1036.
- Sánchez NE, Pereyra PC, Luna MG (2009). Spatial patterns of parasitism of the solitary parasitoid *Pseudapanteles dignus* (Hymenoptera: Braconidae) on *Tuta absoluta* (Lepidoptera: Gelechiidae). *Environ. Entomol.* 38:365-374.
- Schoonhoven LM, Joop JA, Dicke ML (2005). *Insect-Plant Biology* 2<sup>nd</sup> edition, Oxford University Press.
- Scott IM, Jensen HR, Philogène BJ, Arnason JT (2008). A review of *Piper* spp. (Piperaceae) phytochemistry, insecticidal activity and mode of action. *Phytochem. Rev.* 7:65-75.
- Scott IM, Helson BV, Strunz GM, Finlay H, Sanchez-Vindas PE, Poveda L, Lyons B., Philogène BJR, Arnason JT (2007). Efficacy of *Piper nigrum* (Piperaceae) extract for control of insect defoliators of forest and ornamental trees. *Canad. Entomol.* 139:513-522.
- Scott IM, Jensen H, Nicol R, Lesage L, Bradbury R, Sanchez-Vindas P, Poveda L, Arnason JT, Philogène BJR (2004). Efficacy of *Piper* (Piperaceae) extracts for control of common home and garden insect pests. *J. Econ. Entomol.* 97:1390-1403.
- Segarra G, van der Ent S, Trillas I, Pietese CM (2009). MYB72, a node of convergence in induced systemic resistance triggered by a fungal and a bacterial beneficial microbe. *Plant Biol.* 11:90-96.
- Seth RK, Sharma VP (2001). Inherited sterility by substerilizing radiation in *Spodoptera litura* (Lepidoptera: Noctuidae): bioefficacy and potential for pest suppression. *Fla. Entomol.* 84:183-193.
- Shehata I, Ebada I, Ismail IA, Fouda M, Salama SH (2016). On the Population Dynamics of the Tomato Leaf Miner *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in Egypt. *Ecologia Balkanica.* 8:65-75.
- Shiberu T, Getu E (2017). Effects of crude extracts of medicinal plants in the management of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) under laboratory and glasshouse conditions in Ethiopia. *J. Entomol. Nematol.* 9:9-13.
- Silva GA, Picanco MC, Bacci L, Grespo AL, Rosado JF, Guedes RN (2011). Control failure likelihood and spatial dependence of insecticide resistance in the tomato pinworm, *Tuta absoluta*. *Pest Manag. Sci.* 67:913-920.
- Silva VF, Maluf RW, Cardoso GM, Neto GCA, Maciel MG, Nizio CAD, Silva AV (2009). Resistência mediada por aleloquímicos de enótipos de tomateiro à moscabranca e ao ácaro-rajado. *Pesquisa Agropecuária Brasileira*, 44:1262–1269.
- Soleno J, Anguiano L, de D'Angelo PA, Cichón L, Fernández D, Montagna C (2008). Toxicological biochemical response to azinphos-methyl in *Cydia pomonella* L. (Lepidoptera: Tortricidae) among orchards from the Argentinian Patagonia. *Pest Manag. Sci.* 64:964-970.
- Stein E, Molitor A, Koqel KH, Waller F (2008). Systemic resistance in *Arabidopsis* conferred by the mycorrhizal fungus *Piriformospora indica* requires jasmonic acid signaling and the cytoplasmic function of NPR1. *Plant Cell Physiol.* 49:1747-1751.
- Suckling DM, Brockerhoff EG (2010). Invasion biology, ecology, and management of the light brown apple moth (Tortricidae). *Annu. Rev. Entomol.* 55:285-306.
- Theoduloz C, Vega A, Salazar M, Gonzalez E, Meza-Basso L (2003). Expression of a *Bacillus thuringiensis* delta-endotoxin cry1Ab gene in *Bacillus subtilis* and *Bacillus licheniformis* strains that naturally colonize the phylloplane of tomato plants (*Lycopersicon esculentum*, Mills). *J. Appl. Microbiol.* 94: 375-381.
- Tonnang Z, Mohamed F, Khamis F, Ekesi S (2015). Identification and risk assessment for worldwide invasion and spread of *Tuta absoluta* with a focus on sub-saharan Africa: Impl. *Phyto Meas Manag.*
- Torres JB, Faria CA, Evangelista WS Pratisoli D (2001). Within-plant distribution of the leaf miner *Tuta absoluta* (Meyrick) immatures in processing tomatoes, with notes on plant phenology. *Int. J. Pest Manage.* 47:173-178.
- Trillas MI, Segarra G (2009). Interactions between nonpathogenic fungi and plants. *Adv. Bot. Res.* 51:321-359.
- Tumuhaise V, Khamis MF, Agona A, Sseruwu G, Mohamed AS (2016). First record of *Tuta absoluta* (Lepidoptera: Gelechiidae) in Uganda. *Int. J. Tropical Insect Sci.* 36:135-139.
- Urbaneja A, González-Cabrera J, Arnó J, Gabarra R (2012). Prospects for the biological control of *Tuta absoluta* in tomatoes of the Mediterranean basin. *Pest Manage. Sci.* 68:1215-1222.
- Urbaneja A, Vercher R, Navarro V, Garcí'a Mari' F, Porcuna L. (2007) La polilla del tomate, *Tuta absoluta*. *Phytoma Espan'a* 194:16–23.
- Van Damme V, Berkvens N, Berckmoes R Wittemans L, De Vis R, Casteels H, Tirry L, De Clercq P (2015). Overwintering potential of the invasive leafminer *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) as a pest in greenhouse tomato production in Western Europe. *J Pest Sci.* 88:533-541.
- Van der Ent S, van Hulten M, Pozo MJ, Czechowski T, Uduardi MK, Pieterse CM, Ton J (2009). Priming of plant innate immunity by rhizobacteria and beta-aminobutyric acid: Differences and similarities in regulation. *New Phytol.* 183:419-431.
- Van der Hoeven RS, Monforte AJ, Breeden D, Tanksley SD, Steffens JC, (2000). Genetic control and evolution of sesquiterpene biosynthesis in *Lycopersicon esculentum* and *L. hirsutum*. *Plant Cell* 12:2283-2294.
- Van Lenteren C, Bueno P (2003). Augmentative biological control of arthropods in Latin America. *BioCont.* 48:123-139.
- Van Oosten VR, Bodenhausen N, Reymod P, van Pelt JA, van Loon LC, Dicke M, Pieterse CM (2008). Differential effectiveness of microbially induced resistance against herbivorous insects in *Arabidopsis*. *Mol. Plant-Microbe Interact.* 21:919-930.
- Van Wees SCM, van der Ent S, Pieterse CM (2008). Plant immune responses triggered by beneficial microbes. *Curr. Opin. Plant Biol.* 11:443-448.
- Vannette RL, Hunter MD (2008). Mycorrhizal fungi as mediators of defense against insect pests in agricultural systems. *Agric. For. Entomol.* 11: 351-358.
- Vasconcelos GR (2013). Strain selection and host effect on *Trichogramma pretiosum* Riley 1879 (Hymenoptera:Trichogrammatidae) quality for *Tuta absoluta* (Meyrick, 1917) (Lepidoptera: Gelechiidae) control in tomato crops. Thesis, Univ. Moura Lacerda, Brasil.
- Visser D, Uys MV, Nieuwenhuis JR, Pieterse W (2017). First records of the tomato leafminer *Tuta absoluta* (Meyrick, 1917) (Lepidoptera: Gelechiidae) in South Africa. *BiolInvasio. Recor.* 6:1-5.
- Wang D, Gong YP, Li M, Qiu HX, Wang YK (2009). Sublethal effects of spinosad on survival, growth, and reproduction of *Helicoverpa armigera* (Lepidoptera: Noctuidae). *Pest Manag. Sci.* 65:223-227.
- Wieczorek H, Beyenbach KW, Huss MO (2009). Vacuolar-type proton

- pumps in insect epithelia. *J. Exper. Biol.* 212:161-169.
- Wu LC, Fan NC, Lin MH, Chu IR, Huang SJ, Hu CY (2008). Anti-inflammatory effect of spilanthol from *Spilanthes acmella* on murine macrophage by down-regulating LPS-induced inflammatory mediators. *J. Agric. Food Chem.* 56:2341-2349.
- Wu S, Yang Y, Yuan G, Campbell MP, Teese GM, Oakeshott GJ, Wu Y (2011). Over expressed esterases in a fenvalerate resistant strain of the cotton bollworm, *Helicoverpa armigera*. *Insect Biochem. Mol. Biol.* 41:14-21.
- Yalcin M, Mermer S, Kozaci DL, Turgut C (2015). Insecticide resistance in two populations of *Tuta absoluta* (Meyrick, 1917) (Lepidoptera: Gelechiidae) from Turkey. *Türk. Entomol. Derg.* 39:137-145.
- Yankova V, Valchev N, Markova D (2014). Effectiveness of phytopesticide Neem Azal T/S ® against tomato leaf miner (*Tuta absoluta* Meyrick) in greenhouse tomato. *Bul. J. Agric. Sci.* 20:1116-1118.
- Youssef AN, Hassan M (2013). Bioinsecticide activity of *Bacillus thuringiensis* isolates on tomato borer, *Tuta absoluta* (Meyrick) and their molecular identification. *Afr. J. Biotechnol.* 12: 3699-3709.
- Yu N, Christiaens O, Liu J, Niu J, Cappelle K, Caccia S, Huvenne H, Smaghe G (2012). Delivery of dsRNA for RNAi in insects: an overview and future directions. *In. Sci.* 20:4-14.
- Zekeya N, Ndakidemi AP, Chacha M, Mbega E (2016). Tomato Leafminer, *Tuta absoluta* (Meyrick 1917), an emerging agricultural pest in Sub-Saharan Africa: Current and prospective management strategies. *Afr. J. Agric. Res.* 12:389-396.
- Zhang H, Li HC, Miao XX (2013). Feasibility, limitation and possible solutions of RNAi based technology for insect pest control. *Insect Sci.* 20:15-30.
- Zhang HM, Xie X, Kim MS, Korneyew DA, Holaday S, Pere, PW (2008). Soil bacteria augment Arabidopsis photosynthesis by decreasing glucose sensing and abscisic acid levels in planta. *J. Plant* 56:264-273.
- Zibae I, Bandani RA, Sabahi G (2016). The expression profile of detoxifying enzyme of tomato leaf miner, *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae) to chlorpyrifos. *Arthro.* 5:77-86.