

Full Length Research Paper

Influence of Zn stresses on growth and physiology in Khus-khus (*Vetiveria zizanoides* Nash.) and its essential sesquiterpene oil(s), in relation to roots diameter circumferential positions

A. MISRA*, N. K. SRIVASTAVA and A. K. SRIVASTAVA

Central Institute of Medicinal and Aromatic Plants, P. O. CIMAP, Kukrail Picnic Spot Road Lucknow – 226015, India.

Accepted 8 November, 2011

Culturing Khus in controlled glasshouse condition in sand culture techniques for maximum essential monoterpene oil(s) was found (0.21%) in young developed middle position circumferences of roots. At middle position of leaf, net photosynthetic rate and contents of chlorophyll (Chl) were affected. The maximum peroxidase activity was obtained at the middle position of leaf and roots circumferences area, with the maximum production of biomolecule of khusimol and khusinol at 250 mg Zn/L. The maximum monoterpene oil(s) (0.21%) was also found at middle position of developed roots. However, the relative contents of khusimol and Khusinol varied at different circumference area of positions. As a result of different root developmental positions, the contents of Fe, Mn, Zn and Cu were smaller in quantity in Vetiver. Their maximum contents were observed at middle positions of developed roots. Thus, the value addition of essential monoterpene oil(s) seems to be used for commercial exploitation at large scales, to be for the collection of developed roots position at middle levels (roots area of 5th positions developments) and for better quality of total essential oil of Khus.

Key words: Chlorophyll, dry mass, leaf area, net photosynthetic rate, plant height, saccharides, Zn.

INTRODUCTION

Khus-khus (*Vetiveria zizanoides* Nash.) is an aromatic grass of the family Poaceae and it is the only source of one of the most important essential monoterpene oil(s) called the oil of Khus. It is commonly known as Khus oil. It is distinctly different from the horticultural khus, which are basically used for soil erosion and have no commercial usage in perfumery industries (Douglas, 1969). *V. zizanoides* Nash widely grown cultivars improved variety 'Gulabi' (Rajeshwar and Bhattacharya, 1992) are commonly cultivated in India. Steam distillation of root biomass of Khus-Khus yields Khusimol and Khusinol rich monoterpene Vetiver oil(s), extensively used for perfuming soaps and cosmetics and in

aromatherapy to which they impart a pronounced and lasting Vetiver oil peculiar odour. It is also largely used in flavouring soft drinks products and other pharmaceutical preparations, and also the plant is used in soil erosion as the roots are having soil binding properties. Apart from this, it is widely used in phytochelatin properties and in phytoremediation of heavy metals extraction from the heavily polluted environment.

Zn is an essential micronutrient and acts as a phytochelatin that acts either as a metal component of various enzymes or as a functional, structural, or regulatory cofactor, and is thus, associated with saccharide metabolism, photosynthesis and protein synthesis (Marschner, 1986). Zn deficiency reduces plant growth and inhibits photosynthesis in many plants including forest trees (Dell and Wilson, 1985), fiber crops (Ohki, 1986), rice (Ajay and Rathore, 1985) and spinach (Randal and Bouma, 1973). Zn retards the activity of carbon metabolism enzymes such as carbonic anhydrase

*Corresponding author. E-mail: amisracimap@yahoo.co.in or a.misra@cimap.res.in. Tel: 91 522 2342676. Fax: 91 522 2342666.

(Ohki, 1976; Ohki, 1978), ribulose 1,5-bisphosphate carboxylase/oxygenase and fructose-1,6-bisphosphate (Marschner, 1986). Zn, Se and Cr are antioxidants scavenging free radicals. Zn stimulates the removal of free radicals (Chakmak and Engles, 1999).

Essential oil biosynthesis in geranium is strongly influenced by Zn acquisition and the stresses caused by Zn on nutrition and growth. Zn is involved in carbon assimilation, saccharide accumulation, free radical removal, antioxidant enzymes, carbon utilization in terpene biosynthesis and the overall growth of the plants. The requirement of Zn for Japanese mint and its limitations imposed on photosynthetic carbon metabolism and translocation in relation to essential oil accumulation in mint were shown by Misra and Sharma (1991), whereas antioxidants enzymes for free radical quenching in geranium have not been fully documented.

In the present paper, we report on the role of Zn as a stimulant of quenching of free radicals through Zn affected antioxidant enzyme activity. Simultaneously, photosynthetic efficiency in terms of net photosynthetic rate (P_N), content of chlorophyll (Chl), leaf fresh and dry mass, leaf area, Zn content in plant shoot biomass and oil yield were also determined.

MATERIALS AND METHODS

Plant tips (12.5 to 15.0 cm) with 3 to 4 leaves of *V. zizanoides* L. genotype of diploid – Gulabi were obtained from the farm nursery of the Central Institute of Medicinal and Aromatic Plants (CIMAP), Lucknow, India. Uniform slips were initially planted in 10 000 cm³ earthen pots filled with purified silica sand (Agrawala and Sharma, 1961) for the development of roots. After 15 days, rooted cuttings were transferred to 2 500 cm³ pots. The salts used in nutrient solution of Hoagland and Arnon (1952) were purified for Zn (Hewitt, 1952). The nutrient solution was used in the experiment except Fe which was supplied as Fe- ethylenediaminetetra acetic acid (EDTA). Three pots each of Zn treatments ranging from 0.0 to 1.0 g (Zn) m⁻³ were maintained in controlled glasshouse condition at ambient temperature (30±5°C) and irradiance (800 to 1 000 $\mu\text{mol m}^{-2} \text{s}^{-1}$). The nutrient solution in each treatment was added at alternate days. With onset of deficiency and toxicity (after 20 days), growth and detailed physiological, and biochemical data characteristics were determined. P_N was measured using a computerized portable photosynthesis system *Li-COR 6000* (LiCOR, USA) (Srivastava, 1991). Chl amount in 80% acetone extracts from 3rd leaf was determined spectrophotometrically on *Pye Unicam PU8610* according to Arnon (1949). Leaf fresh and shoot dry mass, and area (area meter *Li-3000*) were also recorded. For tissue element analysis, 1 g dried leaf samples were digested with 1 M HCl at 60°C for 24 h. Aliquot samples of the clear digest were diluted with water (10 cm³) and analyzed for Zn by atomic absorption spectrophotometer (*Pye Unicam SP 2800*) (Misra and Sharma, 1991). Antioxidant reactive peroxidase enzyme activity was estimated as described in Sharon et al. (1966). Two gram (2 g) of freshly chopped leaves at 3rd position was homogenized with 5 cm³ of 0.1 M phosphate buffer (pH 6.8). Each treatment was replicated 3 times and assayed by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE).

Vetiver oil was estimated by steam distillation of 100 g freshly plucked roots area from circumferences (Root area #1 to 7, that is, outer area (ar.#1:1.5 cms, then the b following ar.2 to 7 in descending orders), in an apparatus of Clevenger (1928). Khusimol

and Khusinol, and other associated oil contents were determined by gas liquid chromatography (*Perkin-Elmer model 3920 B*). The stainless steel column was packed with 10% carbowax (20 meshes) on *Chromosorb WNAW*. Injector and detector temperature were maintained at 200°C. The flow of H₂ was 0.47 cm s⁻¹; data processing for area % was done on a *Hewlett- Packard* integrator model *HP-33*.

RESULTS AND DISCUSSION

The fresh and dry biomasses increased with increase in the supply of Zn (Table 1). Maximum fresh and dry biomass, and leaf area were observed at Zn_{0.250}. Plant height was maximum at Zn_{0.500}. Zn_{1.000} was toxic to all growth parameters. The Chl content increased up to Zn_{0.250} and then decreased. The maximum P_N was found at Zn_{0.250}; at this Zn supply also the saccharide content was the highest. Zn deficiency and toxicity inhibited P_N in cotton (Ohki, 1976), peppermint (Srivastava et al., 1997), soybean (Ohki, 1978) and sweet mint (Misra et al., 2003). A decrease in Chl content represents a decline in photochemical capacity of leaf at deficient Zn supply (Ohki, 1976).

Maxima of peroxidase activity were observed at Zn_{0.250}. The Zn deficient and toxic cultured plants revealed lesser peroxidase activity with lesser peroxidase isoenzyme band profiles. In Japanese, mint similar report was given for Mn nutrition (Misra, 1996). The maximum of monoterpene oil(s) was found at Zn_{0.250}. However, relative contents of citronellol, geraniol, linalool and nerol varied at different Zn treatments. As a result of different Zn supply, the contents of Fe, Mn, Zn and Cu were smaller in shoots of geranium. Their maximum contents were observed at Zn_{0.250}. Statistical analysis showed a positive significant association between Zn content in leaf and P_N (γ , 0.924 $\leq p$, 0.5%) and between P_N and content of saccharides (γ , 0.879 $\leq p$, 0.05%). However, Zn content in leaf was negatively correlated with Chl *a/b* ratio. P_N showed a positive significant association with leaf fresh mass (γ , 791 $\leq p$, 0.05%), leaf dry mass (γ , 692 $\leq p$, 0.05%), leaf area and total monoterpene oil(s) (γ , 0.721 $\leq p$, 0.01). A positive significant correlation was also observed between saccharides and total oil (γ , 0.695 $\leq p$, 0.01%). A quadratic trend was observed for all these characters which were comparable in +Zn than in plants grown at Zn deficit or much higher Zn supply.

We found that optimum supply of Zn is Zn_{0.250}. Utilization of metabolites from primary photosynthetic process in secondary metabolism regulates monoterpene production (Gershezon and Croteau, 1991). Thus, a close relation between photosynthesis, photorespiration and terpenoid synthesis exists in essential monoterpene oil(s) bearing plants (Maffei and Codignola, 1990). Moreover, the actively growing leaves require a larger supply of an antioxidants stimulator Zn, in association with greater supply of photosynthates. Since essential oil biosynthesis occurs in these rapidly growing leaves, the initial growth period would require a greater supply of photosynthates

Table 1. Effect of root positions on parameters of *V. zizanioides*.

Growth attribute	Root ar. #1	Root ar. #2	Root ar. #3	Root ar. #4	Root ar. #5	Root ar. #6	Root ar. #7	LSD	LSD
	0.00	0.01	0.100	0.200	0.250	1.0	1.5 mg Zn/L	At 5%	at 1%
Plant height [cm]	57.0	58.0	61.0*	62.5**	63.4**	64.1**	59.0	2.5	4.1
No. of branches	9	10*	13**	18**	10*	10*	8	1.1	3.2
Fresh mass [g plant ⁻¹]	218.8	238.6*	224.8	252.1**	282.5**	215.5**	196.2	11.1	16.3
Dry mass [g plant ⁻¹]	14.11	16.33*	16.81*	17.37**	19.36**	18.46**	15.85	2.10	3.30
Leaf area [cm ²]	8.2	12.1*	25.2**	39.1**	40.3**	37.2**	11.2	3.5	6.2
Chl a [g kg ⁻¹ (FM)]	0.68	0.79*	0.94**	1.35**	1.48**	1.01**	0.82*	0.11	0.15
Chl b [g kg ⁻¹ (FM)]	0.50	0.56	0.61*	0.69**	0.79**	0.40	0.29	0.08	0.12
Chl a/b	1.36	1.41	1.54	1.96	1.87	2.53	2.83	-	-
P _N [µg(CO ₂) m ⁻² s ⁻¹]	0.15	0.19*	0.75**	0.76**	0.82**	0.71**	0.42**	0.03	0.06
Saccharides [µg (CH ₂ O) m ⁻² s ⁻¹]	0.102	0.129	0.510	0.516	0.558	0.483	0.286	-	-
Oil %	0.15	0.16	0.17*	0.19	0.21**	0.16	0.15	0.02	0.04
Khusimol [% of total oil]	0.21	0.27**	0.29**	0.32**	0.25**	0.18**	0.17**	0.01	0.02
Khusinol [% of total oil]	0.09	0.09	0.10**	0.11**	0.07**	0.12**	0.10**	0.01	0.01
Roots tissue concentrations									
Fe [mg kg ⁻¹]	98	112	142**	249**	537**	419**	312**	21	42
Mn [mg kg ⁻¹]	26	37**	41**	57**	98**	62**	53**	9	11
Zn [mg kg ⁻¹]	12	19*	34**	45**	64**	41**	36**	7	9
Cu [mg kg ⁻¹]	7	9	11**	11	12**	7	5	3	5

Chl: Chlorophyll; P_N, net photosynthetic rate; oil amounts in % of total oil. *, ** Values are significant at P, 0.05 or P, 0.01 levels, respectively.

and energy.

Conclusion

The Zn at 0.250 mg/L application and at # 5 diameter circumference position has given the maximum total oil % as well as the maximum khusimol and khusinol oil contents. Therefore, 0.250 Zn mg/L application is more useful for better growth of Khus-khus and maximum oil production.

REFERENCES

- Agrawala SC, Sharma CP (1961). The standardization of sand and water culture technique for the study of macro and micronutrients (trace) element deficiencies under Indian conditions. *Curr. Sci.*, 40: 424-428.
- Ajay, Rathore VS (1985). Effect of Zn²⁺ stress in rice (*Oryza sativa* cv. Manhar) on growth and photosynthetic processes. *Photosynthetica*, 31: 571-584.
- Arnon DI (1949). Copper enzymes in isolated chloroplasts. *Polyphenoloxidase in Beta vulgaris*. *Plant Physiol.*, 24: 1-15.
- Chakmak I, Engels C (1999). Role of mineral nutrients in photosynthesis and yield formation – In: Ringel, Z. (ed.): *Mineral Nutrition of Crops*. Haworth Press, New York, pp. 141-168.
- Clevenger JF (1928). Apparatus for determination of essential oils. *J. Am. Pharm. Assoc.*, 17: 346.
- Dell B, Wilson SA (1985). Effect of Zn supply on growth of three species of Eucalyptus seedlings and wheat. *Plant Soil*, 88: 377-384.
- Douglas JS (1969). Essential oil crops and their uses. *World Crops*, 21: 49-54.
- Hewitt EJ (1952). Sand and water culture methods used in the study of plant nutrition. *Commonwealth Bureau bot. Plantation Crops Tech. Commun.*, 22: 405-439.
- Hoagland DR, Arnon DI (1952). The water culture method for growing plants without soil. *Calif. Agr. Exp. Stat. Circ.*, 347: 1-32.
- Maffei M, Codignola A (1990). Photosynthesis, photorespiration and herbicide effect on terpene production in peppermint (*Mentha piperita* L.). *J. Essent. Oil Res.*, 2: 275-286.
- Marschner H (1986). Function of mineral nutrients. *Micronutrients*. – In: *Mineral Nutrition of Higher Plants*. Academic Press, New York, pp. 269-300.
- Misra A (1996). Genotypic variation of manganese toxicity and tolerance of Japanese mint. *J. Herbs Spices Medic. Plants*, 4: 3-13.
- Misra A, Sharma S (1991). Zn concentration for essential oil yield and menthol concentration of Japanese mint. *Fertilizer Res.*, 29: 261-265.
- Misra A, Srivastava NK, Sharma S (2003). Role of an

- antioxidant on net photosynthetic rate, carbon partitioning and oil accumulation in sweet mint. *Agrotropica*, 15: 59-74.
- Ohki K (1976). Effect of zinc nutrition on photosynthesis and carbonic anhydrase activity in cotton. *Physiol. Plant*, 38: 300-304.
- Ohki K (1978). Zinc concentration in soybean as related to growth, photosynthesis, and carbonic anhydrase activity. *Crop Sci.*, 18: 79-82.
- Rajeshwar-Rao BR, Bhattacharya AK (1992). History and botanical nomenclature of rose scented geranium cultivars grown in India. – *Indian Perfumer*, 36: 155-161.
- Sharon LM, Kay E, Lew JY (1966). Peroxidase isoenzymes from horse radish roots 1-1 Isolation and physical properties. *J. Biol. Chem.*, 241: 2166-2172.
- Srivastava NK, Misra A (1991). Effect of the triacontanol formulation "Miraculon" on photosynthesis, growth, nutrient uptake and essential oil yield of lemon grass (*Cymbopogon flexuosus steud*, Watts). *Plant Growth Regul.*, 10: 57-63.
- Srivastava NK, Misra A, Sharma S (1997). Effect of Zn deficiency on net photosynthetic rate, ¹⁴C partitioning, and oil accumulation in leaves of peppermint. *Photosynthetica*, 33: 71-79.