

Full Length Research Paper

Variation in icariin and flavonoid contents of Barrenwort species

Qiu-Mei Quan¹, Wei Wu¹, Yun-Xiang Li^{1,2*} and Qian-Rong Cai¹

¹Sichuan Agricultural University, Yaan, Sichuan 625014, China.

²College of Life Sciences, China West Normal University, Nanchong, Sichuan 637002, China.

Accepted 27 May, 2010

Flavonoid and icariin contents of five Barrenwort species native to Sichuan, China, of the genus *Epimedium* were determined by HPLC and UV. The icariin and flavonoid contents of the root, stem and leaf were differentiated significantly among five Barrenwort species. The contents showed the same trend in different tissues of the five Barrenwort species and differed from each other, that's leaf > root > stem. Although, only the leaf of five Barrenwort species meet the standards, the root and stem still can be utilized for herbalist and manufacturing medicines. The icariin content extracted from long-term storage Barrenwort was little available, but not flavonoid contents.

Key words: Barrenwort, icariin, flavonoid, Sichuan.

INTRODUCTION

Barrenwort is a perennial plant of the genus *Epimedium* L. (Berberidaceae) and is a popular botanical supplement used to improve menopausal symptoms and bone health (Shen et al., 2008). Internationally more than 60 species are known as barrenwort, whereas about 50 are detected in China (CCP, 2005). Most of the Barrenwort grow in the southwestern China. Sichuan is one of the most important centers of diversity of the genus *Epimedium*. Medicinal composition of this plant is effective in strengthening the functional of normal kidney and curing the rheumatism. It is widely used in the treatment of osteoporosis, hypertension, coronary heart disease (Guo, 1996), to strengthen immunity, prevent caducity and cure cancer (Xue et al., 2005; Zhang et al., 2002). It is already known that the most prominent medicinal compounds in the genus are flavonoids and icariin which are used to monitor quality of Barrenwort as defined by the Committee of China Pharmacopoeia (CCP, 2005). Icariin and flavones were reported to enhance the osteogenic differentiation of rat primary bone marrow stromal cells

(Chen et al., 2005), increase osteoblastic proliferation (Meng et al., 2005), reduced osteoclastic bone resorption (Yu et al., 1999) and to increase mineral content and prevent osteoporosis in ovariectomized rats (Zhang et al., 2006; Shen et al., 2008).

Many Barrenwort species are used in Chinese medicine (Zhang et al., 1995; Xie and Sun, 2006). They can vary in their icariin and flavonoid contents and because raw and processed materials can be mixed from different species, the quality of medicinal preparations is not assured (Sheng et al., 2008). So far, most of the research on *Epimedium* has focused on extracting and identifying novel compounds. Li and Liu (1988) described the isolation and chemical characterization of novel flavonoid compounds from *Epimedium davidii*. Chen et al. (2007) studied Simultaneous determination of 15 flavonoids in *Epimedium brevicornu*, *Epimedium wushanense* and *E. davidii* using pressurized liquid extraction and high-performance liquid chromatography. And Wu et al. (2008) reported simultaneous determination of seven flavonoids in *E. brevicornu*, *E. wushanense* and *E. davidii* by liquid chromatography-tandem mass spectrometry method.

Zhang et al. (2008) studied simultaneous extraction of epimedin A, B, C and icariin from *E. brevicornu*,

*Corresponding author. E-mail: yx_li@263.net. Fax: +86-0817-2314311.

Table 1. Five species of Barrenwort used in the study.

Species	Native to	Plant number	Sample	Symbol
<i>E. brevicornu</i>	Nanchong, Sichuan	10	Root	BRr
			Stem	BRs
			leaf	BRI
<i>E. wushanense</i>	Nanchong, Sichuan	7	Root	SAr
			Stem	SAs
			leaf	SAI
<i>E. davidii</i>	Baoxing, Sichuan	12	Root	DAr
			Stem	DAs
			leaf	DAI
<i>E. chlorandrum</i>	Baoxing, Sichuan	6	Root	CHr
			Stem	CHs
			leaf	CHI
<i>E. flavum</i>	Tianquan, Sichuan	11	Root	FLr
			Stem	FLs
			leaf	FLI

E. wushanense and *Epimedium flavum* by ultrasonic technique. However, there are few reports on the variation in flavonoid and icariin contents of many *Epimedium* species native to Sichuan, China. And according to the Chinese Pharmacopoeia (CCP, 2005), many researches used the dried aerial parts (leaf) of *Epimedium* (family Berberidaceae), but underground parts of *Epimedium* were usually collected as medicinal materials by herbalist. So, this paper describes flavonoid and icariin contents of the dried aerial (leaf, stem) and underground parts (root) of five Barrenwort species native to Sichuan, China, using high-performance liquid chromatography (HPLC) and ultraviolet (UV). This can provides some clues on the quality of whole-plant, rational use of wild *Epimedium* resource and directing the cultivation.

MATERIALS AND METHODS

We collected five Barrenwort species from field and were divided into root, stem, leaf, respectively (Table 1). The researchers obtained 90 samples from Barrenwort species.

Determination of icariin and flavonoid contents

Icariin and flavonoid contents were estimated by following the standard methods (CCP, 2005) recommended for studies using HPLC and UV. All analyses of icariin were performed on an Agilent Series 1100 (Agilent Technologies, USA) LC/MSD Trap system, equipped with a vacuum degasser, a quaternary pump, an autosampler, a column compartment, a diode-array detector and an ion-trap mass spectrometer with electrospray ionization interface,

controlled by Agilent LC/MSD Trap Software. A Zorbax SB-C18 column (250 × 4.6 mm I.D., 5 μm) was used. The mobile phase consisted of acetonitrile and water (70:30). The flow rate was 1.0 ml/min and the injection volume was 10 μl. The column temperature was maintained at 30°C. The analytes were monitored at 270 nm. Flavonoid are test by UV-2450 (SHIMADZU, Japan), the analytes also were monitored at 270 nm.

Statistical analyzes

All data were analyzed using one way ANOVA with Duncan analysis as a posterior test and Nonparametric test with SPSS11.0 software.


RESULTS

Variation in icariin content and flavonoid content among five Barrenwort species

Nonparametric-test was used for pooled data on icariin and flavonoid contents among five Barrenwort species (Table 2). Table 2 shows that the icariin and flavonoid contents of the root, stem and leaf differed significantly among five Barrenwort species (p < 0.05). Firstly, the icariin content of BRr is much higher than other four and the CHr have the lowest icariin content. While the flavonoid contents of CHr is much more than other four, the flavonoid contents of DAr is lowest among them. Secondly, DAs contained the greatest amount both in icariin and flavonoid content, whereas, SAs and BRs contained less in icariin and flavonoid content, respectively. Lastly, DAI recorded the highest icariin and

Table 2. Variation in icariin content and flavonoid content among five Barrenwort species.

Sample	N	Icariin content Mean \pm S.E (%)	Chi-Square	Sig.	Flavonoid content Mean \pm S. E (%)	Chi-Square	Sig.
BRr	10	0.243 \pm 0.008			4.643 \pm 0.077		
SAr	7	0.072 \pm 0.002			4.529 \pm 0.067		
DAr	12	0.169 \pm 0.011	12.833	0.012	4.133 \pm 0.199	10.333	0.035
CHr	6	0.047 \pm 0.001			4.711 \pm 0.003		
FLr	11	0.174 \pm 0.018			4.130 \pm 0.125		
BRs	10	0.036 \pm 0.001			1.014 \pm 0.021		
SAs	7	0.013 \pm 0.001			1.260 \pm 0.070		
DAs	12	0.202 \pm 0.030	13.5	0.009	3.028 \pm 0.210	13.233	0.01
CHs	6	0.056 \pm 0.005			1.538 \pm 0.091		
FLs	11	0.091 \pm 0.012			2.541 \pm 0.081		
BRI	10	0.350 \pm 0.004			5.461 \pm 0.064		
SAI	7	0.146 \pm 0.005			6.067 \pm 0.414		
DAI	12	0.557 \pm 0.002	13.5	0.009	8.639 \pm 0.169	13.033	0.011
CHI	6	0.220 \pm 0.017			5.372 \pm 0.127		
FLI	11	1.030 \pm 0.021			9.949 \pm 0.443		

**Figure 1.** Variation in icariin about aerial and underground parts of five Barrenwort species (The bars indicate mean \pm SE, and the different letters indicate significant differences according to the test at $p < 0.05$).

flavonoid content, while, the SAI and CHI recorded the lowest icariin and flavonoid content, respectively.

Variation in icariin and flavonoid contents about aerial and underground parts of five Barrenwort species

The change trends of the icariin (Figure 1) and flavonoid (Figure 2) contents about aerial and underground parts were the same in the five Barrenwort species, that's leaf > root > stem and one way ANOVA indicated the icariin

and flavonoid contents of leaf, root and stem were remarkably different from each other, respectively ($p < 0.01$).

Variation in flavonoid and icariin contents between different deposit time of *E. brevicornu* and *E. wushanense*

Nonparametric-test was used for pooled data on icariin and flavonoid (Figure 4) contents between different deposit time (freshly and one years later) of *E. brevicornu*




Figure 2. Variation in flavonoid about aerial and underground parts of five Barrenwort species.

and *E. wushanense*. It is clear that the fresh *E. brevicornu* and *E. wushanense* contained the greatest amount icariin contents, and are significantly higher than those stored for a years at room temperature ($p < 0.05$) (Figure 3). The average flavonoid contents of the aerial and underground parts in *E. brevicornu* and *E. wushanense* are remarkably higher than those stored for a years at room temperature ($p < 0.05$), except for BRr, BRl and SAl, which showed no significant difference between the no storage and storage ($p > 0.05$) (Figure 4).

DISCUSSION

Different Barrenwort species can vary in the icariin and flavonoid contents (Sheng et al., 2008). The authors studied five Barrenwort species native to Sichuan in China, and discovered that the icariin and flavonoid contents of the root, stem and leaf differed significantly among five Barrenwort species. The BRr, DAs and DAi contained the greatest amount in icariin content and flavonoid contents were rich in the CHr, DAs and DAi. The icariin and flavonoid contents about aerial and underground parts showed the same trend in the five Barrenwort species and significantly differed from each other, that's leaf > root> stem. Based on the standard laid down by the Chinese Pharmacopoeia (2005), it requires

that flavonoid content as determined by UV is more than 5.0% and icariin content as determined by HPLC is more than 0.5%. The authors found that, only the leaf of the five Barrenwort species meet the standards, whereas, the icariin and flavonoid contents of root and stem do not meet the standards. But both the root and stem, especially the root still can be commercially utilized for herbalist and manufacturing medicines.

Five species of *Epimedium*, namely *Epimedium wushanense*, *E. wushanense*, *Epimedium koreanum* Nakai, *Epimedium pubescens* Maxim and *E. brevicornu* Maxim were embodied in the medicinal species of this genus according to the Chinese Pharmacopoeia. Because of their higher flavonoid and icariin contents, abundance in nature (Takahashi, 1989) and extensive distribution (Wang et al., 2001; Zhang et al., 2002), resulting in great scarcity of naturally occurring Chinese medicinal plants (Lubell and Brand, 2005; Xie and Sun, 2006). The study indicated that *E. davidii* and *E. flavum* are richer in flavonoids and icariin than *E. wushanense* and *E. brevicornu* in the leaf. So, there exists some high-quality *Epimedium* of the species that can meet the standards even when not recorded by the Chinese Pharmacopoeia. These wild *E. davidii* and *E. flavum* may prove to possess great medicinal value.

Although, pure icariin is very stable (Li et al., 1996), the authors found highly significant differences between the




Figure 3. Variation in icariin contents in two treatments of *E. brevicornu* and *E. wushanense*.




Figure 4. Variation in flavonoid contents in two treatments of *E. brevicornu* and *E. wushanense*.

icariin content of freshly prepared samples and those stored for one year at room temperature. The result was analogous to *Epimedium acuminatum* (Sheng et al., 2008). But the icariin of *E. acuminatum* decomposition was faster than *E. brevicornu* and *E. wushanense*. This may be caused by the different materials and the icariin of *E. acuminatum* stored in the solvent, while *E. brevicornu* and *E. wushanense* were crushed during storage. Flavonoid content of the leave in *E. brevicornu* and *E. wushanense* showed no significant difference between freshly prepared samples and those stored. The study suggested that flavonoid content Barrenwort could be stored in a long-term.

ACKNOWLEDGMENTS

We are very grateful to Prof. Qing-Fu Chen, Dr. Ai-Min Shi, Shen Hao, Wen-Ting Yang for the instruction and help. We also thank the support from Qian Liu and Xiao-Hong Ye.

REFERENCES

- Chen KM, Ge BF, Ma HP, Liu XY, Bai MH, Wang Y (2005). Icariin, a flavonoid from the herb *Epimedium* enhances the osteogenic differentiation of rat primary bone marrow stromal cells. *Pharmazie* 60: 939-942.
- Chen X, Guo BL, Li SP, Zhang QW, Tub PF, Wang YT (2007). Simultaneous determination of 15 flavonoids in *Epimedium* using pressurized liquid extraction and high-performance liquid chromatography. *J. Chromatograph. A* 1163: 96-104.
- Guo BL (1996). A Study of Phylogenesis of *Epimedium* L. Institute of Medicinal Plant Development, Chinese Academy of Medical Science, Beijing 36(9): 719-723.
- Li F, Liu YL (1988). Studies on the isolation and structure of baohuoside-1, -VI and -VII and baohuosu. *Acta Pharmaceutica Sinica* 23: 739-748.
- Li WK, Zhang RY, Xiao PG (1996). Flavonoids from *Epimedium wushanense*. *Phytochemistry* 43(2): 527-530.
- Lubell JD, Brand MH (2005). Division size and timing influence propagation of four species of *Epimedium* L. *Hortscience* 40(5): 1444-1447.
- Meng FH, Li YB, Xiong ZL, Jiang ZM, Li FM (2005). Osteoblastic proliferative activity of *Epimedium brevicornu* Maxim. *Phytomedicine* 12: 189-193.
- Shen P, Guo BL, Gong Y, Hong DYQ, Hong Y, Yong EL (2007). Taxonomic, genetic, chemical and estrogenic characteristics of *Epimedium* species. *Phytochemistry* 68: 1448-1458.
- Takahashi C (1989). Karyomorphological studies on speciation of *Epimedium* and its allied *Vancouveria* with special reference to C-bands. *J. Sci. Hiroshima Univers. Ser. B. Div. 2 (Botany)* 22(2): 159-269.
- Wu CS, Guo BL, Sheng YX, Zhang JL (2008). Simultaneous determination of seven flavonoids in *Epimedium* by liquid chromatography-tandem mass spectrometry method. *Chinese Chem. Lett.* 19: 329-332.
- Wang T, Su YJ, Zhu JM (2001). RAPD analysis on some species of *Berberidaceae*. *Bull. Bot. Res.* 21(3): 428-431.
- Xie JP, Sun WJ (2006). Progress of chemical materials and pharmacy of genus *Epimedium* plants. *Strait Pharm. J.* 18(5): 17-20.
- Xue Y, Wang P, Qi QH (2005). The experimental study of the effects of icariin on increasing Smad4 mRNA level in MC3T3-E1 cell *in vitro*. *Chin. J. Orthopaedics* 25(2): 119-124.
- Yu S, Chen K, Li S, Zhang K (1999). *In vitro* and *in vivo* studies of the effect of a Chinese herb medicine on osteoclastic bone resorption. *Chin. J. Dent. Res.* 2: 7-11.
- Zhang L, Wang Y, Mao HT (2002). Study on the inhibition of telomerase activity and regulated mechanism in human cancer cell by icariin. *Chinese J. Immunol.* 18 (3): 191-196.
- Zhang G, Qin L, Hung WY (2006). Flavonoids derived from herbal *Epimedium brevicornu* Maxim prevent OVX-induced osteoporosis in rats independent of its enhancement in intestinal calcium absorption. *Bone* (January 12) [Epub] PMID: 16413840.
- Zhang Y, Xiao CH, Meng XL (1995). Flavonoids contents determined and resources utilized of 8 species for sale native to Sichuan. *China J. Chinese Mat. Med.* 20(4): 201-202.
- Zhang HF, Yang TS, Li ZZ, Wang Y (2008). Simultaneous extraction of epimedin A, B, C and icariin from *E. brevicornu*, *E. wushanense*, *E. flavum* by ultrasonic technique. *Ultrasonics Sonochemistry* 15: 376-385.