

Full Length Research Paper

# Antioxidative properties of leaf extracts of a popular Malaysian herb, *Labisia pumila*

Mohamad Norhaiza, Mahmood Maziah\* and Mansor Hakiman

Department of Biochemistry, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia 43400 UPM Serdang, Selangor Darul Ehsan, Malaysia.

Accepted 15 March, 2009

A study was undertaken to examine the presence of antioxidative activities of two varieties of *Labisia pumila*; *L. pumila* var. *Alata* and *L. pumila* var. *Pumila* using DPPH, FRAP and  $\beta$ -carotene bleaching methods. In addition, ascorbic acid,  $\beta$ -carotene, anthocyanin, total flavonoid and total phenolic content were also analyzed. In eight methods studied, six of them showed high activities of antioxidant in *L. pumila* var. *Alata* compared to that of *L. pumila* var. *Pumila*. The results obtained showed that *L. pumila* var. *Alata* contained higher antioxidative activities in all three methods applied compared to var. *Pumila*. In DPPH, FRAP and  $\beta$ -carotene bleaching methods, *L. pumila* var. *Alata* had high antioxidant activities with 299.84  $\mu$ M trolox/g db, 164.16  $\mu$ M trolox/g db and 89.22%, respectively. The same pattern of antioxidant activities also can be observed in ascorbic acid,  $\beta$ -carotene and anthocyanin in *L. pumila* var. *Alata* compared to var. *Pumila* with 0.022, 3.175 and 0.328 mg/g FW, respectively. *L. pumila* var. *Pumila* had higher total flavonoid content than *L. pumila* var. *Alata* with 1.281 mg/g FW. For total phenolic content, no significant different was observed because the amount of total phenolic content ranging from 2.53 to 2.55 mg/g FW. There is a positive correlation between antioxidant capacities and individual antioxidative compounds in the following order  $\beta$ -carotene>flavonoid>vitamin C>total anthocyanins >phenolics.

**Keywords:** *Labisia pumila*, antioxidants,  $\beta$ -carotene, flavonoid, vitamin C, anthocyanin, phenolics.

## INTRODUCTION

*Labisia pumila* is a popular herb from the family Myrsinaceae locally known as Kacip Fatimah used to induce and facilitate childbirth as well as a post partum medication to help contract the birth channel, to tone the abdominal muscles and to regain body strength (Wan Ezumi et al., 2007). Other traditional uses include treatment for dysentery, rheumatism and gonorrhoea. *L. pumila* products are widely available commercially as health supplements. In spite of the wide usage of this plant as a traditional herb no information on its chemical constituents has been documented. However, it was reported that plants from the same genus (Myrsinaceae) are being used to treat respiratory tract infections and menstrual disorders (Huang et al., 2000). Many herbal plants are known to contain high bioactive compounds such as saponins, steroids and flavonoids. Results of previous studies (Huang

et al., 2003; Jansakul et al., 1987; De Tommasi et al., 1993) have shown that steroid and triterpenoid saponins have been isolated from *Ardisia* plants (Myrsinaceae). Several studies on the antioxidative activities properties from herbal plants have been reported (Djeridane et al., 2006; Chanwitheesuk et al., 2005; Saha et al., 2004; Velioglu et al., 1998). Plant antioxidants are believed to play a role in protection against a variety of diseases and delaying ageing processes. The health promoting effect of antioxidants from plants could be due to their protective effects by counter acting reactive oxygen species (ROS) (Wong et al., 2006). There are several compounds which contribute to the antioxidative properties, these include polyphenols (Marinova et al., 2005), vitamin C (Chanwitheesuk et al., 2005),  $\beta$ -carotene (Lisiewska et al., 2006), anthocyanins (Longo and Vasapollo, 2006) and flavonoids (Harborne and Williams, 2000). There are three varieties of *L. pumila* namely, *L. pumila* var. *Pumila*, *L. pumila* var. *Alata* and *L. pumila* var. *Lanceolata*. However, only two varieties were available, *L. pumila* var. *Pu-*

\*Corresponding author. E-mail: [maziahm@biotech.upm.edu.my](mailto:maziahm@biotech.upm.edu.my).  
Phone: +603 89466703. Fax: +603 89430913.

*mila* and *L. pumila* var. *Alata*. The antioxidative capacities of *L. pumila* plants have not been reported. A study was, therefore, undertaken to examine the presence of antioxidative activities of *L. pumila*. The information will be useful in preparing the herbal formulations for health supplements.

## MATERIALS AND METHODS

### Chemicals and reagents

Gallic acid, Folin-Ciocalteu phenol reagent, 2, 2-diphenyl-1-picrylhydrazyl radical (DPPH), 2, 4, 6 - tri (2-pyridyl) 2, s-trizine (TPTZ) and kaemferol were purchased from Sigma Chemical Co. Ltd (USA). The 2, 6-dichlorophenolindophenol disodium salt was obtained from Sigma (USA). All other chemicals and solvents used in this study were of analytical grades.

### Plant materials

Two varieties of *L. pumila*, var. *Pumila* and var. *Alata* were used. The first fully expanded leaves from one-year old plants, grown in the greenhouse, were sampled. Leaf samples were dried in a convection oven at 45°C for 72 h or until there is no change in weight. The samples were placed in air tight plastic bags and kept in the refrigerator (-20 °C) until ready to be analyzed.

### Extraction method for antioxidant activity

For DPPH and FRAP assay, 0.2 g of fresh weight were extracted in 10 ml of double distilled water. The mixture was allowed to stand at room temperature for 1 h in the dark with occasional agitation. The aqueous extract was obtained by filtering the mixture through Whatman No. 1 filter paper and used for analysis. Ground sample (1 g) was extracted for  $\beta$ -carotene bleaching assay, 25 ml of 80 % methanol on orbital shaker for 120 mins at room temperature. The mixture was filtered using Whatman No. 1 filter paper and the filtrate was assayed for antioxidant activity.

### 1, 1-diphenyl-2-picrylhydrazyl (DPPH) free radical scavenging activity

The DPPH free radical scavenging activity of each sample was determined according to the method described by Wong et al. (2006). A solution of 0.1 mM DPPH in methanol was prepared. The initial absorbance of the DPPH in methanol was measured at 515 nm. An aliquot (40  $\mu$ l) of an extract was added to 3 ml of methanolic DPPH solution. The change in absorbance at 515 nm was measured after 30 mins. The antioxidant capacity based on the DPPH free radical scavenging ability of the extract was expressed as  $\mu$ mole Trolox equivalent per gram of dried plant material.

### Ferric reducing antioxidant potential (FRAP) assay

The ability to reduce ferric ions was measured using modifying methods of Wong et al. (2006), Benzie and Strain (1996). An aliquot (200  $\mu$ l) of the extract with appropriate dilution was added to 3 ml of FRAP reagent (10 parts of 300 mM sodium acetate buffer at pH 3.6, 1 part of 10 mM TPTZ solution and 1 part of 20 mM  $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$  solution) and the reaction mixture was incubated in a water bath at 37°C. The increase in absorbance at 593 nm was measured after 30 mins. The antioxidant capacity based on the ability to reduce ferric ions of the extract was expressed as  $\mu$ mole Trolox equivalents per gram of plant material based on dry weight.

### $\beta$ -Carotene bleaching assay

The  $\beta$ -carotene bleaching method (Velioglu et al., 1998) was carried out to measure the antioxidant activity.  $\beta$ -carotene (0.2 mg/ml) dissolved in chloroform, was added to round bottom flasks (50 ml) containing 20  $\mu$ l of linoleic acid and 200  $\mu$ l of Tween 20. A volume of 200  $\mu$ L of 80% MeOH (as control) or corresponding plant extract or BHT (as standard) was added to the mixture. The concentrations of the BHT and plant extract are same (40 mg/ml). After evaporation to dryness under vacuum at room temperature, oxygenated distilled water (50 ml) was added and the mixture was shaken to form a liposome solution. The samples were then subjected to thermal autoxidation at 50°C for 2 h. The absorbance of the solution was monitored at 470 nm by taking measurements at 10 mins intervals for 120 min and the rate of bleaching of  $\beta$ -carotene was recorded.

### Ascorbic acid content

The ascorbic acid content was measured using a modified method of Davis and Masten (1991). The leaf samples were extracted in 1% of phosphate-citrate buffer, pH 3.5 using chilled pestle and mortar. The homogenate was filtered. The filtrate was added to the 1 ml of 1.7 mM 2, 6-dichloroindophenol (2, 6-DCPIP) in 3 ml cuvette. The absorbance at 520 nm was read within 10 mins of mixing the reagents. The extraction buffer was used as a blank.

### $\beta$ -Carotene content

$\beta$ -carotene content was determined using the method described by (Harborne, 1973). Fresh leaves were ground in the presence of cold acetone and light petroleum. The extract was applied on to Thin Layer Chromatography (TLC) plate silica gel 60  $F_{254}$  plate (20x20 cm). The plate was developed with hexane: acetone (1:1). The spot was scraped and diluted back to the extraction solvent. The absorbance was read at 451 nm. The amount of  $\beta$ -carotene was determined as mg of  $\beta$ -carotene equivalent by using an equation obtained from the standard curve of  $\beta$ -carotene (Merck, Darmstadt, Germany).

### Anthocyanin content


Anthocyanin content was determined according to Bharti and Khurana (2003). Fresh leaves were added in 10 ml acidic methanol (1% v/v HCl) and incubate overnight. Anthocyanin was partitioned from chlorophyll with 10 ml chloroform, followed by adding 9 ml of double deionised water. The test tubes containing the samples were shaken gently and allowed the mixture to settle down. The absorbance was read at 505 nm. Petunidin was used as a standard.

### Total flavonoid content

The total flavonoid content was determined using the aluminium chloride assay (Zhishen et al., 1999). The fresh samples were added in 2 ml HCl in test tube and reflux it at 100°C for 30- 40 min. The hydrophilic extract was top up to 5 ml with distilled water. The mixture was added with 5% w/v  $\text{NaNO}_2$  and 10 %  $\text{AlCl}_3$  was added. 1 M NaOH was added a min later. Absorption reading was read at 510 nm. Kaemferol was used as the standard.

### Total phenolic content

The total phenolic content of leaves was determined by using the Folin-Ciocalteu method (Singleton and Rossi, 1965). The dried



**Figure 1.** Antioxidant activities of *L. pumila* leaf extracts based on their abilities to reduce the ferric ion-TPTZ complex (FRAP) and scavenge DPPH free radicals. Data were analyzed using One-Way ANOVA and the different contrasted using Tukey's multiple range tests. Different letter(s) indicate the values are significantly different ( $p < 0.05$ ).

sample was extracted with the 80 % methanol that contains 1 % HCl at room temperature on an orbital shaker set at 200 rpm. The mixture was centrifuged at 1000 g for 15 min. The extract was added to the 0.2 ml Follin-Ciocalteau reagent and mixed thoroughly. After 4 mins, 15 %  $\text{Na}_2\text{CO}_3$  was added. After 2 h, the absorbance was read at 760 nm with spectro uv-vis auto, LaboMed. Inc. The amount of the total phenolic content was determined as mg of gallic acid equivalent using the standard curve.

#### Statistical analysis

Data were analyzed using One-Way ANOVA and the different contrasted Tukey's multiple range tests. Different letter(s) indicate the values are significantly different ( $p < 0.05$ ).

## RESULTS


### Total antioxidant activities

DPPH, FRAP and  $\beta$ -carotene bleaching methods were carried out to determine the total antioxidative potential of two varieties of *L. pumila* leaves. The results obtained showed that *L. pumila* var. *Alata* contained higher antioxidative activities for all the methods employed.


### DPPH and FRAP assays

DPPH is a very stable free radical and it is widely used to evaluate antioxidant activities in a relatively short time (Mokbel and Hashinaga, 2006). The assay is based on the reduction of DPPH radicals in methanol which causes

an absorbance to drop at 515 nm. In this study, the antioxidant activity was expressed as Trolox equivalents per gram of plant material on a dry basis. Wong et al. (2006) reported that this method is a more meaningful and descriptive expression than the assay that used percent of the antioxidant to reduce ferric ions of the extract though both methods are expressed as  $\mu\text{mol}$  Trolox equivalents per gram of plant material on dry weight basis. The control used was butylated hydroxytoluene (BHT). The DPPH free radical scavenging activity (Figure 1) showed that *L. pumila* var. *Alata* had higher scavenging activity ( $299.84 \pm 34.05 \mu\text{M}$  trolox equivalent per g db) as compared to *L. pumila* var. *Pumila* ( $167.6 \pm 68.44 \mu\text{M}$  trolox equivalent per g db). This accounts to an increase of scavenging power by almost two-fold in *L. pumila* var. *Alata*. The FRAP values were also expressed as Trolox equivalent capacity (TEAC) of the two varieties as shown in Figure 1. A similar trend was obtained whereby the *L. pumila* var. *Alata* showed higher  $\text{TEAC}_{\text{FRAP}}$  values ( $164.16 \pm 6.89 \mu\text{M}$  trolox equivalent per g db) followed by var. *Pumila* ( $45.47 \pm 3.02 \mu\text{M}$  trolox equivalent per g db). This further indicates that *L. pumila* var. *Alata* had the higher ability to reduce ferric ions. The procedure of FRAP assay is relatively simple and easy to be standardized. But, this assay has been reported not to react fast enough with some antioxidants, such as glutathione (Guo et al., 2003). Although the trend for both DPPH and FRAP free radical scavenging activity appeared the same the absolute values obtained were higher in DPPH assay. The  $\text{TEAC}_{\text{DPPH}}$  values were consistently higher than those obtained for  $\text{TEAC}_{\text{FRAP}}$ . Gil et al. (2002) reported that the FRAP values were higher than the



**Figure 2.** Percent antioxidant activities (%) of *L. pumila* leaf extract using  $\beta$ -carotene bleaching method. Data were analyzed using One-Way ANOVA and the different contrasted using Tukey's multiple range tests. Different letter(s) indicate the values are significantly different ( $p < 0.05$ ).



**Figure 3.** Vitamin C content (mg/g FW) of *L. pumila* leaf extracts. Data were analyzed using One-Way ANOVA and the different contrasted using Tukey's multiple range tests. Different letter(s) indicate the values are significantly different ( $p < 0.05$ ).

DPPH free radical scavenging activities of stone fruits. Similarly, Wong et al. (2006) reported that the ferric ions reducing activities of 25 tropical plants, expressed as Trolox equivalent antioxidant capacity were higher to that of DPPH free radical scavenging activities. In this study the reason is probably due to the difference in compounds that are reactive towards the two different methods. Wong et al. (2006) suggested that there are some

compounds such as polyphenols which may be more efficient as reducing agents for ferric ions but some may not scavenge DPPH free radicals due to steric differences. In both cases, however, the antioxidative ability of the control BHT was significantly higher than in the *L. pumila* leaf extracts.

### $\beta$ -Carotene bleaching assay

Besides the two methods described above  $\beta$ -carotene bleaching method expressed as percent (%) of total antioxidant activity (Figure 2) was used in determining antioxidant activity. Again the values indicated that *L. pumila* var. *Alata* has higher percentage of antioxidative activity ( $89.72 \pm 0.95\%$ ) than *L. pumila* var. *Pumila* ( $59.09 \pm 2.24\%$ ). *L. pumila* var. *Alata* had exceptionally high antioxidant level.

### Ascorbic acid content


Ascorbic acid also known as vitamin C is one of the most abundant antioxidants in plant where the role of ascorbate is to protect plants against oxidative stress (Smirnov, 2000). It is a powerful water soluble antioxidant and its established role is to prevent scurvy (Padayatty et al., 2003). The result in Figure 3 further showed that there is a slight different in vitamin C content and *L. pumila* var. *Alata* appeared to have a higher vitamin C content ( $0.022 \pm 0.001$  mg/g FW) when compared with var. *Pumila* ( $0.019 \pm 0.001$  mg/g FW).

### $\beta$ -Carotene content

Carotenoids are also classified among the basic constituents of the antioxidative effect (Duthie et al., 2003; Kidmose et al., 2001). Figure 4 shows the  $\beta$ -carotene content of *L. pumila* var. *Alata* and *L. pumila* var. *Pumila*. The highest content of  $\beta$ -carotene is in *Labisia pumila* var. *Alata* followed by var. *Pumila*,  $3.175 \pm 0.12$  mg/g FW and  $2.34 \pm 0.16$  mg/g FW, respectively. The result indicated that  $\beta$ -carotene content was higher in var. *Alata*. The red colored leaves of var. *Alata* may contribute to the high  $\beta$ -carotene content as opposed to var. *Pumila* where the leaves appear greener. This finding is in parallel with the report by Gil et al. (2002) which reports that total carotenoids is higher in yellow-colored nectarines compared to that of white-flesh.  $\beta$ -carotene is a precursor of vitamin A, which is important in human vision and also to prevent certain types of cancer (Craft et al., 1993; Colditz, 1987).

### Total anthocyanin content

Anthocyanins are the naturally occurring phenolic compounds responsible for the color of many flowers, fruits, and berries (Cooper-Driver, 2001). It is the most impor-



**Figure 4.** β-Carotene, flavonoid, anthocyanin, phenolic content (mg/g FW) of *L. pumila* leaf extracts. Data were analyzed using One-Way ANOVA and the different contrasted using Tukey's multiple range tests. The result are the mean  $\pm$  SE. (n = 5). Different letter(s) indicate the values are significantly different (p < 0.05).

tant group of water soluble-pigments in plants and had the beneficial health effects as antioxidant and anti-inflammatory agents (Wang and Jiao, 2000; Wang et al., 1999; Tamura and Yamagami, 1994). The total anthocyanin content (Figure 4) was higher in *L. pumila* var. *Alata* ( $0.328 \pm 0.07$  mg/g FW) compared to that of *L. pumila* var. *Pumila* ( $0.058 \pm 0.01$  mg/g FW). From this result, it shows that anthocyanin-rich samples generally showed very strong activities (var. *Alata*) and confirm that anthocyanins possess strong antioxidant activities (Wang et al., 1997; Tsuda et al., 1994). Anthocyanins are probably the largest group of phenolics compounds in the human diet, and their strong antioxidant activities suggest their importance in maintaining health. Anthocyanin is also important as antioxidant, which have roles in promoting good health and reducing the risk of chronic disease and also as anti-inflammatory agents. It was reported by (Wang and Jiao 2000; Wang et al., 1999; Tamura and Yamagami, 1994) that anthocyanins possess some positive therapeutic effects, mainly associated with their antioxidant activities.

#### Total flavonoid content

Flavonoids are naturally occurring substances in plants that are thought to have positive effects on human health (Montoro et al., 2005). The most important function of flavonoids is the antioxidants properties. Flavonoids have been shown to be highly effective scavengers of most types of oxidizing molecules, including singlet oxygen and various free radicals (Bravo, 1998). The flavonoid

content was higher in *L. pumila* var. *Pumila* ( $1.281 \pm 0.11$ mg/g FW) compared to that of *L. pumila* var. *Alata* ( $0.732 \pm 0.05$  mg/g FW) (Figure 4). There are different classes of flavonoids and the protective activity may differ with each of the compounds present. It was reported that quercetin, kaemferol, catechin and taxifolin were shown to suppress the cytotoxicity of O<sub>2</sub> and H<sub>2</sub>O<sub>2</sub> on Chinese hamster V79 cells in a protective manner (Nakayama et al., 1993). Flavonoid distribution in plants depends on the several factors including variation according to plant phyla/order/family and population variations within species (Harborne, 1986). The distribution pattern depends on the degree of accessibility to light and previous illumination because formation of the higher oxidized flavonoids is accelerated by light (Smith et al., 2000). Lee et al. (2005) reported that immature fruit generally contained lower levels of lutein and zeaxanthin than mature, color fruit, but the differences were not always statistically significant. The antioxidant property of flavonoids was the first mechanism of the action studied, particularly with regard to their protective effects against cardiovascular diseases. Flavonoids have been shown to be highly effective scavengers of most types of oxidizing molecules, including singlet oxygen and various free radicals (Bravo, 1998) that are probably involved in several diseases. Harborne and Williams (2000) suggested that additional benefit of flavonoids is their ability to stabilize membranes by decreasing membrane fluidity.

#### Total phenolic content

Phenolics are class of low molecular weight secondary

**Table 1.** Correlation ( $R^2$ ) between antioxidant activities (TEAC<sub>DPPH</sub> and TEAC<sub>FRAP</sub>) and  $\beta$ -carotene, flavonoid, vitamin C, anthocyanin and phenolic content ( $R^2$ ) in *L. pumila* var. *Alata*

	DPPH ( $R^2$ )	FRAP ( $R^2$ )
<b>Beta-carotene</b>	0.99	0.96
<b>Flavonoid</b>	0.97	0.95
<b>Vitamin C</b>	0.92	0.94
<b>Anthocyanin</b>	0.87	0.86
<b>Phenolic</b>	0.81	0.88

metabolite that found in most land plants (Akowuah et al., 2004). Many phenolic compounds have been reported to possess potent antioxidant activity and have anti-cancer or anti-carcinogenic, anti-bacterial, anti-viral or anti-inflammatory activities to a greater or lesser extent (Chung et al., 1998; Cassidy et al., 2000; Gao et al., 2000; Tapiero et al., 2002). The phenolic content of *L. pumila* was shown in Figure 4. There was no significant difference of total phenolic content in *L. pumila*. The content of phenolics is more or less the same among the two varieties which ranged from  $2.53 \pm 0.04$  mg/g FW to  $2.55 \pm 0.06$  mg/g FW (Figure 4).

## DISCUSSION

The antioxidant properties of *L. pumila* have not been reported previously. Its use as a traditional herb for women appears to have some basis in relation to the total antioxidant content and in relation to report that an *Alectoria crispa* plants which comes from the family as *L. pumila* have utero wall-contracting saponins (Jansakul et al., 1987). Further studies on the specific components such as the saponins, enzymatic antioxidants will be carried out. *L. pumila* var. *Alata* had higher total antioxidant activity compared to that of var. *Pumila* in all the three different methods which is FRAP, DPPH and  $\beta$ -carotene bleaching assays. There is a positive correlation between total antioxidant activities and the individual antioxidant compounds with a range correlation coefficient of  $R^2 = 0.81$  to  $0.99$  (Table 1). The highest correlation based on DPPH/FRAP assays is with  $\beta$ -carotene (0.99/0.96) followed by flavonoids (0.97/0.95), vitamin C, (0.92/0.94) anthocyanins (0.87/0.86) and total phenolics (0.81/0.88). It is therefore, appears that the antioxidative activities are primarily due to  $\beta$ -carotene and flavonoids. Such correlations have been observed in other studies. Chanwitheesuk et al. (2005) reported that the contents of vitamins C and E, total carotenes, total xanthophylls, tannins and total phenolics of some plants are correlated with antioxidant activities. Such correlations have been widely reported. Studies on the antioxidant activities of Algerian medicinal plants linear correlation of Trolox equivalent antioxidant capacity (TEAC) with respect to the total and flavonoid content (Djeridane et al., 2006) where there is a

positive correlation between the total phenolic content of a given sample and its antioxidant activity. However the value is  $R^2 = 0.7931$ . There was a strong correlation (0.93-0.96) between total phenolics and antioxidant activity in stone fruits nectarines, peaches and plums but low correlation with vitamins C and carotenoids (Gil et al., 2002). A similar correlation was obtained by Wong et al. (2006). Further studies to identify the different flavonoids will be identified as each flavonoid exhibits different scavenging capacities (Cai et al., 2006).

## ACKNOWLEDGEMENTS

The authors wish to thank Universiti Putra Malaysia for the financial support.

## REFERENCES

- Akowuah GA, Ismail Z, Norhayati I, Sadikum A, Khamsah S (2004). Sinensetin, eupatorin, 3'-hydroxy-5, 6, 7, 4'-tetramethoxyflavone and rosmarinic acid content and antioxidative effects of *Orthosiphon stamineus* from Malaysia. Food Chem. 87: 559-566.
- Benzie IFF, Strain JJ (1996). The ferric reducing ability of plasma (FRAP) as a measure of 'Antioxidant Power': the FRAP assay. Anal. Biochem. 239: 70-76.
- Bharti AK, Khurana JP (2003). Molecular characterization of *transparent testa (tt)* mutants of *Arabidopsis thaliana* (ecotype Estland) impaired in flavonoid biosynthetic pathway. Plant Sci. 165: 1321-1332.
- Bravo L (1998). Polyphenols: chemistry, dietary sources, metabolism, and nutritional significance. Nutr. Rev. 56: 317-333.
- Cai YZ, Sun M, Xing J, Luo Q, Corke H (2006). Structure-radical scavenging activity relationships of phenolic compounds from traditional Chinese medicinal plants. Life Sci. 78: 2872-2888.
- Cassidy A, Hanley B, Lamuela-Raventos RM (2000). Isoflavones, lignans and stilbenes: origins, etabolism and potential importance to human health. J. Sci. Food Agric. 80 (7): 1044-1062.
- Chanwitheesuk A, Teerawutgulrag A, Rakariyatham N (2005). Screening of antioxidant activity and antioxidant compounds of some edible plants of Thailand. Food Chem. 92: 491-497.
- Chung KT, Wong TY, Huang YW, Lin Y (1998). Tannins and human health: a review. Crit. Rev. Food Sci. Nutr. 38 (6): 421-464.
- Colditz GA (1987). Beta-carotene and cancer. In: Quebedeaux, B, Bliss, FA (Eds), Horticulture and Human Health, Contributions of Fruits and Vegetables. Proceedings of ASHS Symposium Series No. 1 held at Prentice-Hall, Upper Saddle River, New Jersey, pp. 150-159.
- Cooper-Driver GA (2001). Contributions of Jeffrey Harborne and co-workers to the study of anthocyanins. Phytochemistry 56: 229-236.
- Craft NE, Wise SA, Soares JH Jr (1993). Individual carotenoid content of SRM 1548 total diet and influence of storage temperature, lyophilization and irradiation on dietary carotenoids. J. Agric. Food Chem. 41: 208-213.
- Davies SHR, Masten SJ (1991). Spectrophotometric method for ascorbic acid using dichlorophenolindophenol: elimination of the interference due to iron. Anal. Chim. Acta 248: 225-227.
- De Tommasi N, Piacente S, De Simone F, Pizza C, Liang ZZ (1993). Characterization of three new triterpenoid saponins from *Ardisia japonica*. J. Nat. Prod. 56: 1669.
- Djeridane A, Yousfi M, Nadjemi B, Boutassouna D, Stocker P, Vidal N (2006). Antioxidant activity of some Algerian medicinal plant extracts containing phenolic compounds. Food Chem. 97: 654-660.
- Duthie GG, Gardner PT, Kyle JAM (2003). Plant polyphenols: are they a new magic bullet. Proc. Nutr. Soc. 62: 599-603.
- Gao D, Kakuma M, Oka S, Sugino K, Sakurai H (2000). Reaction of  $\beta$ -alkannin (shikonin) with reactive oxygen species: detection of  $\beta$ -alkannin free radicals. Bioorg. Med. Chem. 8: 2561-2569.
- Gil MI, Thomas-Barberan FT, Hess-Pirce B, Kader AA (2002). Antioxi-

- dant capacities, phenolic compounds, carotenoids and vitamin C contents of nectarine, peach and plum cultivars from California. *J. Agric. Food Chem.* 50: 4976-4982.
- Guo C, Yang J, Wei J, Li Y, Xu J, Jiang Y (2003). Antioxidant activities of peel, pulp and seed fractions of common fruits as determined by FRAP assay. *Nutr. Res.* 23: 1719-1726.
- Harborne JB (1973). *Phytochemical Methods*. In: Harborne JB (eds) *A Guide to Modern Techniques of Plant Analysis*, Chapman and Hall, London, pp. 33-80.
- Harborne JB (1986). Nature, distribution, and function of plant flavonoids. In: Cody V, Middleton CE, Harborne JB (eds) *Plant flavonoids in biology and medicine: biochemical, pharmacological and structure-activity relationships*. Proceedings of conference held at Buffalo, New York, pp 15-24.
- Harborne JB, Williams CA (2000). Advances in flavonoid research since 1992. *Phytochemistry* 55: 481-504.
- Huang J, Ogihara Y, Zhang H, Shimizu N, Takeda T (2000). Triterpenoid saponins from *Ardisia mamillata*. *Phytochemistry* 54: 817-822.
- Huang J, Zhang H, Shimizu N, Takeda T (2003). Ardisiamamillosides G and H, two new triterpenoid saponins from *Ardisia mamillata*. *Chem. Pharm. Bull.* 51 (7): 875-877.
- Jansakul C, Baumann H, Kenne L, Samuelsson G (1987). Ardisiacrispin A and B, two utero-contractin saponins from *Ardisia crispata*. *Planta Med.* 53(5): 405-409.
- Kidmose U, Knuthsen P, Edelenbos M, Justesen U, Hegelund E (2001). Carotenoids and flavonoids in organically grown spinach (*Spinacia oleracea* L.) genotypes after deep frozen storage. *J. Sci. Food Agric.* 81: 918-923.
- Lee JJ, Crosby KM, Pike LM, Yoo KS, Leskovaar DI (2005). Impact of genetic and environmental variation on development of flavonoids and carotenoids in pepper (*Capsicum* spp.). *Sci. Hort.* 106: 341-352.
- Lisiewska Z, Kmiecik W, Korus A (2006). Content of vitamin C, carotenoids, chlorophylls and polyphenols in green parts of dill (*Anethum graveolens* L.) depending on plant height. *J. Food Compos. Anal.* 19(2-3): 134-140.
- Longo L, Vasapollo G (2006). Extraction and identification of anthocyanins from *Smilax aspera* L. berries. *Food Chem.* 94: 226-231.
- Marinova D, Ribarova F, Atanassova M (2005). Total phenolics and total flavonoids in Bulgarian fruits and vegetables. *J. Univ. Chem. Technol. Metallurgy* 40(3): 255-260.
- Mokbel MS, Hashinaga F (2006). Evaluation of the antioxidant activity of extracts from buntan (*Citrus garandis* Osbeck) fruit tissues. *Food Chem.* 94: 529-534.
- Montoro P, Braca A, Pizza C, De Tommasi N (2005). Structure-antioxidant activity relationships of flavonoids isolated from different plant species. *Food Chem.* 92: 349-355.
- Nakayama T, Yamada Y, Osawa T, Kawakishi S (1993). Suppression of active oxygen-induced cytotoxicity by flavonoids. *Biochem. Pharmacol.* 45: 265-267.
- Padayatty SJ, Katz A, Wang Y, Eck P, Kwon O, Lee JH, Chen S, Corpe C, Dutta A, Dutta SK, Levine M (2003). Vitamin C as an antioxidant: Evaluation of its role in disease prevention. *J. Am. Coll. Nutr.* 22 (1): 18-35.
- Saha K, Lajis NH, Israf DA, Hamzah AS, Khozirah S, Khamis S, Syahida A (2004). Evaluation of antioxidant and nitric oxide inhibitory activities of selected Malaysian medicinal plants. *J. Ethnopharmacol.* 92: 263-267.
- Singleton VL, Rossi JA (1965). Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagent. *Am. J. Soc. Enol. Viticul.* 16: 144-158.
- Smirnoff N (2000). Ascorbic acid: metabolism and functions of a multifaceted molecule. *Curr. Opin. Plant Biol.* 3: 229-235.
- Smith G, Thomsen SJ, Markham KR, Andary C, Cardon D (2000). The photostabilities of natural occurring 5-hydroxyflavones, flavonols, their glycosides and their aluminum complexes. *J. Photochem. Photobiol. A: Chem.* 136: 87-91.
- Tamura H, Yamagami A (1994). Antioxidative activity of monoacylated anthocyanins isolated from Muscat bailey A grape. *J. Agric. Food Chem.* 42: 1612-1615.
- Tapiero H, Tew KD, Nguyen Ba G, Mathé G (2002). Polyphenols: do they play a role in the prevention of human pathologies? *Biomed. Pharmacother.* 56: 200-207.
- Tsuda T, Ohshima K, Kawakishi S, Osawa T (1994). Antioxidative pigments isolated from seed of *Phaseolus vulgaris* L. *J. Agric. Food Chem.* 42: 248-251.
- Velioglu YS, Mazza G, Gao L, Oomah BD (1998). Antioxidant activity and total phenolics in selected fruits, vegetables, and grain products. *J. Agric. Food Chem.* 46: 4113-4117.
- Wan Ezumi MF, Siti Amrah S, Suhaimi AWM, Mohsin SSJ (2007). Evaluation of the female reproductive toxicity of aqueous extract of *Labisia pumila* var. *alata* in rats. *Indian J. Pharmacol.* 39 (1): 30-32.
- Wang H, Cao G, Prior RL (1997). Oxygen radical absorbing capacity of anthocyanins. *J. Agric. Food Chem.* 45: 304-309.
- Wang H, Nair MG, Strasburg GM, Chang YC, Booren AM, Gray JI, DeWitt DL (1999). Antioxidant and anti-inflammatory activities of anthocyanins and their aglycon, cyanidin, from tart cherries. *J. Nat. Prod.* 62: 294-296.
- Wang SY, Jiao H (2000). Scavenging capacity of berry crops on superoxide radicals, hydrogen peroxide, hydroxyl radicals and singlet oxygen. *J. Agric. Food Chem.* 48: 5677-5684.
- Wong CC, Li HB, Cheng KW, Chen F (2006). A systematic survey of antioxidant activity of 30 Chinese medicinal plants using the ferric reducing antioxidant power assay. *Food Chem.* 97: 705-711.
- Zhishen J, Mengcheng T, Jianming W (1999). The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chem.* 64: 555-559.