

Full Length Research Paper

Prevalent home management techniques and outcome among mothers of febrile children in Eastern Nigeria

U. M. Chukwuocha*, G. C. Nwakwuo, C. Emerole, I. N. S. Dozie and O. E. C. Nwuda

Department of Public Health Technology, School of Health Technology, Federal University of Technology, P.M.B 1526, Owerri, Imo State, Nigeria.

Received 6 November, 2013; Accepted 13 February, 2014

The magnitude of the consequences and the extent of problem associated with febrile illnesses in children have been shown to be associated to mothers' behaviour in treatment, decision making and prevention of childhood illnesses which are in turn influenced by social, cultural and intra-household relations. This study employed a cross-sectional survey design to assess the method employed by caregivers in the management of febrile conditions of their children in parts of Eastern Nigeria. About 100 women were sampled using multistage sampling techniques. Common actions taken by mothers and caregivers in the management of febrile include administration of home drug (32%), tepid sponging (4%), striping the baby naked (16%) and took baby/child to hospital (48%). Result from the study showed significant relationship between age of mother ($p<0.05$), educational attainment of mother ($p<0.05$), number of children ($p<0.05$) as well as occupation of mothers ($p<0.05$) and management action taken. Caregivers need to be properly aware of ill health conditions of their children and be properly equipped with prowess to manage them. There is also need for the governments through the primary health care system to improve their interaction with the grass root women and educate them more on integrated management of childhood illness.

Key words: Children, febrile, management techniques, outcome, Eastern Nigeria.

INTRODUCTION

Febrile illness refers to a feverish condition which is usually associated with malaria, measles, neonatal tetanus, pneumonia, whooping cough (Feyisetan et al., 1997; Ewoigbokhan, 2000).

Traditionally, malaria has been regarded as the most common and important febrile illness in Sub-Saharan Africa (Greenwood et al., 2005). However, many of the physicians' diagnosis of febrile conditions in children also include pneumonia, diarrhoea, and other medical conditions such as otitis media, anaemia and acute eye

problems (Madubuike, 1997; Osikoya and Sebanjo, 2008).

A study on the magnitude of the consequences and the extent of problems associated with febrile illnesses in children reported that more than 4.4 million children die every year in sub-Saharan Africa (Black et al., 2003) and most children die from fever at home without receiving adequate therapy. Only a small number of children with fever are treated at health-care facilities.

A number of elements are critical to management of the

*Corresponding author. E-mail: uchukwuocha@gmail.com. Tel: +2348034712957.

Author(s) agree that this article remain permanently open access under the terms of the [Creative Commons Attribution License 4.0 International License](http://creativecommons.org/licenses/by/4.0/)

child with fever. The caregiver must recognize the illness and provide the appropriate supportive care. According to Chukwuocha (2011), diagnosis of malaria at home by caregivers was mostly by recognising symptoms such as constant fever, headache, coughing and pains, loss of appetite, etc.

An efficacious drug must be available either at home or nearby when the child becomes ill and the drug must be affordable for the caregiver. But these drugs are not readily available to the first calls of most caregivers (the Patent Medicine Dealers-PMDs) and even when available, they are usually of high cost (Chukwuocha et al., 2013). While some mothers in the community think that drugs provided in the community are very weak or of a lower quality, some believe that the drugs provided in the clinics and shops are better off, whereas others do not seek for drug treatment at all. This may be as a result of cultural or religious beliefs (Kidane and Morrow, 2000).

Earlier studies on home health care had pointed out that women assume the burden of the care for their children (Sims et al., 1992; Molyneux, 2002). And with issues surrounding pregnancy extremely viewed as women issues (Jhpiego, 2001; Nwakwuo and Oshonwoh, 2013), the health beliefs of these women can greatly influence the outcome of febrile illnesses in children such that action taken results in good or poor outcomes. Most mothers according to Nsungwa et al. (2007) become upset and aggressive in management of fever conditions, and these may be determined by their level of education, age of the mother, occupations, religion and other socio economic/cultural factors. However, the fact that some caregivers prefer handling the case at home before going to hospital for further treatment might be greatly influenced by their knowledge of necessary health management techniques.

Home treatment is a common practice among caregivers of under-five-year old children (Demming et al., 1999; Fawole and Onadeko, 2001; Olaogun, 2005). The finding of Ajibade (2013) affirms that caregivers of children play active role in the health care of these children. Thus, the challenge is to work on this natural tendency for self-treatment, as a means of ensuring early recognition of illness and ensure safe, prompt and appropriate action.

Further studies on home health care have also shown that behaviour in treatment, decision making and prevention of childhood illnesses are influenced by some factors such as social, cultural and intra-household relations (Feyisetan et al., 1997; Molyneux, 2002). This by implication indicates that the adequacy of care provided for the children is a reflection of capability of taking control of the difficulties associated with complexity and dynamism of these factors (Ajibade, 2013). This study intends to access the prevalent home management techniques and outcome among mothers of febrile children in Eastern Nigeria in relation to factors that influence the use of such techniques.

MATERIALS AND METHODS

This study employed a cross-sectional survey design to assess the methods employed by caregivers in the management of febrile conditions of their children.

Out of a total of 154 women of child bearing age permanently residing in Umuevu (resided in Umuevu in the 12 months), 100 respondents were selected. Using a multistage sampling technique, 3 villages that make up Umuevu in Eastern Nigeria were clustered and 100 women were randomly selected proportionate to the number of women of child bearing age in each cluster.

Questionnaire was used to seek information from mothers/caregivers on their knowledge of signs and symptoms of ill health conditions among children, cultural dispositions and techniques and factors in their management.

The questionnaire was validated for both content and criterion by an expert and the reliability (0.78) of the questionnaire was assured through pilot study.

Data collected were collated and compiled into a spreadsheet. Raw data was analyzed using SPSS version 20 and tables and charts were presented. Fishers' exact and chi-square was used to provide inferential conclusion.

RESULTS

Table 1 shows the socio-demographic characteristic of the respondents. Greater percentages of the mothers are aged 18 to 30 years (48%), 31 to 42 years (42%) and 98% of them are of Christians background. About 54% of the respondents has attained a form of tertiary level of education with few (8%) having no formal education. The common occupations of the mothers are trading (38%) and civil service (46%) with few farmers (10%). Large number of the mothers in the study has one child (64%) with 26% having around 2 to 4 children. More febrile condition is reported for female children (54%) and is common around age 1 to 2 years (32%), 3 to 4 years (30%) and less than 1 year (20%).

Table 2 shows mothers' knowledge of the child's febrile condition. More febrile were noticed in prior four days to the study and most reported episode has occurred only but once (60%) with malaria remaining the common cause (64%), followed by pneumonia (28%). Although mothers and mother in-laws play greater role in the identification of febrile conditions (50%), husbands (22%) has been helpful with only 8% of conditions identified at the health care facility level by health workers.

Table 3 shows that 48% of mothers prefer taking children to hospitals, while 32% prefer to administer drugs at home. Greater percentage (50%) of actions taken by mothers is usually immediately a sign febrile is noticed with 40% taken in less than 24 h of the identification of signs of febrile condition. Apparently, 88% of actions taken have resulted in improved health conditions and 43% intends to take children to the hospital in future occurrences, while 38% will prefer to give drug at home even if subsequent action is required.

From Table 4, mothers were shown to have identified febrile through loss of appetite 46 (24%), dullness 32 (16.7%), hand touching 30 (15.6%), shivery and

Table 1. Demographic characteristics.

| Variable | Frequency | Percentage |
|-------------------------------------|-----------|------------|
| Age of mother | | |
| 18 - 30 years | 48 | 48.0 |
| 31 - 42 years | 42 | 42.0 |
| 43 years and above | 10 | 10.0 |
| Religion | | |
| Traditional Christian | 80 | 80.0 |
| Modern Christian | 18 | 18.0 |
| Traditionalist | 2 | 2.0 |
| Level of education of mother | | |
| No formal education | 8 | 8.0 |
| Primary | 6 | 6.0 |
| Secondary | 32 | 32.0 |
| Tertiary | 54 | 54.0 |
| Occupation of mother | | |
| Farming | 10 | 10.0 |
| Trading | 38 | 38.0 |
| Civil servant | 46 | 46.0 |
| Student | 6 | 6.0 |
| Number of children | | |
| 1 child | 64 | 64.0 |
| 2 – 4 children | 26 | 26.0 |
| Above 4 children | 10 | 10.0 |
| Sex of febrile child | | |
| Female | 54 | 54.0 |
| Male | 46 | 46.0 |
| Age of febrile child | | |
| Less than 1 year | 20 | 20.0 |
| 1 - 2 years | 32 | 32.0 |
| 3 - 4 years | 30 | 30.0 |
| Above 4 years | 18 | 18.0 |

restlessness 24 (12.5%).

Table 5 shows that 72 (35%) of causes of febrile are noticed by laboratory test, fever 54 (26.2%). Table 6 shows relationships between different variables and common actions taken by mothers and/or caregivers. A significant relationship was shown between age of mother and action taken (fishers exact, $p=0.000$) as younger mothers are more likely to administer drugs at home whereas elderly mothers tend to take their children to the hospital.

The relationship between educational attainment of mother and action taken (fishers exact, $p=0.000$) was also significant. Mothers with no formal education, primary and secondary educations are more likely to administer

home drugs aside other actions, whereas mothers with advance education go for hospitals as their first action for febrile children.

Also from the same table, a significant relationship was shown between the number of children and action taken (fishers exact, $p=0.000$). Mothers with single child prefer to administer drugs at home, tepid sponge or stripe the baby naked compared to those with more children (2 and above) who prefer to use the hospital.

Occupation of mother and/or caregivers was also significantly (fishers exact, $p=0.000$) related to their different action or actions taken. Farmers and traders are more likely to administer drug at home with formers tepid sponging and striping babies more. Figure 1 shows that the relationship between mother's monthly income and actions taken (fishers exact, $p=0.000$) is significant. Mothers with high income are more likely to take their children to hospitals.

DISCUSSION

Parents and care providers have in several studies identified "fever" as a significant indicator of illness in children (Osikoya and Sebanjo, 2008; Poirier et al., 2000; Akpede, and Akenzua, 2001). A number of malaria-related symptoms particularly fever, shivering, chills, loss of appetite and headache were presented, which taken together may approximate a clinical diagnosis of malaria although using these symptoms, particularly fever, as a proxy for malaria appears to be neither sensitive nor specific when compared with parasitologically confirmed diagnosis (Font et al., 2001; Amexo et al., 2004). This implies that fever dictates the needs of treatment and what mothers do.

Traditionally, malaria has been regarded as the most common and important febrile illness in sub-Saharan Africa (Greenwood et al., 2005). This study indicates that 64% of reported febrile illnesses are due to malaria. Only 28% is due to pneumonia.

Most children die from fever at home without receiving adequate therapy and although the recognition of illness and provision of the appropriate dose of drugs, together with supportive care is a critical element in the management of febrile, access to health services and the quality of care administered at all levels of health care have been considered as the central determinants of health outcomes (WHO, 2000). The survival of children in developing countries depends on the family's and community's ability to access basic needs of life (Nnebue et al., 2010).

Some studies have shown that home treatment is a common practice among caregivers of under-five-year old children (Demming et al., 1999; Fawole and Onadeko, 2001; Olaogun et al., 2005; Nwankwo et al., 2009). Other studies on home health care pointed out that women assume the burden of the care (Sims et al., 1992; Molyneux et al., 2002). The perceived illness severity, maternal recognition of certain signs and symptoms of




Figure 1. Relationship between monthly income of mothers and action taken (p=0.000).

Table 2. Mothers' knowledge of child's febrile condition.

| Variable | Frequency | Percentage |
|--|-----------|------------|
| When the child had febrile condition | | |
| Last four days | 30 | 30.0 |
| Last one week | 26 | 26.0 |
| Last two weeks | 20 | 20.0 |
| Last one month | 24 | 24.0 |
| Frequency of episodes | | |
| Just once | 60 | 60.0 |
| Two days interval | 26 | 26.0 |
| Three days interval | 14 | 14.0 |
| Cause of febrile condition | | |
| Pneumonia | 28 | 28.0 |
| Malaria | 64 | 64.0 |
| Measles | 8 | 8.0 |
| Who noticed febrile condition if not mother | | |
| Sister/sister in-law | 8 | 8.0 |
| Brother/Brother in-law | 2 | 2.0 |
| Mother/Mother in-law | 50 | 50.0 |
| Neighbour/friend | 10 | 10.0 |
| Husband | 22 | 22.0 |
| Health worker | 8 | 8.0 |

Table 3. Mother's action during child's illness episode.

| Variable | Frequency | Percentage |
|--|-----------|------------|
| Action taken | | |
| Administer home drugs | 32 | 32.0 |
| Tepid sponge | 4 | 4.0 |
| Striped the baby naked | 16 | 16.0 |
| Took to hospital | 48 | 48.0 |
| Time between sign and action taken | | |
| Immediately | 50 | 50.0 |
| Action in less than 24 hours | 40 | 40.0 |
| Action more than 24 hours | 10 | 10.0 |
| Who informed action taken | | |
| Sister/sister in-law | 8 | 8.0 |
| Brother/Brother in-law | 2 | 2.0 |
| Mother/Mother in-law | 50 | 50.0 |
| Neighbour/friend | 10 | 10.0 |
| Husband | 22 | 22.0 |
| Health worker | 8 | 8.0 |
| Confident of action | | |
| Yes | 84 | 84.0 |
| No | 10 | 10.0 |
| Not sure | 6 | 6.0 |
| Result of action taken | | |
| Condition improved | 88 | 88.0 |
| Condition remain the same | 12 | 12.0 |
| Action to be taken in future episodes | | |
| Given drugs | 38 | 38.0 |
| Strip the body/baby naked | 43 | 43.0 |
| Take to hospital | 19 | 19.0 |

of childhood illness were critical factors determining health care seeking behaviour.

Rural residence has been shown by several studies to have the highest risk factor for febrile illnesses in under fives (Yoder and Hornik 1996; D'Souza, 1999; Goldman and Heuveline, 2000; Amarasiri et al., 2001). Children of mothers in rural settings with lower levels of education are more likely to suffer from fever as compared to urban children with better educated women (Yoder and Hornik, 1996; D'Souza, 1999; Goldman and Heuveline, 2000; Amarasiri et al., 2001). But according to Olaogun et al. (2006), education was however insignificant at 5% level. This is in contrast with the report that educational status has a positive effect on the health status of the family (Nnebue et al., 2010). Another study has also reported a strong relationship between mothers' education and

health seeking behaviour (Nigeria Demographic and Health Survey, 2003) and this is consistency with the significant relationship (fishers exact=111.493, $p<0.05$) shown in this study between educational attainment of mother and action taken.

A study by Olaogun et al. (2006) in Osun State Nigeria showed that mothers' age was negatively correlated with under-fives mothers' action. In contrary, (Nnebue, Ileadike, Nworah and Duru (2010) showed that the older the mothers' are, the more likely they would not take action when their under-fives develop febrile illness. This is study shows a significant relationship between age of mother and action taken (fishers exact=107.066, $p<0.05$) with younger mothers are more likely to administer drugs at home whereas elderly mothers tend to take their children to the hospital.

Further studies on behaviours in treatment decisions making and prevention of childhood illnesses are influenced by some factors such as social, cultural and intra-household relations (Feyisetan et al., 1997; Molyneux et al., 2002).

Occupation in this study was shown to be significantly (fishers exact=112.014, $p<0.05$) related with management action taken by caregivers. This is consistent with a positive correlation with under-five mothers' action reported in a study in Osun State (Olaogun et al., 2006) and the assertion by Hobbs and Blanks (1975) that occupation plays a major role in shaping the life style of individuals. Hobbs and Blanks (1975) reported increased women's access to resources and strengthens their bargaining power within the household and the workplace. The women's access to resources and their bargaining power within the household have a significant influence on their treatment seeking behaviour for their children (Nigeria Demographic and Health Survey, 2004).

Income according to Olaogun et al. (2006) was insignificant at 5% level. This result is contrary to the result of this study which shows a significant relationship (fishers exact=39.562, $p<0.05$) between monthly income of mothers and their action on management of children febrile conditions.

The findings from this study showed that about 48% of respondents took febrile children with 50% of decision taking, immediately febrile condition is noticed and 40% within 24 h. This is contrary to the findings by Olaogun et al. (2005) who found that only one respondent in their study reported to the clinic within 24 h of the child been sick (Olaogun et al., 2005).

Fever was the mostly mentioned sign observed by mothers in their sick children which prompted them to take action aside result from the laboratory. This finding affirms the reports from various studies that fever is a good sign for mothers or parents to seek for treatment of childhood febrile illnesses (Poirier et al., 2000; Agbolosu et al., 1997). Findings of this study affirm the several reports of the high prevalence of home treatment for under-five year children with febrile illnesses (Olaogun et al., 2005; Audu and Ogala, 1997; Lubanga et al., 1997).

Table 4. Mother's perception and understanding of child's febrile condition.

| Variable | Frequency | Percentage |
|---|-----------|------------|
| *How febrile condition was noticed | | |
| Hand washing | 30 | 30.0 |
| Shivery | 24 | 24.0 |
| Uses of clinical thermometer | 10 | 10.0 |
| Restlessness | 24 | 24.0 |
| Loss of appetite | 46 | 46.0 |
| Dullness in movement | 32 | 32.0 |
| Joint/body pain | 12 | 12.0 |
| Rashes | 4 | 4.0 |
| Others | 10 | 10.0 |

*Multiple responses

Table 5. Mothers knowledge of causes of febrile condition.

| Variable | Frequency | Percentage |
|-------------------------------------|-----------|------------|
| Causes of febrile conditions | | |
| Fever | 54 | 54.0 |
| Laziness/dullness | 26 | 26.0 |
| Loss of appetite | 32 | 32.0 |
| Rashes | 4 | 4.0 |
| Convulsion | 2 | 2.0 |
| Joint/body pains | 16 | 16.0 |
| Laboratory test result | 72 | 72.0 |

According to the reports of study carried out in Uganda, mothers gave modern drugs as action taken before coming to health facility. In Nigeria, reports showed that parents used more of herbs than modern drugs (Salako et al., 2001), but this study also indicates 32% administration of home drugs by caregivers. Tepid sponging and use of orthodox drugs are two common practices of home management and patent medicine dealers remains the main source of drugs. The practices of home management have to be exploited so that childhood febrile illnesses could be controlled through appropriate treatment (Ajibade, 2013). The children would be highly benefited from pre-packaged drugs, and mothers from integrated management of childhood illness education.

Conclusion

The magnitude of the consequences and the extent of problems associated with febrile illnesses in children is enormous (Black et al., 2003).

Caregivers must therefore be able to recognize the illness so as to provide the appropriate supportive care and access to health services, since the quality of care administered at all levels of health care have been considered as the central determinants of health outcomes (WHO, 2000).

Although this study shows a good relationship between ages, education, household size and occupation of mothers and the action taken on their children's health and the fact that greater percentage visits the hospital during febrile conditions. Younger mothers with low education are more likely to administer drugs at home or apply one prevention method or the other. Fever has been shown not to be the only way of recognizing febrile illness, but also loss of appetite, body pain and restlessness.

Further more, although greater actions taken by caregivers (mothers) resulted in improved health conditions, caregivers and mothers need to be properly aware of ill health conditions of their children and be properly equipped with prowess to manage them, because the adequacy of care knowledge for children will improve their capability of taking control of the difficulties associated with complexity and dynamism of factors influencing their decisions and improve integrated management of childhood illness.

LIMITATION OF STUDY

As most women of child bearing age in the community were not permanently residing in the village, it was difficult to reach more respondents or increase the sample

Table 6. Relationship between variables.

| Parameter | Action taken | | | | Total |
|---|-----------------------|--------------|------------------------|------------------|----------|
| | Administer home drugs | Tepid sponge | Striped the baby naked | Took to hospital | |
| Age of mother and action taken: p=0.000 | | | | | |
| Age of mother | | | | | |
| 18 – 30 years | 32 (66.7) | 4 (8.3) | 12 (25.0) | 0 (0.0) | 48 (100) |
| 31 – 42 years | 0 (0.0) | 0 (0.0) | 4 (9.5) | 38 (90.5) | 42 (100) |
| 43 years and above | 0 (0.0) | 0 (0.0) | 0 (0.0) | 10 (100) | 10 (100) |
| Level of education of mother and action taken: p=0.000 | | | | | |
| Level of education | | | | | |
| No formal education | 8 (100) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 8 (100) |
| Primary | 6 (100) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 6 (100) |
| Secondary | 18 (56.3) | 4 (12.5) | 10 (31.3) | 0 (0.0) | 32 (100) |
| Tertiary | 0 (0.0) | 0 (0.0) | 6 (11.1) | 48 (88.9) | 54 (100) |
| Number of children and action taken: p=0.000 | | | | | |
| Number of children | | | | | |
| 1 child | 32 (50) | 4 (6.3) | 16 (25) | 12 (18.8) | 64 (100) |
| 2 – 4 children | 0 (0.0) | 0 (0.0) | 0 (0.0) | 26 (100) | 26 (100) |
| Above 4 children | 0 (0.0) | 0 (0.0) | 0 (0.0) | 10 (100) | 10 (100) |
| Occupation and action taken: p=0.000 | | | | | |
| Occupation | | | | | |
| Farming | 10 (100) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 10 (100) |
| Trading | 22 (57.9) | 4 (10.5) | 12 (31.6) | 0 (0.0) | 38 (100) |
| Civil servant | 0 (0.0) | 0 (0.0) | 4 (8.7) | 42 (91.3) | 46 (100) |
| Student | 0 (0.0) | 0 (0.0) | 0 (0.0) | 6 (100) | 6 (100) |

size. Even the available women are either farmers or traders, thus difficult to reach except during antenatal clinics or vital clinic days.

Also, the difficulty of mothers and some women to recall any previous episode of febrile illness of their children and the fear to comment on their practices of self medication. There was also a low level of knowledge about integrated management of childhood illness.

RECOMMENDATIONS

Based on the findings of this study, the flowing recommendations have been proffered: the government through the primary health care sector has to improve their interaction with the grass root women and educate them more on integrated management childhood illness; provision of insecticide treated bed nets will also help reduce the prevalence of malaria among children especially the under five; emphasis should me made on seeking adequate health care for children to mothers during antenatal and postnatal care visits to health care workers especially to young and inexperience mothers; the health care providers, non governmental and support

organization should also help mothers with requisite preventive emergency techniques for management of febrile conditions at home/community level; and access to health care facilities should be at a reasonable proximity.

Conflict of Interests

The author(s) have not declared any conflict of interests.

ACKNOWLEDGEMENTS

The authors gratitude goes to the management of the Department of Public Health Technology, Federal University of Technology, Owerri, Nigeria and all health facilities whose participation make this piece of research a success.

REFERENCES

Agbolosu NB, Cuevas LE, Milligan PE, Boardhead RL, Brewster D, Graham SM (1997). Efficacy of Tepid Sponging versus Paracetamol

- in Reducing Temperature in Febrile Children. *Annals of Tropical Paediatrics*, 17:283-288.
- Ajibade BL (2013). Mothers' Action and Preferences of Treatment of Febrile Illnesses among under- five -year- old Children in Osun State. *Journal of Biology, Agric. Healthcare* 3(7):148-155
- Akpede GO, Akenzua GI (2001). Management of children with prolonged fever of Unknown Origin and Difficulties in the Management of Fever of Unknown Origin in Children in Developing Countries. *Paediatric Drug* (3- 4):247-262
- Amarasiri de Silva MW, Wijekoon A, Hornik R, Martines J (2001). Care seeking in Sri Lanka: one possible explanation for low childhood mortality. *Soc. Sci. Med.* 53(10):1363-72.
- Amexo M, Tolhurst R, Bamish G, Bates I (2004). Malaria misdiagnosis: effects on the poor and vulnerable. *Lancet*, 364:1896-1898.
- Audu L, Ogala WN (1997). Home treatment of children with fever in Zaria, Nigeria. *Niger. Med. Pract.* 34:17-19.
- Black RE, Morris SS, Bryce JP (2003). Where and Why are 10 million dying: Quality of Treatment among Children Dying every year. *Lancet* 361:2226 -2234.
- Chukwuocha UM (2011). Rapid assessment of home management of malaria among caregivers in parts of South Eastern Nigeria. *Pan Afr. Med. J.* 10:29.
- Chukwuocha UM, Nwakwuo GC, Mmrole I (2013). Artemisinin-Based Combination Therapy: Knowledge and Perceptions of Patent Medicine Dealers in Owerri Metropolis, Imo State, Nigeria and Implications for Compliance with Current Malaria Treatment Protocol. *J. Community health* 38(4):759-765.
- Demming MS, Gayibor A, Murphy K, Jones TS, Karsa T (1999). Home Treatment of Febrile Children with Antimalarial Drugs. *Bull. WHO* 67(6):695-700.
- D'Souza RM (1999). Care seeking behavior. *Clin. Infect. Dis.* 28:234.
- Ewoigbokhan SE (2000). Morbidity Data and their Interpretation for Nigerian Children. Paper presented at a seminar on Applied Research on Child Health Sept 26th to 28th.
- Feyisetan BJ, Asa S, Ebigbola JA (1997). Mothers' Management of Childhood diseases in Yoruba land: The influence of cultural beliefs. *Health Transition Rev.* 7:221-234.
- Font F, González MA, Nathan R, Kimario J, Lwilla F, Ascaso C, Tanner M, Menéndez C, Alonso PL (2001). Diagnostic accuracy and case management of clinical malaria in the primary health services of a rural area in south-eastern Tanzania. *Trop. Med. Int. Health* 6:423-428.
- Goldman N, Heuveline P (2000) Health-seeking behaviour for child illness in Guatemala. *Trop. Med. Int. Health* 5(2):145-55.
- Greenwood BM, Bajang K, Whitty CJ, Target GA (2005).. Malaria. *Lancet* 365:1487-1498
- Hobbs DA, Blank SJ (1975). *Sociology and the Human Experience.* John Wiley Incorporation.
- Jhpiego (2001). Maternal and Neonatal health (MNH) Program: Birth preparedness and complication readiness. A matrix of shared responsibilities. *Maternal and Neonatal Health* 23-31.
- Kidane G, Morrow RH (2000). Teaching mothers to provide home treatment of malaria in Tigray, Ethiopia: a randomized trial. *Lancet*, 356: 550-555.
- Lubanga RG, Norman S, Ewbank D, Karamagi C (1997). Maternal Dagnosis and Treatment of Children's Fever in an Endemic Malaria Control Programme – *Acta Tropica*.
- Madubuike I (1997). Improving childhood mortality: Nigeria adopts the Integrated management of childhood illnesses. *WHO Newslett.* 12(14).
- Molyneux CS, Murira G, Masha J, Snow RW (2002). Intra-Household Relations and Treatment Decision-Making for Childhood Illness: A Kenyan Case Study. *J. Biol. Soc. Sci.* 34(1):109-131.
- Nigeria Demographic and Health Survey (2004). Calverton, Maryland: National Population Commission (Nigeria) and ORC Macro; Nigeria Demographic and Health Survey 1999.
- Nigeria Demographic and Health Survey (2003). Calverton, Maryland: National Population Commission (Nigeria) and ORC Macro; Nigeria Demographic and Health Survey 2003.
- Nnebue CC, Ifeadike GO, Nworah NG, Duru CB (2010). The Effect of Socio-demographic Status of Mother s on the Management of Febrile Illnesses Among Under-Fives that Presented in a Tertiary Hospital In South-east Nigeria: A Two-Year Review. *Afr. Med. J.* 1(1):27-30
- Nwakwuo GC, Oshonwoh FE (2013). Assessment of the level of male involvement in Safe motherhood in Southern Nigeria. *J. Community Health* 38:349-356.
- Nwankwo BO, Chukwuocha UM, Amadi AN, Esomonu OC, Dozie INS, Ikegwuoha AE, Nwabueze PO, Mbagwu SO (2009). Treatment seeking behaviour of mothers for febrile children in some rural parts of Imo State Nigeria: Implications for home management of malaria in endemic areas. *Int. J. Trop. Med.* 4(3):132-135
- Olaogun AAE, Adebayo AA, Ayandiran OE, Olasode OA (2006). Effect of mother's socio-economic status on the management of febrile conditions in their under-five children in resource limited settings. *BMC* 6(1):1-6.
- Olaogun AAE, Ayandiran OE, Olasode AO, Adebayo AA, Omokhodion F (2005) Home Management of Childhood Febrile Illnesses in a Rural Community in Nigeria. *Australia Journal of Rural Health* 13:97-107
- Osikoya KA, Sebanjo IO (2008) Fever in children: Mothers' Perceptions and their Home Managements. *Iranian J. Paediatrics* 18(3):225-229.
- Poirier MP, Davies PH, Gonzalez-dei JM, Maroe KW (2000) Paediatric Emergency Department Nurses' Perspective on fever and children. *Paediatric emergency Care* 16(1):9-12.1
- Sims SL, Boland DL, O'Neill CA (1992). Decision Making In Home Health Care. *West J. Nurs. Res.* 14(2):186-197.
- WHO (2000). World Health Report 2000 Health Systems: Improving Performance. World Health Organization. In Nnebue C.C, Ifeadike G.O, Nworah N.G, Duru C.B (2010). The Effect of Socio-demographic Status of Mother s on the Management of Febrile Illnesses Among Under-Fives that Presented in a Tertiary Hospital In South-east Nigeria: A Two-Year Review. *Afrimed. J.* 1(1): 27-30.
- Yoder PS, Hornik RC (1996). Symptoms and perceived severity of illness as predictive of treatment for diarrhea in six Asian and African sites. *Soc. Sci. Med.* 43(4):429-39.