

## Full Length Research Paper

## Acute toxicity studies, antioxidant and *in vitro* antibacterial activities of extract from the barks of *Ricinodendron heudelottii* (Euphorbiaceae)

Victor Alain Oyono<sup>1</sup>, Charles Fokunang<sup>1</sup>, Jean Paul Assam Assam<sup>2</sup>, Steve Voundi<sup>3</sup>, Peguy Tsafack<sup>3</sup>, Emmanuel Tekwu Mouafo<sup>3</sup>, Bonaventure Tchaleu Ngandjui<sup>1</sup> and Véronique Penlap Beng<sup>3\*</sup>

<sup>1</sup>Department of Pharmacognosia and Traditional Pharmacopoeia, Faculty of Medicine and Biomedical Sciences, University of Yaoundé I, Cameroon.

<sup>2</sup>Department of Biochemistry, Faculty of Science, University of Douala, Cameroon.

<sup>3</sup>Department of Biochemistry, Faculty of Science, University of Yaoundé I, Cameroon.

Received 17 February, 2014; Accepted 9 May, 2014

Many bacteria are involved in infectious diseases. Most of these bacteria become resistant to the most commonly used synthetic drugs. In Cameroon, natural substance seem to be an alternative to this problem. Thus the aim of this research was to investigate the acute toxicity, antioxidant activities and the *in vitro* antibacterial of the methanol extract of *Ricinodendron heudelottii* (Euphorbiaceae) against twelve pathogenic bacteria involved in infectious diseases. The major bioactive components were also screened. The antibacterial activity of the extract was investigated against 12 strains including 10 Gram- and 2 Gram+ bacteria by disc diffusion method and micro dilution method, followed by another agar disc diffusion for the determination of inhibition diameters, the minimum inhibitory concentration (MICs) and the minimum bactericidal concentration (MBC), respectively. 2,2-Diphényl-1-Picrylhydrazyl (DPPH) assay was used to evaluate antiradical activity. The acute toxicity study was performed according to World Health Organization (WHO) protocol. The results of the antibacterial assays indicated that the crude extract was active on 8 of 12 strains tested, with MIC ranging from 188 to 750 µg/ml and MBC from 375 to 1500 µg/ml for the extract from barks of *R. heudelottii*. Overall, the results of this study indicated that the crude extract represented a potential source of antibacterial and antiradical compounds as shown in previous studies and justified their traditional use in the treatment of bacterial infections and other diseases in Cameroon.

**Key words:** *Ricinodendron heudelottii*, antibacterial activities, bacteria, toxicity.

### INTRODUCTION

*Ricinodendron heudelottii* is a perennial native tree in the tropic and subtropic areas, reaching 40 m in height and 1.2 m of diameter, belonging to the Euphorbiaceae family

and widely distributed in the Centre, South and West regions of Cameroon (Noumi and Yomi, 2001). The barks are brown or grey color and smooth. *R. heudelottii*

\*Corresponding author. E-mail: v.penlap@yahoo.fr. Tel: (237) 77 69 54 59.

Author(s) agree that this article remain permanently open access under the terms of the [Creative Commons Attribution License 4.0 International License](http://creativecommons.org/licenses/by/4.0/)

morphology varies from one region to another by the color of the bark, the foliage and the fruits produced ; some species like those found in South region of Cameroun have deciduous leaves and big fruits, with 2 or 3 lobes in yellow green color and black color in maturity and this is different from those in other regions of Cameroon (Noumi and Yomi, 2001). The bark extract of this plant is used against cough, as poison antidote and for the treatment of intestinal diseases. Barks is also used to treat yellow fever, malaria, headache, stomach pains and some times it can help pregnant women (Momeni et al., 2005). Infusion of barks of *R. heudelotii* has been shown to possess diuretic and aphrodisiac effects. The leaves are used to treat dysentery and the fruits can be used as spices (Momeni et al., 2005). Plants possess secondary metabolites different from primary metabolites such as proteins, lipid and carbohydrate. Secondary metabolites are classified into many groups: phenolic compounds, terpenes, essential oils and nitrogenous compounds as alkaloids. Each group contains a diversity of components with biological activities (Li et al., 2007). The stem bark of *R. heudelotii* contains dinorditerpenoids for example, heudelotenol, heudelotinone as well as E-ferulic acid octacosylate and some natural chemopreventive agents (Suh et al., 1995).

Several researches demonstrated that many strains of Gram-positive and Gram-negative bacteria currently developed outstanding drug resistance, making the search of new, safe, non toxic and effective antibacterial agents to become strictly a necessity. Many antibacterial agents are available in nature for the treatment of systemic infections. Plants therefore constitute good source of active agents for this purpose and many plant extracts have been reported to possess various antimicrobial activities (Nawel et al., 2005). Although *R. heudelotii* is widely used for the mentioned biological activities, no toxicological study of the plant has been reported previously as well as its *in vitro* antibacterial and antioxidant activities in Cameroon. As a contribution to the search of non toxic, novel antibacterial principle from medicinal plants of Cameroon, results of *in vitro* antibacterial and antioxidant investigations is being reported here.

## METHODOLOGY

### Chemicals

Gentamicin usually used was acquired from a local pharmacy. Pure methanol, chloroform, nutrient agar and nutrient broth were purchased from Merck Company and other chemicals used were from Sigma Company.

### Plant

The barks of *R. heudelotii* were collected in Minlamizibi, a village in the South region of Cameroon in January, 2009. Botanical identification was made in the Cameroon National Herbarium and

the voucher specimen was deposited under the number 16610/SFR/CAM. Preparation of extract the collected plant was dried at room temperature ( $30 \pm 3^\circ\text{C}$ ), pulverized and sieved. For the extraction, the powders of 500 g each were separately macerated in pure methanol (MeOH) solvents for 72 h. The solution was filtered using Whatman filter paper N°1 and concentrated in an air circulating oven at  $54^\circ\text{C}$  until total dryness.

## Antibacterial assay

### Microorganisms

Twelve species of strains isolated from patients at "Centre Pasteur of Cameroon", a reference center for diagnostic and identified by lab technicians of Centre Pasteur were used for the evaluation of antibacterial activity. These micro-organisms were Gram-negative (*Escherichia coli*, *Salmonella typhi*, *Shigella flexneri*, *Pseudomonas aeruginosa*, *Enterobacter cloacae*, *Klebsiella pneumoniae*, *Klebsiella oxytoca*, *Morganella morganii*, *Citrobacter freundii* and *Proteus vulgaris*) and positive gram (*Staphylococcus aureus* and *Streptococcus faecalis*).

### Inocula preparation

An inoculum for each micro-organism was prepared from broth cultures containing approximately  $5.10^5$  to  $9.10^6$  colony forming units per milliliter (CFU/ml). Each diluted (1:50) inoculum was applied as a lawn with a micropipette calibrated to deliver 50  $\mu\text{l}$  containing around  $9.10^6$  CFU. The discs impregnated with extract and pure methanol only to show if there is any activity noticed using pure methanol solvent, were evaporated for 24 h at the sterile condition. The inoculated plates were incubated at  $37^\circ\text{C}$  for 24 h. Each assay in this experiment was repeated 3 times.

### Determination of the diameters of inhibition zone

The extracts was tested *in vitro* for antibacterial activity by the standard disc diffusion method against the micro-organisms at a concentration of 80 mg/ml with pure methanol. Gentamycin used as standard antibiotic (positive control) was tested at a concentration of 1  $\mu\text{g}/\mu\text{l}$ . The diameters of inhibition zones produced by these extracts and discs impregnated with pure methanol were then compared to standard antibiotic (gentamycin).

### MIC determination

For MICs determination, only the most sensitive microorganisms were tested with the extract. Serial dilutions were from 93.75 to 3000  $\mu\text{g}/\text{ml}$  of extract in the nutrient broth medium. 100  $\mu\text{l}$  of the suspension of each pathogenic bacterium ( $10^6$  cells/ml) were added and incubated at  $37^\circ\text{C}$  for 24 h. Standard antibiotic (positive control) was tested in the concentration between 2.5 to 80  $\mu\text{g}/\text{ml}$ . The lowest concentration which did not show any macroscopic growth of tested microorganism was identified as the MIC.

### Minimum bactericidal concentration (MBC) determination

For MBC determination, 100  $\mu\text{l}$  of each tube following the MIC tube was subcultured in the nutrient broth medium at  $37^\circ\text{C}$  for 24 h. The absence of any macroscopic growth of the nearest MIC tube was identified as the MBC (Mims et al., 1993).

### Antioxidant activity

Evaluation of the curd extract of *R. heudelotti* is based on the reduction of 2,2-Diphényl-1-Picrylhydrazyl (DPPH) by antioxidant component (Brand-Williams et al., 1995). 100 mM of concentrated methanolic solution of DPPH is prepared and 2 ml shared in tubes where 10 µl of ascorbic acid added. The content was mixed on a plate shaker at 300 rpm for 30 min and incubated at 25°C. Antioxidant activity was determined by absorbance at 517 nm using spectrophotometer UV-120-01 at different concentrations 1, 2, 3, 4, 5 and 6 mg/ml. Each test was repeated 3 times.

### Toxicity study

#### Experimental animals

Albino Wistar rats (100 to 200 g) of about two and half months were obtained from the animal laboratory of the Biochemistry Department of the Yaounde I University, Cameroon. All the rats were kept under environmental conditions ( $27 \pm 2^\circ\text{C}$ ) and they had free access to water and food. These rats were deprived of food but not water (16 to 18 h) prior to the administration of the extract. The principles of laboratory animal care were followed while the Department's ethical committee approved the use of the animals and the study design.

#### Acute toxicity

The bioassay was conducted according to the World Health Organisation guideline for the evaluation of safety and efficacy of herbal medicines (WHO, 2002). For the study, albino Wistar rats were divided into six groups of five animals each. The methanol extract of *R. heudelotii* was suspended in the distilled water. This extract was employed because of its moderate or high antibacterial activity against all the bacteria considered. This solution was administrated *per os* to rats groups in a single oral dose of 2, 4, 6, 8, and 10 g/kg body weight (bw), respectively by intra gastric gavages using a feeding needle. The control group received an equal volume of distilled water as vehicle. Observations of toxic symptoms were made and recorded systematically at 1, 2, 4, 6 and 24 h after administration of the extract. The number of rats that survived were noted after 24 h and then maintained for the further 8 days with daily observations. This visual observation included skin changes, mobility and aggressiveness, sensitivity to sound and pain, as well as respiratory movements. The toxic effects of the extract were assessed on the basis of mortality which was expressed as LD<sub>50</sub> (Schorderet, 1992). During the experiment, the animals were weighed, food and water intake were monitored. At the end of the experiment, all surviving animals fasted overnight and were sacrificed by decapitation. The organs such as liver, lungs, heart and kidneys were excised and weighed. The pathological observations of these tissues were performed on gross. The blood samples were also collected freshly in the dry non-heparinised centrifuge tubes. This blood was allowed to coagulate before being centrifuged to separate the serum. This serum was assayed for biochemical parameters. The liver was excised, rinsed in ice-normal solution followed by cold 0.1 M Tris-HCl (pH 7.5), blotted, dried and weighed. The 20% (w/v) liver homogenates was prepared in the 0.1 M Tris-HCl buffer and the supernatant was used for biochemical analysis. Lung, kidney and heart were removed, washed in 0.9% NaCl weighed and examined as previously mentioned. Determination of biochemical parameters blood collected into non heparinised tubes was centrifuged at 3,000 rpm for 10 min to separate the serum. This serum was used to evaluate the liver enzyme function through some biochemical parameters such as aspartate aminotransferase (AST), alanine aminotransferase (Reitman and Frankel, 1957), alkaline phosphatase

(Oliver et al., 1954), creatinine (Bartels et al., 1972), urea and total protein (Allan et al., 1949), total bilirubine and direct bilirubin (George, 1959).

### Phytochemical screening

Qualitative phytochemical tests of *R. heudelotti* methanolic extract were carried out according to Odebiyi and Sofowora (1978) methods to identify some components as alkaloids, saponins, tannins, flavonoids, polyphenols, anthraquinones.

**Test for alkaloids:** 0.5 g of the sample was stirred with 5 ml of 1% aqueous HCl on a steam bath and then filtered. 1 ml of the filtrate was treated with a few drops of Mayer's reagent and a second 1 ml portion was treated similarly with Dragendroff reagent. Turbidity or precipitation with either of these reagents was taken as evidence for the presence of alkaloids in the extract.

**Test for saponins:** The ability of saponins to produce frothing in aqueous solution and to haemolyse red blood cells was used for the screening test. 0.5 g of plant extract was shaken with water in a test tube. Frothing which persisted on warming was taken as evidence for the presence of saponins.

**Test for tannins:** 0.5 g of dried extract was stirred with 5.0 ml of distilled water. This was filtered and ferric chloric reagent was added to the filtrate. A blue-black precipitate was taken as evidence for the presence of tannins.

**Test for phenol and polyphenols:** 0.5 g of plant extract was heated for 30 min in a water bath. 3 ml of 5% FeCl<sub>2</sub> was added to the mixture then followed by the addition of 1 ml of 1.00% potassium ferrocyanide. The mixture was filtered and green (phenol) and blue (polyphenol) colours were observed.

**Test for anthraquinones:** 0.5 g of plant extract was shaken with 5 ml of benzene, filtered and 2 ml of 10% ammonia solution was added to the filtrate. The mixture was shaken and the presence of a pink or violet colour in the ammoniacal (lower) phase indicated the presence of free hydroxy anthraquinones.

**Test for flavonoids:** 0.5 g of plant extract was dissolved in 5 ml of NaOH at 1 N. The change of the yellow colour obtained after adding HCl 1 N indicated the presence of flavonoïds.

### Statistical analysis

The values were expressed as mean  $\pm$  standard deviation (SD). Each value was a mean of five or six tests. The one-way analysis of variance (ANOVA) was used to determine the significant differences between parameters and the student-Newman Keurls test served to compare these differences at  $p < 0.05$  using statistical package used was SPSS 10.1.

## RESULTS

### Antibacterial activity

The results of the *in vitro* antibacterial activity of methanol extracts determined by diameters of inhibition zones are presented in Table 1. These results indicated that the diameters of inhibition zones varied from 12 to 19.67 mm

**Table 1.** Antibacterial activities of *R. heudelotii* from the disc diffusion method.

Extract	Bacterial strains											
	<i>S.aureus</i>	<i>Kb. pneumo</i>	<i>S. feacalis</i>	<i>S. typhi</i>	<i>Kb. oxy</i>	<i>S. flexneri</i>	<i>E. coli</i>	<i>P. vulgaris</i>	<i>P. aeruginosa</i>	<i>M. morgani</i>	<i>C. freundi</i>	<i>E. cloacae</i>
Gentamicin	34.33±1.53	26±2.65	31±2	27.33±6.51	32±1	27.33±1.53	26±1	26±3.61	34.67±1.53	30.33±1.53	30.67±2.52	28±1.73
<i>R. heudelotii</i>	12±0	15.33±0.58	12.67±0.58	-	-	-	12.33±0.58	-	19.67±1.59	14.67±0.58	16±1	12.33±1.15

*S.aureus* =*Staphylococcus aureus*; *Kb. pneumo* =*Klebsiella pneumonia*; *S. feacalis* =*Streptococcus feacalis*; *S. typhi* =*Salmonella typhi*; *Kb. oxy* =*Klebsiella oxytoca*; *S. flexneri* = *Shigella flexneri*; *E. coli* =*Escherichia coli*; *P. vulgaris* =*Proteus vulgaris*; *P. aeruginosa* =*Pseudomonas aeruginosa*; *M. morgani* = *Morganella morgani*; *C. freundi* =*Citrobacter freundi*; *E. cloacae* = *Enterobacter Cloacae*.

**Table 2.** MIC, MBC values ( $\mu\text{g/ml}$ ) of *R. heudelotii* extract in the macro dilution assay comparable to gentamicin.

Extract/ gentamicin	Parameter ( $\mu\text{g/ml}$ )	Bacterial strain											
		SA	KP	AB	ST	KO	SF	EC	PV	PA	MM	CF	EC*
<i>R. heudelotii</i>	MIC	750	375	750	-	-	-	750	-	188	375	375	750
	MBC	1500	750	1500	-	-	-	1500	-	375	750	750	1500
	MBC/MIC	2	2	2	-	-	-	2	-	2	2	2	2
Gentamicin	MIC	10	10	10	20	10	10	10	10	10	5	10	10
	MBC	20	20	20	40	20	20	20	40	20	10	20	20
	MBC/MIC	2	2	2	2	2	2	2	2	2	2	2	2

SA=*Staphylococcus aureus*; KB=*Klebsiella pneumonia*; AB=*Streptococcus feacalis*; ST=*Salmonella typhi*; KO=*Klebsiella oxytoca*; SF= *Shigella flexneri*; EC=*Escherichia coli*; PV=*Proteus vulgaris*; PA=*Pseudomonas aeruginosa*; MM= *Morganella morgani*; CF=*Citrobacter freundi*; EC\*=*Enterobacter cloacae*. Minimum inhibitory concentration (MICs); minimum bactericidal concentration (MBC).

and 26 to 34.67 for the extracts and gentamicin, respectively. Gentamicin used as a standard antibiotic at the concentration of 1  $\mu\text{g}/\mu\text{l}$  exhibited higher diameters of inhibition than other extracts. No diameters of inhibition zones was obtained with discs impregnated with pure methanol. Among the twelve isolates, eight bacteria (*S. aureus*, *K. pneumonia*, *S. feacalis*, *E. coli*, *P. aeruginosa*, *M. morgani*, *C. freundi* and *E. cloacae*) were sensitive to the extract. However, *P. aeruginosa* was the most sensitive, with 19.67 mm. *S. typhi*, *K. oxytoca*, *S. flexneri* and *P. vulgaris* were the most resistant bacteria isolate with no diameters

of inhibition zones. The results of the antibacterial activity of the extract determined by minimal inhibitory concentration (MIC) and minimal bactericide concentration (MBC) are represented in Table 2. The MICs ranged between 188 and 750  $\mu\text{g/ml}$  and the MBC ranged between 375 and 1500  $\mu\text{g/ml}$ . These results revealed that *P. aeruginosa* was the most sensitive with values of 188  $\mu\text{g/ml}$  (Table 2).

The MIC of the extract of *R. heudelotii* was less active than standard antibiotic (gentamicin) ranged between 5 and 10  $\mu\text{g/ml}$ . No result of MIC and MBC was determined for *S. typhi*, *K. oxytoca*,

*S. flexneri* and *P. vulgaris* isolates.

### Antioxidant activity

2,2-Diphényl-1-Picrylhydrazyl (DPPH) radical scavenging assay revealed a high antiradical activity of the extract from barks of *R. heudelotii* with a  $\text{SC}_{50}$  of 7.21 mg/ml (Table 3).

### Acute toxicity

For the acute toxicity studies, no death of rats was

**Table 3.** Trapping percentage of the extract/ascorbic acid at different concentration.

Extracts	Concentration of extract/ascorbic acid (mg/ml)						Concentration of antioxidant 50 of extract/ascorbic acid	
	1	2	3	4	5	6	SC <sub>50</sub> (mg/ml)	R <sup>2</sup>
Ascorbic acid	36.67±0.53	40.91±0.00	50.30±1.39	55.46±0.91	60.30±0.51	63.64±0.91	3.28	0.978
<i>R. heudelotii</i>	20.30±0.53	24.85±0.53	27.88±0.53	35.46±0.91	40.00±0.00	43.64±0.91	7.21	0.988

**Table 4.** Biomedical parameters values of male rats in acute toxicity of the methanol extract from the barks of *R. heudelotii*.

Parameter tests	Extract dose (g/kg body weight)					
	0	2	4	6	8	10
Weight	124.15±10.15	116.87±20.70	117.9±21.46	102.80±15.65	109.56±18.64	124.36±13.34
Urea (mg/ml)	0.90±0.74	0.94±0.69	1.23±0.67	1.65±1.05	2.04±0.55	2.48±1.27
Creatinin (mg/ml)	1.1±0.08	1.2±0.04	1.6±0.15	2±0.47	2.01±0.13	2.45±0.22
Total protein (mg/ml)	3.77±0.53	3.93±0.51	4.55±0.55	5.5±0.71	6.1±1.19	7.35±2.31
ALP (UI/L)	10.69±0.59	11.95±0.29	13.24±1.2	14.9±0.56	16.2±1.58	18.56±1.98
AST (UI/L)	42±2.62	50.75±4.49	52.50±4.79	61.5±6.78	76±1.84	93.5±13.88
ALT (UI/L)	26±3.18	29.9±0.62	29.15±4.27	36.75±8.82	39.5±4.99	54.1±7.84
Total bilirubin (mg/l)	1.05±0.05	1.07±0.10	1.14±0.04	1.09±0.04	1.09±0.065	1.36±0.09
Conjugated bilirubin (mg/l)	0.32±0.001	0.33±0.07	0.34±0.005	0.31±0.05	0.29±0.01	0.29±0.04

neither recorded in the control nor in the treated groups. The animals exhibited slight changes in general behaviour (slow response to stimuli, stretching and sluggishness) but did not expressed changes in their physiopatological activities. Although there was an increase in the weights of rats and their organs in the treated groups, this addition remained statistically not significant compared with control. The pathological examinations of the tissues on a gross basis indicated no detectable abnormalities at the end of the experiment. Table 4 shows the results of the blood and hepatic parameters. These results indicated that alkaline phosphatase (ALP), alanine aminotransferase (ALT), aspartate aminotransferase (AST) of the treated rats

increased significantly ( $p > 0.05$ ) at the higher concentration of the extract compared to the control. We noticed also an increase of urea, total protein, creatinin (CRT) and bilirubin of the treated rats but the difference is not significant ( $p < 0.05$ ).

#### Phytochemical screening

The phytochemical screening of the extract of *R. heudelotii* indicated the presence of three main classes of compounds: polyphenols, alkaloids and steroids. The polyphenols group constitutes the principal component of *R. heudelotii* extract with three bioactives subgroups: tannins, flavonoids and saponins. Anthraquinone was absent in

the extract of *R. heudelotii*. Alkaloids and steroids whose antimicrobial activities have been demonstrated were also found in this extract (Table 5)

#### DISCUSSION

This work permitted the evaluation of some biological properties of *R. heudelotii*; among which the antimicrobial activity directed on some pathogens frequently encountered in infections as well as the toxicological and antioxidant studies of this extract. The diameters of inhibition zones varied from 12 to 19.67 mm and 26 to 34.67 for the extracts and gentamycin, respectively. The

**Table 5.** Phytochemical screening of crude methanol extract.

Chemical group	polyphenols	phenols	alkaloids	favanoids	saponins	tanins	Anthraquinone
Results	+	+	+	+	+	+	-

(diameter of inhibition between 7 and 10 mm), moderate activity (diameter of inhibition between 11 and 16 mm) and good or higher activity (diameter of inhibition between > 16 mm). All the micro-organisms (*S. aureus*, *K. pneumonia*, *S. fecalis*, *E. coli*, *P. aeruginosa*, *M. morgani*, *C. freundii* and *E. cloacae*) tested here were harmful and involved in infectious diseases. The broad antimicrobial activity of the extract in our study can be attributed to the presence of various bio-actives components such as tannins, polyphenols, alkaloids, glycosides, flavonoids, steroids and saponins found in this extract (Dinan et al., 2000). The variation observed from the diameters of inhibition zone of the bacteria tested can be attributed either to the difference of the bioactive molecules present in the extract or to their mechanism of action on Gram-positive and Gram-negative bacteria. The mechanism of action of the glycosides, polyphenols, tannins and alkaloids on Gram-positive and Gram-negative bacteria was demonstrated (Deeni and Sadiq, 2002; Kalemba, 2003). The MICs ranged between 188 and 750 µg/ml and the MBC ranged between 375 and 1500 µg/ml. These results revealed that *P. aeruginosa* was the most sensitive with values of 188 µg/ml. The MIC of the extract of *R. heudelotii* was less active than standard antibiotic (gentamycin) ranged between 5 and 10 µg/ml. The antibacterial property of *P. aeruginosa* is well documented (Kuete et al., 2007; Armelle et al., 2008). The rate MBC/MIC revealed that the extract presents a bactericidal action (MBC/MIC < 4). This action can justify the use of this plant against bacterial infections. No result of MIC and MBC was determined for *S. typhi*, *K. oxytoca*, *S. flexneri* and *P. vulgaris* whose isolates were the most resistant bacteria isolate with no diameters of inhibition zones and this can explain the result obtained.

The evaluation of trapping of DPPH radical revealed a high antiradical activity of the extract from barks of *R. heudelotii* with a SC<sub>50</sub> of 7.21 mg/ml. This antioxidant activity observed in our study can be attributed to the presence of various bio-actives components such as tannins, polyphenols, alkaloids, flavonoids and steroids found in this extract (Sokol-Letowska et al., 2007). However this trapping rate remained lower than the one of ascorbic acid (SC<sub>50</sub> = 3,28 mg/ml). The work done by Afolabi et al., in 2007 (Afolabi et al., 2007) showed a large correlation between the percentage of phenolic components of the extract and the inhibitory activity of the extract on the DPPH.

The acute toxicity test has been investigated to establish the adverse effects of the administration of the

methanol extract of *R. heudelotii* on some behavioural and biochemical parameters. The results indicated that up to 10 g/kg bw, no abnormal symptoms and no death of the rats was observed. According to the OCED protocol (Organisation of Economic Co-operation and Development, 2001) *R. heudelotii* extract can be classified as non toxic since the limited dose of an acute toxicity is generally considered to be 5.0 g/kg bw (Schorderet, 1992). If no mortality is observed at this level, a higher dosage is generally not necessary (Wallace, 1989). However, the body weight of the rats increased during the experiment. Compared to the control, the weight gained by the treated rats was higher but statistically not significant. This result showed that the extract slightly stimulates the appetite of the rats and probably not irritates directly the gastrointestinal tract.

The significant changes noted in some of the blood chemistry parameters such as ALT, AST, ALP and CRT were an indicator of some toxic effects undoubted. It should be noted that these modifications were observed at high dose (more than 6 g/kg bw). Even though the changes noted were not significant, they were statistically different compared to control. The variation of biochemical parameters indicated the malfunctioning of one or many organs. The increase of the activity of ALP after the administration of the extract may indicate the obstruction of the bile duct. However, this variation can not be attributed only to a dysfunction of the bile duct since many sources of ALP are known (liver cells; osteoblast, intestinal cells and placenta tissue). Although the variation of ALT and AST activities are associated with the hepato-cellular damage, only ALT is specific for the evaluation of liver damage. AST is highly concentrated in cardiac muscle, liver, skeletal muscle and kidneys. The significant increase of ALT and AST activities after treatment of the rats with 8 and 10 g/kg bw of extract implies an injury of the liver as well as the heart or other sources of these enzymes. The creatinine level of treated rats increased compared with the control. This last result confirms that kidneys are slightly affected at the dose of 10 g/kg body weight of the extract. The variation of other parameters such as protein and bilirubin were not statistically significant.

It has been recognized that eventual therapeutic bioactive products from plants may also contain substances which act as poisons in human (Ekramul et al., 2002). Several researches demonstrated that phenolics and polyphenols compound have antimicrobial activities. Other workers have shown that the sites and the number of hydroxyl groups on the phenol are thought to be related

to their relative toxicity to microorganisms, with the evidence that an increase of hydroxylation will result to an increased toxicity (Kareem et al., 2008). The presence of three subclasses of polyphenols (tannins, flavonoids and saponins) in the extract may have significant inhibitory effect on all isolates tested. The antibacterial effects of each of these subclasses of polyphenols on Gram-positive and Gram-negative have been demonstrated as well as the microbicidal effects of alkaloids (Kalemba and Kunicka, 2003). The presence of multiple phytochemical components confers to *R. heudelotii* extracts *in vitro* antibacterial and antioxidant activities.

## Conclusion

The methanol extract of *R. heudelotii* demonstrated effective *in vitro* antibacterial and antioxidant activities. However some toxic effects have been discovered after administration of high dose of this extract. Further research needs to be carried out to identify the active molecules and evaluate the *in vivo* antibacterial and antioxidant activities as well as sub-acute or chronic toxicities.

## ACKNOWLEDGEMENTS

The authors acknowledge the Laboratory for Tuberculosis Research of the Biotechnology Centre of Nkolbisson (University of Yaoundé I) for the important technical support and the Cameroon National Herbarium for the identification of the plant. A special vote of thanks to the Higher Institute of Health Professions staff for their support.

## Conflict of Interests

The author(s) have not declared any conflict of interests.

## REFERENCES

- Afolabi C, Akinmoladun EO, Ibukun, Dan-Ologe IA (2007). Phytochemical constituents and antioxidant properties of extracts from the leaves of *Chromolaena odorata*. *Sci. Res. Essay* 2:191-194.
- Allan GG, Charles BJ, Maxima DM (1949). Determination of serum proteins by means of biuret reaction. *J. Biol. Chem.* 177(2):751-762.
- Armelle MT, Ngameni B, Kuete V, Ingrid SK, Ambassa P, Roy R, Bezabih M, Xavier EF, Bonaventure NT, Berhanu AM, Marion MJJ, Namrita L, Veronique PB (2008). Antimicrobial activity of the crude extracts and five flavonoids from the twigs of *Dorstenia barteri* (Moraceae). *J. Ethnopharmacol.* 116:483-489.
- Bartels H, Bohmer M, Heierli C (1972). Serum creatinine determination without protein precipitation. *Clin. Chem. Acta.* 37:193-197.
- Brand-Williams W, Cuvelier ME, Berset C (1995). Use of a free radical method to evaluate antioxidant activity. *Lebensmittel-Wissenschaft und-Technologie* 28:25-30.
- Deeni Y, Sadiq N (2002). Antibacterial properties and phytochemical constituents of the leaves of African mistletoe (*Tapinanthus dodoneifolius* (DC) Danser) (*Loranthaceae*): and ethno medicinal plants of Hausaland, Northern Nigeria. *J. Ethnopharmacol.* 83:463-480
- Dinan L, Burne P, Whiting P (2000). Pytoecdysteoid profiles in seeds of *Sida* spp (Malvaceae). *Phytochem. Ann.* 12:110-119.
- Ekramul IM, Naznin KA, Ekramul HM (2002). *In vitro* antibacterial activity of the extracts and glycoside from *Sida rhombifolia* Linn. *J. Med. Sci.* 2(3):134-136.
- George EL (1959). Tissue sulfhydryl groups. *Arch Biochem. Biophys.* 82(1):70-77.
- Kalemba D, Kunicka A (2003). Antibacterial and antifungal properties of essential oils. *Curr. Med. Chem.* 10:813-829.
- Kareem SO, Akpan I, Ojo OP (2008). Antimicrobial activities of *Calotropis procera* on Selected Pathogenic Microorganisms. *Afr. J. Biomed. Res.* 11:105-110
- Kuete V, Oben EK, Gustave FN, Veronique PB, Hussain H, Krohn K, Augustin NE (2007). Antimicrobial activity of the methanolic extract and of the chemical constituents isolated from *Newbouldia laevis*. *Pharmazie* 62(7):552-556.
- Li HB, Cheng KW, Wong CC, Fan KW, Chen F, Tian Y (2007). Evaluation of antioxidant capacity and total phenolic content of different fraction of selected microalgae. *Food Chem.* 102:771-776
- Mims CA, Playfair JHL, Roi HIM, Wakelin D, Williams R (1993). Antimicrobial agents and chemotherapy. Mims Ed, *Medical Microbiology*. St Louis 35:1-35
- Momeni J, Djoulde RD, Akam MT, Kimbu SF (2005). Chemical constituents and antibacterial activities of the stem bark extracts of *Ricinodendron heudelotii*(Euphorbiaceae). *Indian J. Pharm. Sci.* 67:386-399.
- Monks R, Noel CL, Amelia HT, Farias FM, Elfrides SES, Suyenga SS, DaRocha AB, Schwartzmann G, Mothes B (2002). Anticancer, antichemotactic and antimicrobial activities of marine sponges collected off the coast of Santa Catarina, Southern Brazil. *J. Exp. Mar. Biol. Ecol.* 281:1-12.
- Nawel H, Ammar RB, Afef W, Soumaya K, Amor M, Chibani JB, Mariotte AM, Ghedria K, Dijoux-Franca MG, Ghedira LC (2005). Antibacterial and antimutagenic activity of extracts and essential oil from (Tunisia). *Pistacia lentiscus*. *Toxicol. Environ. Chem.* 87(4):567-573.
- Noumi E, Yomi A (2001). Medicinal plants used for intestinal diseases in Mbalmayo Region, Central Province, Cameroon. *Fitoterapia* 72:246-254.
- Oliver LH, Nira RR, Wu ML, Walther HS, Elizabeth CF (1954). The quantitative histo-chemistry of brain II Enzyme measurement. *J. Biol. Sci.* 207(1):19-37.
- Organisation of Economic Co-operation and Development (OECD) (2001). The OECD guideline for testing of chemical: 420 Acute Oral Toxicity. OECD, Paris, pp. 1-14.
- Reitman S, Frankel S (1957). A colorimetric method for the determination of serum glutamic oxalacetic and glutamic pyruvic transaminases. *J. Clin. Pathol.* 28(1):56-63.
- Schorderet M (1992). *Pharmacology. The fundamental concepts for therapeutics applications* Geneva, Slatkine, Paris Frison-Roche.
- Sokol-Letowska A, Oszmianski J, Wojdylo A (2007). Antioxidant activity of the phenolic compounds of Hawthorn, pine and skullcap. *Food Chem.* 103:853-859.
- Suh N, Luyengi L, Fong HHS, Kinghorn AD, Pezzuto J (1995). Discovery of natural product chemopreventive agents utilizing HL-60 cell differentiation as a model. *Anticancer Res.* 15:233-9.
- Wallace HA (1989). *Principle and Methods of Toxicology*, 2nd Edition. Raven Press, New York, NY 10036.
- WHO (2002). *Stratégie de l'OMS pour la médecine traditionnelle pour 2002-2005*. WHO/EDM/TRM/2002. Genève: OMS. P 65.